

Reclassification of *Sphaerobacter thermophilus* from the subclass *Sphaerobacteridae* in the phylum *Actinobacteria* to the class *Thermomicrobia* (emended description) in the phylum *Chloroflexi* (emended description)

Philip Hugenholtz^{1†} and Erko Stackebrandt²

Correspondence
Philip Hugenholtz
phugenholtz@lbl.gov

¹Department of Environmental Science, Policy and Management, 151 Hilgard Hall, University of California Berkeley, Berkeley, CA 94720-3110, USA

²DSMZ – Deutsche Sammlung von Mikroorganismen und Zellkulturen, Mascheroder Weg 1b, D-38124 Braunschweig, Germany

Sphaerobacter thermophilus was originally classified as the deepest branching member of the phylum *Actinobacteria* (high-G + C, Gram-positive bacteria) based on 16S rRNA gene comparative analysis. However, the analysis lacked suitable outgroups, and additional 16S rRNA gene sequences indicate that it is most closely related to *Thermomicrobium roseum*, which it also resembles phenotypically. Furthermore, both species are reproducibly affiliated with the phylum *Chloroflexi* (green non-sulfur bacteria), despite *T. roseum* currently being classified in its own phylum, the *Thermomicrobia*. Transfer of *Sphaerobacter* to the class *Thermomicrobia*, and transfer of the class *Thermomicrobia* to the phylum *Chloroflexi*, are proposed. Descriptions for the phylum *Chloroflexi* and the class *Thermomicrobia* are emended to reflect the proposed changes in classification.

Sphaerobacter thermophilus is a Gram-positive, non-spore-forming bacterium that was isolated from an aerobic thermophilic sludge (Hensel *et al.*, 1989). Comparative analysis of a partial 16S rRNA sequence obtained by reverse transcriptase sequencing (GenBank/EMBL/DDBJ accession no. X53210) suggested, at the time, that *S. thermophilus* was the most deeply branching member of the phylum *Actinobacteria* (high-G + C, Gram-positive bacteria; Demharter *et al.*, 1989). This phylum-level assignment has been disseminated through the literature (Carreto *et al.*, 1996; Embley & Stackebrandt, 1994; Monciardini *et al.*, 2003; Rainey & Stackebrandt, 1993; Rappé *et al.*, 1999; Shooner & Tyagi, 1995; Wiegel, 1998), culminating in a formal classification as a novel suborder of the *Actinobacteria* (Stackebrandt *et al.*, 1997). However, the initial analysis by Demharter *et al.* (1989) was compromised by insufficient reference sequences (outgroups) and, as more 16S rRNA gene sequences have become available, it is apparent that *S. thermophilus* is not reproducibly affiliated

with the *Actinobacteria* and is instead a member of the phylum *Chloroflexi* (green non-sulfur bacteria; Björnsson *et al.*, 2002; Hugenholtz *et al.*, 1998).

We have obtained an almost complete 16S rRNA gene sequence for the type strain of *S. thermophilus* (DSM 20745^T) using methods described previously (Rainey *et al.*, 1996) and confirm that its phylogenetic placement is within the *Chloroflexi* (Fig. 1), on the basis of evolutionary distance and Bayesian inference as previously described (Janssen & Hugenholtz, 2003). Fig. 1 shows only the *Chloroflexi* phylum sequences (ingroup) used in the analysis; they were rigorously tested for monophyly using a variety of bacterial outgroup sequences representing multiple phyla (not shown; datasets available on request), as described previously (Dalevi *et al.*, 2001).

The closest relative of *S. thermophilus* is *Thermomicrobium roseum* (Fig. 1), which it resembles phenotypically. Both are thermophilic, obligately aerobic, chemoheterotrophic, catalase-positive, non-motile, non-spore-forming, short irregular-shaped rods with similar DNA G + C content (Demharter *et al.*, 1989; Garrity & Holt, 2001b). The cell membrane of *T. roseum* contains distinctive long-chain diols instead of the familiar glycerol-based lipids (Pond *et al.*, 1986) and these have been postulated to contribute to membrane stabilization at high temperatures (Pond &

Published online ahead of print on 20 August 2004 as DOI 10.1099/ij.s.0.03028-0.

†Present address: DOE Joint Genome Institute, 2800 Mitchell Creek Bldg 400-404, Walnut Creek, CA 94598, USA.

The GenBank/EMBL/DDBJ accession number for the 16S rRNA gene sequence of *Sphaerobacter thermophilus* DSM 20745^T is AJ420142.




Fig. 1. Evolutionary distance tree of the phylum *Chloroflexi* based on comparative analysis of 16S rRNA gene sequences showing the phylogenetic affiliation of *S. thermophilus* (shown in bold). Branch points supported by evolutionary distance bootstrap resampling and Bayesian posterior probabilities are indicated by filled circles (bootstrap proportion and posterior probability values > 90%) and open circles (both values > 75%). *Escherichia coli* (X80725), *Bacillus subtilis* (D26185) and *Corynebacterium diphtheriae* (X84248) were used as outgroups (not shown). Subdivision (class)-level groupings are indicated by brackets to the right of the figure. Class names follow Garrity & Holt (2001a, b), and informally proposed classes (in quotation marks) are based on representative species names. The numbers in brackets following these names are according to the subdivision numbering in Hugenholtz *et al.* (1998). Bar, 0.05 changes per nucleotide.

Langworthy, 1987). *S. thermophilus* has not been investigated for its cell membrane composition, but given its relationship to *T. roseum* and the fact that it is a thermophile, it may also contain diol-based lipids.

T. roseum has long been recognized to be affiliated with the phylum *Chloroflexi* on the basis of 16S rRNA comparative analysis (Woese, 1987), but is currently classified in its own bacterial phylum, *Thermomicrobia* (Garrity & Holt, 2001b). We propose to transfer the class *Thermomicrobia* to the phylum *Chloroflexi* to acknowledge this phylogenetic relationship formally (Fig. 1). In addition to the class *Thermomicrobia*, at least four other major lineages exist in the phylum *Chloroflexi* (Fig. 1), three of which now have cultivated representatives: the well-characterized class *Chloroflexi*, containing the phototrophic genera *Chloroflexus*, *Chloronema*, *Heliobacter* and *Oscillochloris* and the chemoheterotrophic genus *Herpetosiphon* (Garrity & Holt, 2001a); the 'Dehalococcoidetes', containing the tetrachloroethene-respiring coccus 'Dehalococcoides ethenogenes' (Maymó-Gatell *et al.*, 1997); and, most recently, the 'Anaerolineae', containing the thermophilic filamentous bacteria *Anaerolinea thermophila* and *Caldilinea aerophila* (Sekiguchi *et al.*, 2003). The 'Anaerolineae' are not monophyletic in all analyses (note node support < 90% in Fig. 1) and may need to be subdivided as new sequences

come to hand. Detailed comparative studies of *S. thermophilus* and other members of the *Chloroflexi* may reveal unifying chemotaxonomic and ultrastructural features of this interesting phylum. For example, some members of the *Chloroflexi*, including *S. thermophilus*, have unusual cell envelopes rich in protein and lacking significant amounts of peptidoglycan (Demharter *et al.*, 1989; Maymó-Gatell *et al.*, 1997; Merkel *et al.*, 1980).

We conclude, on the basis of 16S rRNA gene sequence comparative analysis, that *S. thermophilus* should be reassigned to the class *Thermomicrobia* in the phylum *Chloroflexi*. We suggest retaining the order and family proposed for *S. thermophilus* (*Sphaerobacterales* and *Sphaerobacteraceae*, respectively; Stackebrandt *et al.*, 1997) in the class *Thermomicrobia* until a more rigorous comparison of *S. thermophilus* and *T. roseum* is performed and further representatives of this class are characterized.

Emended description of the phylum *Chloroflexi* Garrity and Holt 2001

The phylum *Chloroflexi* is a deep-branching lineage of the *Bacteria*. On the basis of comparative 16S rRNA gene sequence analysis of isolates and environmental clones, the phylum can be divided into at least five major subdivisions,

including the classes *Chloroflexi* and *Thermomicrobia*. Mostly Gram-negative, filamentous bacteria with unusual cell envelopes. No peptidoglycan or atypical peptidoglycan present in cell walls. Lipopolysaccharide-containing outer membrane not present in studied representatives.

Emended description of the class *Thermomicrobia* Garrity and Holt 2001

The class *Thermomicrobia* subdivides into two orders with validly published names: *Thermomicrobiales* Garrity and Holt 2001 and *Sphaerobacterales* Stackebrandt, Rainey and Ward-Rainey 1997. Pleomorphic, non-motile, non-spore-forming rods. Atypical proteinaceous cell walls.

Acknowledgements

We thank I. Kramer for determining the full-length 16S rRNA gene sequence of *S. thermophilus*, H. Trüper for advice on the names of higher taxa and two anonymous reviewers for helpful comments on the manuscript.

References

- Björnsson, L., Hugenholtz, P., Tyson, G. W. & Blackall, L. L. (2002). Filamentous *Chloroflexi* (green non-sulfur bacteria) are abundant in wastewater treatment processes with biological nutrient removal. *Microbiology* **148**, 2309–2318.
- Carreto, L., Moore, E., Nobre, M. F., Wait, R., Riley, P. W., Sharp, R. J. & da Costa, M. S. (1996). *Rubrobacter xylanophilus* sp. nov., a new thermophilic species isolated from a thermally polluted effluent. *Int J Syst Bacteriol* **46**, 460–465.
- Dalevi, D., Hugenholtz, P. & Blackall, L. L. (2001). A multiple-outgroup approach to resolving division-level phylogenetic relationships using 16S rDNA data. *Int J Syst Evol Microbiol* **51**, 385–391.
- Demharter, W., Hensel, R., Smida, J. & Stackebrandt, E. (1989). *Sphaerobacter thermophilus* gen. nov., sp. nov., a deeply rooting member of the Actinomycetes subdivision isolated from thermophilically treated sewage sludge. *Syst Appl Microbiol* **11**, 261–266.
- Embley, T. M. & Stackebrandt, E. (1994). The molecular phylogeny and systematics of the actinomycetes. *Annu Rev Microbiol* **48**, 257–289.
- Garrity, G. M. & Holt, J. G. (2001a). Phylum BVI. *Chloroflexi* phy. nov. In *Bergey's Manual of Systematic Bacteriology*, 2nd edn, vol. 1, *The Archaea and the Deeply Branching and Phototrophic Bacteria*, p. 427. Edited by D. R. Boone, R. W. Castenholz & G. M. Garrity. New York: Springer.
- Garrity, G. M. & Holt, J. G. (2001b). Phylum BVII. *Thermomicrobia* phy. nov. In *Bergey's Manual of Systematic Bacteriology*, 2nd edn, vol. 1, *The Archaea and the Deeply Branching and Phototrophic Bacteria*, p. 447. Edited by D. R. Boone, R. W. Castenholz & G. M. Garrity. New York: Springer.
- Hensel, R., Demharter, W. & Hilpert, R. (1989). The microflora involved in aerobic-thermophilic sludge stabilization. *Syst Appl Microbiol* **11**, 312–319.
- Hugenholtz, P., Goebel, B. M. & Pace, N. R. (1998). Impact of culture-independent studies on the emerging phylogenetic view of bacterial diversity. *J Bacteriol* **180**, 4765–4774.
- Janssen, P. H. & Hugenholtz, P. (2003). Fermentation of glycolate by a pure culture of a strictly anaerobic Gram-positive bacterium belonging to the family *Lachnospiraceae*. *Arch Microbiol* **179**, 321–328.
- Maymó-Gatell, X., Chien, Y. T., Gossett, J. M. & Zinder, S. H. (1997). Isolation of a bacterium that reductively dechlorinates tetrachloroethene to ethene. *Science* **276**, 1568–1571.
- Merkel, G. J., Durham, D. R. & Perry, J. J. (1980). The atypical cell wall composition of *Thermomicrobium roseum*. *Can J Microbiol* **26**, 556–559.
- Monciardini, P., Cavaletti, L., Schumann, P., Rohde, M. & Donadio, S. (2003). *Conexibacter woesei* gen. nov., sp. nov., a novel representative of a deep evolutionary line of descent within the class Actinobacteria. *Int J Syst Evol Microbiol* **53**, 569–576.
- Pond, J. L. & Langworthy, T. A. (1987). Effect of growth temperature on the long-chain diols and fatty acids of *Thermomicrobium roseum*. *J Bacteriol* **169**, 1328–1330.
- Pond, J. L., Langworthy, T. A. & Holzer, G. (1986). Long-chain diols: a new class of membrane lipids from a thermophilic bacterium. *Science* **231**, 1134–1136.
- Rainey, F. A. & Stackebrandt, E. (1993). Phylogenetic evidence for the classification of *Acidothermus cellulolyticus* into the subphylum of Actinomycetes. *FEMS Microbiol Lett* **108**, 27–30.
- Rainey, F. A., Ward-Rainey, N., Kroppenstedt, R. M. & Stackebrandt, E. (1996). The genus *Nocardiopsis* represents a phylogenetically coherent taxon and distinct actinomycete lineage: proposal for *Nocardiopsaceae* fam. nov. *Int J Syst Bacteriol* **46**, 1088–1092.
- Rappé, M. S., Gordon, D. A., Vergin, K. L. & Giovannoni, S. J. (1999). Phylogeny of actinobacteria small subunit (SSU) rRNA gene clones recovered from marine bacterioplankton. *Syst Appl Microbiol* **22**, 106–112.
- Sekiguchi, Y., Yamada, T., Hanada, S., Ohashi, A., Harada, H. & Kamagata, Y. (2003). *Anaerolinea thermophila* gen. nov., sp. nov. and *Caldilinea aerophila* gen. nov., sp. nov., novel filamentous thermophiles that represent a previously uncultured lineage of the domain Bacteria at the subphylum level. *Int J Syst Evol Microbiol* **53**, 1843–1851.
- Shoener, F. & Tyagi, R. D. (1995). Microbial ecology of simultaneous thermophilic microbial leaching and digestion of sewage sludge. *Can J Microbiol* **41**, 1071–1080.
- Stackebrandt, E., Rainey, F. A. & Ward-Rainey, N. L. (1997). Proposal for a new hierarchic classification system, Actinobacteria classis nov. *Int J Syst Bacteriol* **47**, 479–491.
- Wiegel, J. (1998). Anaerobic alkalithermophiles, a novel group of extremophiles. *Extremophiles* **2**, 257–267.
- Woese, C. R. (1987). Bacterial evolution. *Microbiol Rev* **51**, 221–271.