A New Species: *Microcycclus marinus*

H. D. RAJ

Department of Microbiology, California State University, Long Beach, California 90840

For a number of years, only a single species of the genus *Microcycclus* Ørskov 1928, namely *M. aquaticus* Ørskov, was reported in the literature. Eventually, two more species, *M. major* Gromov 1963 and *M. flavus* Raj 1970, were named and described. Brief mention was also made in the literature of a marine *Microcycclus* strain (Raj 1970) originally isolated in 1966 from the Pacific Ocean and tentatively, hence not validly, named *M. marinus*. To establish its taxonomic status, the marine isolate was studied extensively and compared with *M. aquaticus* ATCC 25396 and *M. flavus* ATCC 23276, both from freshwater habitats. On the basis of its similarity index, as determined by numerical taxonomy with these previously named species, the marine microcycclus is believed to belong to a new species, for which the name *Microcycclus marinus* is proposed. The type strain of *M. marinus* has been deposited in the American Type Culture Collection under the number 25205 and in the Deutsche Sammlung von Mikroorganismen Collection under the number 745.

Although ring-forming bacteria were reported by Weibel in 1887 (35) and by Migula in 1894 and 1900 (24, 25) it was not until 1928 that scientific names were proposed for these organisms. At that time, Ørskov (27) introduced the generic name *Microcycclus* and the specific name *Microcycclus aquaticus* for these organisms. After a gap of about three decades, various reports of new ring-forming isolates, mostly from soil and water habitats, were made by investigators from different countries: these included Hallock's *Vibrio alternans* (13), Gromov's *Microcycclus major* (12), Raj's *Microcycclus flavus*, and a marine microcycclus, tentatively and therefore not validly named *M. marinus* (5, 29), Claus' *Spirosoma* strains (7), Nikitin's *Renobacter vacuolatum* (26), Van Ert's gas-vacuolated *Microcycclus aquaticus* strains (34), and additional gas-vacuolated strains and a motile, flagellated variant (17; A. E. Konopka, R. L. Moore, and J. T. Staley, personal communication, 1975). Common to all of these gram-negative bacteria is, under suitable cultural conditions, their characteristic closed-ring-like morphology, as implied by the generic name. Such bacterial forms seem to be widespread in nature, as is documented by a recent scanning electron microscope study of the microbial flora in oceans (30). These "doughnut-shaped" bacteria have aroused much interest, and reports concerning their morphogenesis (29; R. G. Freedman and H. D. Raj, Abstr. Annu. Meet. Am. Soc. Microbiol. 1975, I53, p. 125), deoxyribonucleic acid (DNA) base compositions (8), DNA base-sequence homology for genetic relatedness (Konopka et al., personal

communication, 1975), gas-vesicle assembly (17), metabolic pathways and enzymes (18), and taxonomy (5, 8, 29) have appeared in the literature in recent years.

The marine microcycclus tentatively named *M. marinus* but not described by the author in 1970 (29) is described and compared extensively with the two currently recognized species *M. aquaticus* and *M. flavus* in order to establish its taxonomic status.

MATERIALS AND METHODS

Morphological and cultural studies. *M. aquaticus* American Type Culture Collection (ATCC) strain 25396 and *M. flavus* ATCC 23276 were grown and maintained as slant cultures on tryptone glucose extract agar (Difco) medium made with distilled water and fortified with 0.1% yeast extract (TGEY). A marine bacterium was originally isolated (29) from a Dendroaster (sand dollar) off the Pacific Coast in Newport Beach, Calif., during the summer of 1966, and a culture of this isolate was deposited in the ATCC under the number 25205. It was grown and maintained on modified Zobell 2216 (mZ) medium containing peptone (0.5 g), yeast extract (0.1 g), ferrous sulfate (0.02 g), agar (2.0 g), and seawater (100 ml). These three bacteria were incubated at room temperature (20 to 25°C) for growth and were maintained at refrigeration temperature (approximately 7°C), with subculturing every 3 weeks. Young cultures of these bacteria grown in their respective media, as described above, were used to study their morphology and cultural characteristics (Tables 1 and 2).

Electron microscopy. For scanning electron microscopy, exponentially grown cells of the marine isolate were harvested from an mZ agar slant and transferred directly into 3% glutaraldehyde in 10 mM N-2-hydroxyethylpiperazine-N'-2-ethanesul-

TABLE 1. *Morphological characteristics of the strains studied*^a

Characteristic	Marine isolate ^b ATCC 25205	<i>M. flavus</i> ^c ATCC 23276	<i>M. aquaticus</i> ^c ATCC 25396
1. Unicellular cells	+	+	+
2. Curved rods (vibrio-like)	⊥	⊥	+
3. Curved rods (horseshoe-like)	+	+	⊥
4. Closed ring-like cells	+	+	⊥
5. Coils or spiral forms	+	+	—
6. Maximum cell size (width < 1.0, OD ^d < 2.0, length < 5.0 μm)	+	—	+
7. Filamentous forms	+ ^e	+ ^e	—
8. Pleomorphic forms	—	—	+ ^e
9. Cells with rounded ends	+	+	+
10. Cells with pointed ends	—	—	—
11. Cells mostly single	+	+	+
12. Some cells in pairs or chains	—	—	—
13. Branching cells	—	—	—
14. Sheathed cells	—	—	—
15. Flagellated cells	—	—	—
16. Nonmotile cells	+	+	+
17. Common capsule for aggregated cells	—	+	+
18. Cells individually encapsulated	—	+	+
19. Cyst or zoogloea	—	—	—
20. Gas vacuolation	—	—	+
21. Gram-negative cells	+	+	+
22. Cells with giemsa stain	+	+	+
No. of total morphological similarities with the marine isolate		19	13
No. of positive morphological similarities with the marine isolate		11	7
No. of morphological dissimilarities with the marine isolate		3	9

^a Symbols: +, the most common morphological feature seen; ⊥, relatively less common morphological feature seen.

^b Grown on mZ agar medium at room temperature.

^c Grown on TGEY agar medium at room temperature.

^d Abbreviation: OD, outer diameter.

^e Morphological feature seen under certain cultural conditions only.

fonic acid (HEPES) buffer (pH 7.0) at 23°C. The cells were immediately collected on a 0.22-μm membrane filter (Millipore Corp.) and fixed by immersion in 1% OsO₄ in 4 mM potassium phosphate buffer containing 1 mM CaCl₂ (pH 7.0). The cells were then dehydrated by increasing concentrations of ethanol and dried by the critical-point method. Microscopy was carried out with an AMR-1000A scanning electron microscope at 20 kV with a 2-degree specimen tilt.

For transmission electron microscopy, cells of the marine isolate, harvested as above, were fixed in 3% glutaraldehyde in 10 mM HEPES buffer (pH 7.0) for 1.5 h at 0°C. The cells were then centrifuged at 7,000 × *g* for 5 min and suspended in 1% OsO₄ in the above-mentioned phosphate buffer for 4 h at 0°C. The cells were then dehydrated in increasing concentrations of ethanol and embedded in araldite. Ultrathin sections (silver to gray) were cut with a diamond knife on a Reichert ultramicrotome, post-stained with uranyl acetate and lead citrate, and micrographed with a Phillips 300 TEM.

Physiological studies. To determine the optimum growth temperature, cultures of the organisms were

inoculated in their respective media, described above, and incubated at 7, 20 to 25, 37, and 45°C. For determination of the pH which affords optimum growth, the above-mentioned media were adjusted to pH 5.0 through pH 9.0. For heat-sensitivity determinations, cultures were exposed to 55, 80, and 95°C for 10 min, and for disinfecting effect they were exposed to 70% ethanol, 3% H₂O₂, and 0.1% Zephiran chloride at intervals between 2.5 and 15 min. After each exposure, loopfuls of test culture were transported into the growth medium and incubated. For osmotic effect, the above-mentioned media with 5 to 20% sucrose were used. For growth in the presence of NaCl, TGEY broth tubes containing 0 to 20% NaCl were inoculated. The bactericidal effect of ultraviolet (UV) irradiation was determined by exposures to 25-W Hanovia 8341 at a 1-ft (30.48-cm) distance for up to 3.0 min. Various commercially available media (nonselective, selective, and chemically defined), as listed in Table 3, were tested for supporting the growth of these doughnut-shaped bacteria. For the marine isolate, the media were fortified so as to contain 3% NaCl and 0.03% K₂HPO₄. Also,

TABLE 2. *Cultural characteristics, on substrates mZ and TGEY (agar and broth), of the strains studied*

Characteristic		Marine isolate ATCC 25205 (mZ) ^a	<i>M. flavus</i> ATCC 23276 (TGEY) ^a	<i>M. aquaticus</i> ATCC 25396 (TGEY) ^a
<i>Surface colony (agar):</i>				
23. A.	Form:			
	Circular	+	+	+
24.	Irregular	—	—	—
25.	Spindle	—	—	—
26.	Filamentous	—	—	—
27.	Rhizoid	—	—	—
28. B.	Elevation:			
	Flat	—	—	—
29.	Raised	—	—	+
30.	Convex	+	+	—
31.	Pulvinate	—	—	—
32.	Concave	—	—	—
33.	Umbonate	—	—	—
34. C.	Margin:			
	Entire	+	+	+
35.	Undulate	—	—	—
36.	Lobate	—	—	—
37.	Erose	—	—	—
38.	Filamentous	—	—	—
39.	Curled	—	—	—
40.	Granular	—	—	—
41. D.	Size:			
	Punctiform	—	—	—
42.	Small (< 2 mm)	+	—	—
43.	Medium (2-5 mm)	—	+	+
44.	Large (> 5 mm)	—	—	—
45. E.	Color:			
	White	—	—	—
46.	Off-white	—	—	+
47.	Grayish	—	—	—
48.	Yellowish	—	+	—
49.	Pinkish	+	—	—
50. F.	Appearance:			
	Mucoid	+	+	+
51.	Dull	—	—	+
52.	Opaque	+	+	+
53.	Transparent	—	—	—
54.	Contoured	—	—	+
55.	Smooth	+	+	—
<i>Broth culture:</i>				
56. G.	Surface growth:			
	Ring	—	—	—
57.	Pellicle	—	—	+
58. H.	Turbidity:			
	Even	+	+	—
59.	Flocculent	—	—	+
60. I.	Sediment:			
	Flaky or granular	—	—	+
61.	Viscid	+	+	—
No. of total cultural similarities with the marine isolate			35	25
No. of positive cultural similarities with the marine isolate			8	4
No. of cultural dissimilarities with the marine isolate			4	14

^a Substrate.

TABLE 3. *Physiological characteristics of the strains studied*^a

Growth characteristic	Marine isolate ^a ATCC 25205	<i>M. flavus</i> ATCC 23276	<i>M. aquaticus</i> ATCC 25396
62. At 7°C	⊥	⊥	⊥
63. At 20–25°C	+	+	+
64. At 37 and 45°C	—	—	—
65. pH 5.0	—	⊥	—
66. pH 6.0	⊥	⊥	⊥
67. pH 7.0	+	+	+
68. pH 8.0	+	⊥	⊥
69. pH 9.0	⊥	—	⊥
70. In presence of 0% NaCl	⊥	+	+
71. In presence of 1.5% NaCl	+	⊥	+
72. In presence of 3.0% NaCl	+	⊥	⊥
73. In presence of 5.0% NaCl	+	—	—
74. In presence of 10.0% NaCl	⊥	—	—
75. In presence of 15.0% NaCl	⊥	—	—
76. In presence of 20.0% NaCl	—	—	—
77. In presence of 5.0% sucrose	—	—	—
78. In presence of 10.0% sucrose	—	—	—
79. In presence of 15.0% sucrose	—	—	—
80. In presence of 20.0% sucrose	—	—	—
81. After 10-min exposure at 55°C	+	⊥	⊥
82. After 10-min exposure at 80°C	—	—	—
83. After 10-min exposure at 95°C	—	—	—
84. After 0.5-min exposure to UV	—	+	—
85. After 1.0-min exposure to UV	—	+	—
86. After 3.0-min exposure to UV	—	—	—
87. After 2.5-min exposure to 70% ethanol	—	—	—
88. After 2.5-min exposure to 3% H ₂ O ₂	—	—	—
89. After 2.5-min exposure to 0.1% zepharine chloride	—	—	—
90. Brain heart infusion broth (Difco)	+	⊥	+
91. Trypticase soy broth (BBL)	+	⊥	+
92. Fluid thioglycolate medium (Difco)	—	—	—
93. Nutrient broth or agar (Difco)	⊥	⊥	+
94. Modified Zobell 2216 medium	+	⊥	⊥
95. Tryptone glucose extract medium (Difco)	+	+	+
96. Blood agar (Difco)	+	+	+
97. Mueller-Hinton agar (BBL)	+	+	+
98. Cystine trypticase agar with glucose (BBL)	+	+	+
99. Lowenstein-Jensen agar (BBL)	—	—	+
100. Eosin methylene blue agar (Difco)	+	—	+
101. Triple sugar iron agar (Difco): slant/stub	Yellow/yellow	Yellow/yellow	pink/no change
102. Tissue culture medium 199 or L15 (Difco)	+	+	+
103. Brock's synthetic seawater medium	+	⊥	⊥
No. of total physiological similarities with the marine isolate		25	30
No. of positive physiological similarities with the marine isolate		12	14
No. of physiological dissimilarities with the marine isolate		17	12

^a Symbols: +, Good growth; ⊥, slight growth; —, no growth even after 35 days of incubation.^b Media other than the marine medium used for the marine isolate were fortified with 3.0% NaCl and 0.03% K₂HPO₄.

various antibiotic and sulfonamide-susceptibility tests were performed as described in Table 4.

Biochemical studies. The biochemical methods used in this study were described previously (29)

except that, for the marine isolate, the media used were fortified with 3.0% NaCl and 0.03% K₂HPO₄. For detecting the utilization of various carbon sources, a basal medium having the composition of

TABLE 4. Antibiotic and sulfonamide susceptibilities of the strains studied^a

Antimicrobial agent	Marine isolate ATCC 25205	<i>M. flavus</i> ATCC 23276	<i>M. aquaticus</i> ATCC 25396
104. Ampicillin (10 µg)	+	1+	+
105. Aureomycin (30 µg) ^b	—	2+	1+
106. Bacitracin (10 units)	—	+	—
107. Chloramphenicol (30 µg)	1+	1+	—
108. Cephalothin (30 µg)	+	+	+
109. Colymycin (10 µg)	—	—	—
110. Declomycin (30 µg) ^b	—	1+	2+
111. Dihydrostreptomycin (10 µg) ^b	—	+	+
112. Elkosin (50 µg) ^b	—	+	—
113. Erythromycin (15 µg)	2+	+	+
114. Furadantin (300 µg) ^b	2+	+	+
115. Furadantin + macrodantin (300 µg)	2+	+	+
116. Gantrisin (0.25 mg)	—	2+	—
117. Kanamycin (30 µg)	—	+	1+
118. Kynex (1 µg)	—	+	—
119. Lincomycin (2 µg)	1+	—	—
120. Methicillin (5 µg)	—	—	—
121. Nalidixic acid (30 µg)	+	1+	+
122. Neomycin (30 µg) ^b	—	—	+
123. Novobiocin (30 µg) ^b	2+	1+	1+
124. Oleandomycin (15 µg)	2+	+	—
125. Oxacillin (1 µg)	—	—	—
126. Penicillin G (10 U)	—	—	—
127. Polymyxin B (50 U)	—	—	—
128. Ristocetin (30 µg) ^b	+	+	—
129. Streptomycin (5 µg)	—	+	+
130. Streptomycin (10 µg)	—	+	+
131. Sulfadiazine (1 mg)	—	—	—
132. Sulfamerazine (50 µg) ^b	—	+	—
133. Sulfamethoxy-pyridazine (50 µg) ^b	—	+	—
134. Sulfathiazole (1 mg)	+	2+	—
135. Terramycin (30 µg) ^b	1+	1+	2+
136. Tetracycline (5 µg)	—	+	2+
137. Tetracycline (30 µg)	+	1+	2+
138. Thiosulfil (50 µg) ^b	—	1+	—
139. Triple sulfa (0.25 mg)	—	—	—
140. Vibramycin (30 µg)	—	1+	2+
No. of total similarities with marine isolate		21	23
No. of positive similarities with marine isolate		13	9
No. of dissimilarities with marine isolate		16	14

^a Symbols: +, susceptible (10- to 20-mm diameter zone of inhibition); —, resistant; 1+, very susceptible (zone > 20-mm diameter) and 2+, extremely susceptible (zone > 40-mm diameter). For taxonomic purposes, 1+ and 2+ have same weight as +.

^b Difco disks used; all others were BBL disks. Mueller-Hinton agar medium (BBL) was used in all cases, except that it was fortified with 3.0% NaCl + 0.03% K₂HPO₄ for the marine isolate, and was incubated at room temperature.

Simmons citrate agar (Difco), but without citrate, was supplemented with an equal volume of 0.2% yeast extract when used for cultures of *M. aquaticus* and *M. flavus* and with an equal volume of the mZ agar medium when used for the marine isolate. In both cases, the final concentration of carbon source added was 0.1%, except for the controls, to which no carbon source was added. For the determination of the metabolism of carbohydrates by *M. aquaticus* and *M. flavus*, Hugh and Leifson (HL) medium was used in duplicate tubes: one for oxidative reaction,

the other for fermentative reaction (overlaid with a layer of sterile mineral oil) (15). For the marine isolate, modified HL medium containing peptone (0.2%), yeast extract (0.1%), NaCl (3.0%), K₂HPO₄ (0.03%), bromothymol blue (0.008%), and agar (0.4%) (pH 7.8) was used. The guanine plus cytosine (G + C) content of the DNA was determined by the buoyant density-CsCl method described by Mandel (22).

Radiorespirometric studies. The marine isolate was grown in Brock synthetic seawater medium (4)

at room temperature. The preparation of the cells and the apparatus, the analytical procedures, and the specifically labeled [^{14}C]glucoses used were essentially the same as reported previously (18, 28) except that [3- ^{14}C]glucose (specific activity, 10 mCi/mmol) obtained from the New England Nuclear Corp. was employed in this study. All radiochemical recoveries were then calculated to a total of 100% in each case of the specifically labeled carbons before the respiratory $^{14}\text{CO}_2$ recoveries were used to estimate the percent participation of each pathway involved in the metabolism of an organism. The relative contributions of the Embden-Meyerhof (EM) pathway, the Entner-Doudoroff (ED) pathway, and the pentose phosphate (PP) pathway to carbohydrate catabolism are thus calculated from the percent $^{14}\text{CO}_2$ recoveries of glucose carbons by the following set of formulas (H. D. Raj and A. H. Smith. Abstr. Annu. Meet. Am. Soc. Microbiol. 1974, p. 63, p. 155):

$$\% \text{PP} = C_1 + C_3 - C_4 - C_6$$

$$\% \text{EM} = \frac{C_3 - C_6}{C_4 - C_6} (1 - \% \text{PP})$$

$$\% \text{ED} = \frac{C_4 - C_3}{C_4 - C_6} (1 - \% \text{PP})$$

where each numeral subscript indicates the specific position of a carbon (C) atom in the substrate molecule, each C represents percent recovery as CO_2 and where $C_4 = 2 (C_{3,4}) - (C_3)$.

RESULTS

The marine isolate is described and has been compared morphologically, culturally, physiologically, and biochemically with the two recognized species *M. aquaticus* and *M. flavus* (which, unless indicated otherwise, refer to strains ATCC 25396 and ATCC 23276, respectively) as follows.

Morphological characteristics. Cells of the marine isolate are consistently nonmotile, nonflagellated, and gram negative; they occur singly and have the appearance of closed rings (0.9 to 1.5 μm outer diameter) with hollow centers (0.3 to 0.6 μm diameter) or of "horseshoes" (0.8 to 1.3 μm outer diameter and 0.3 to 0.7 μm cell width); some curved or straight rods (0.3 to 0.6 μm wide and 1 to 5 μm long) with rounded ends also occur (see Fig. 1, 2, and 3). Spiral forms and coils of 10 or more cells are common. The electron micrographs clearly show that the ends of a doughnut-shaped cell are not completely joined; sometimes they are positioned one over the other. In some cells, an indentation representing a site of cell division is seen. These cells have never been seen with pointed ends, branches, sheaths, capsules, zoogloae, cysts, spores, or gas vacuoles (Table 1). Morphologically, the

marine isolate resembles *M. flavus* since both characteristically form doughnuts, horseshoes, and coils, whereas *M. aquaticus* is more like a vibrio as it does not commonly form ring-like structures and has never been observed to make spirals or coils (29). However, the dimensions of cells of the marine isolate are about half those of *M. flavus* (doughnut- and horseshoe-like cells: 1.5 to 3.0 μm outer diameter, and 0.4 to 1.0 μm hole diameter; rod-like cells: 0.6 to 1.2 μm wide and 3.0 to 10.0 μm long) and about the same as those of *M. aquaticus* (doughnut- and horseshoe-like cells: 0.7 to 1.3 μm outer diameter, and 0.1 to 0.3 μm hole diameter; rod-like cells: 0.4 to 0.7 μm wide and 1.0 to 3.0 μm long). Unlike the marine isolate and *M. flavus*, *M. aquaticus* shows gas vacuolation; somehow this was overlooked by the author in his previous report on *M. aquaticus*, but it can be seen on close examination of the electron micrographs of this organism published earlier (28). Furthermore, *M. aquaticus* is usually nonmotile; however, a flagellated variant of this organism was recently isolated (17).

Electron micrographs of thin sections of the marine isolate (Fig. 2) show the fine structural features of procaryotic cells in general and terrestrial gram-negative bacteria in particular, especially with regard to cell wall envelope, cytoplasmic granularity, and DNA fibrils. The cell wall of the marine isolate seems to consist of three layers which run unparallel to the more nearly planar cytoplasmic membrane. Although an undulant cell wall is characteristic of most gram-negative bacteria, the extent of undulation, which is slight in this case, varies from organism to organism (36). The cytoplasmic matrix seems to be loosely and randomly packed with densely stained particles of ribonucleoprotein. The DNA is axially disposed and consists of a network of fine fibrils. Similar fine structural features were observed in *M. flavus* (29). Contrary to Ørskov's observations of *M. aquaticus*, septation for cell division is seen in the marine isolate (Fig. 1 and 2) and in *M. flavus* (29); however, it remains conjectural whether the mode of binary fission in these organisms is similar to that of other gram-negative bacteria (11). Structures similar to the blebs observed in several gram-negative bacteria (36) were also found in some electron microscope preparations of cells of the marine isolate. It seems that such structures represent cell wall evaginations due to an osmotic imbalance that results in the leakage of cytoplasmic material through "the pores" in the mucopeptide cell

FIG. 1. Scanning electron micrographs of the marine isolate (*Raj*) showing doughnuts, horseshoes, and coils when grown on mZ marine medium at RT. The cells were washed (A) or not washed (B, C, and D) prior to electron microscopy.

wall layer, thereby protruding the outer wall layer, which is rather elastic (1).

Morphological variation is seen in these organisms under different cultural conditions. For example, *M. aquaticus* shows pleomorphic forms when cultures are old, grown in the presence of an oxidizable carbohydrate, or incubated at 7°C (29). *M. flavus* and the marine isolate, on the other hand, show long filamentous forms instead of the doughnut-like cells under certain cultural conditions. *M. flavus* gives rise to long "spaghetti-like" strings when grown in MR-VP medium, nutrient gelatin medium, or HL medium with glucose (29). Further studies reveal that doughnut cells on TGEY medium transform to long filamentous cells when 0.25 to 0.5% K_2HPO_4 is added. Similarly, when K_2HPO_4 is deleted from MR-VP medium, the spaghetti cells revert to doughnut cells.

Likewise, the doughnut cells of the marine isolate on TGEY medium plus 3.0% NaCl are transformed to long sinuous forms when NaCl is deleted from the medium.

Cultural characteristics. The marine isolate produces pinkish, circular, convex, smooth, mucoid, opaque, viscid colonies of small size (<2 mm in diameter) with an entire edge and a colorless, narrow peripheral zone when grown on mZ medium. Except for the medium size and yellowish pigment, *M. flavus* colonies on TGEY medium have the same characteristics as above for the marine isolate whereas *M. aquaticus* colonies on TGEY medium are circular, raised, opaque, butyrous, off-white, and medium-sized with a "fried-egg" contour (29). Like the yellow pigment of *M. flavus*, the pinkish pigment produced by the marine isolate is insoluble in water but soluble in an acetone-ethanol mixture;

FIG. 2. Electron micrographs of thin sections of the marine isolate showing ring-like cells with indentations representing various stages of cell division.

furthermore, it is possibly carotenoid in nature. In their respective broth cultures, the marine isolate and *M. flavus* never form a ring growth around the container walls or pellicle over the medium surface but produce even turbidity and viscid sediment, whereas *M. aquaticus* gives rise to a pellicle, flocculent turbidity, and a flaky or stringy sediment.

Physiological characteristics. The three *Microcyclops* strains studied here grow optimally at room temperature (20 to 25°C), somewhat at 7°C, and not at all at 37 or 45°C. Also, they grow better with shaking than without. The pH for optimum growth of the marine isolate is around 7.6, and of *M. flavus* and *M. aquaticus* it is about 7.0. Whereas the marine isolate re-

quires 1.5 to 3.0% NaCl for growth, *M. flavus* and *M. aquaticus* can do without NaCl. However, the former can tolerate up to 15.0% NaCl whereas the latter grow only slightly in 3.0% NaCl. All three organisms failed to grow in the presence of 5% or more sucrose. When tested for thermosensitivity, the marine isolate survived 10 min of exposure to 55°C somewhat better than the other two organisms; however, all three were sensitive to the higher temperatures employed. Whereas the marine isolate and *M. aquaticus* were sensitive to UV irradiation, *M. flavus* survived 1 min of exposure but not 3 min. The three organisms were quite sensitive to 70% ethanol, 3% H₂O₂, and 0.1% Zephiran chloride. As shown in Table 3, the marine iso-

FIG. 3. Electron micrographs of thin sections of the marine isolate showing various morphogenic forms when grown on mZ marine medium at room temperature.

late grows well on TGEY medium and various other commercially available media when fortified with 3% NaCl and 0.03% K_2HPO_4 . Similarly, *M. aquaticus* is not fastidious in its growth requirements whereas *M. flavus* grows

only slightly in brain heart infusion and trypticase soy broths and not at all on eosin methylene blue and Lowenstein-Jensen agar media. The three organisms failed to grow in fluid thioglycolate medium. Some of the chemically

defined media commercially available for tissue culture procedures, such as TC medium L15 and TC medium 199 (Difco), supported luxuriant growth of these organisms without any further enrichment except for the marine isolate, which required the addition of 3.0% NaCl and 0.03% K_2HPO_4 to these media. In addition, the marine isolate grew very well in Brock synthetic seawater medium (4).

Of the 37 antibiotics and sulfonamides tested (see Table 4), the three organisms studied were susceptible to the following: ampicillin, cephalothin, erythromycin, furadantin, furadantin plus macrodantin, nalidixic acid, novobiocin, terramycin, and tetracycline (30 μ g); they were resistant to colymycin, methicillin, oxacillin, penicillin G, polymyxin B, sulfadiazine, and triple sulfa. In all, the marine isolate was susceptible to only 14 of the 37 antimicrobial agents tested, and it was extremely susceptible to erythromycin, furadantin, novobiocin, and oleandomycin. Similarly, *M. aquaticus* was susceptible to 18 agents, and it was extremely susceptible to declomycin, terramycin, tetracycline, and vibramycin. On the other hand, *M. flavus* was susceptible to 28 antimicrobial agents and was extremely susceptible to aureomycin, gantrisin, and sulfathiazole.

Biochemical characteristics. The three organisms failed to hydrolyze any of the polysaccharides tested or to produce indole, a positive methyl red reaction, acetyl-methylcarbinol, gelatin liquefaction, or H_2S . However, they produced NH_3 from peptone and possessed catalase and cytochrome oxidase (much weaker oxidase activity in *M. flavus*), but they lacked protease, reductase, urease, and hemolysin activities. Also, only *M. aquaticus* reduced NO_3 to NO_2 and NH_3 , and showed lipase activity. In litmus milk, *M. aquaticus* produced an alkaline reaction, soft curd, and peptonization; the marine isolate gave an acid reaction only; and *M. flavus* failed to cause any reaction. Of the 13 different carbon sources tested, *M. flavus* failed to oxidize any, whereas the marine isolate and *M. aquaticus* utilized acetate, citrate, malonate, pyruvate, and succinate. In addition, the marine isolate oxidized fumarate, malate, and tartrate, whereas *M. aquaticus* attacked formate, lactate, and oxaloacetate (see Table 5).

Of the 22 carbohydrates assayed (Table 5), the marine isolate oxidized 16, *M. flavus* oxidized 13, and *M. aquaticus* oxidized 12; acid but no gas was produced after 2 to 3 weeks of incubation at room temperature. Although none of the organisms attacked any of the carbohydrates fermentatively, they all oxidized arabinose, fructose, galactose, glucose, inulin, man-

nose, melibiose, sucrose, and xylose; adonitol, dulcitol, glycerol, and inositol were not oxidized.

Radiorespirometric studies. Since carbohydrate metabolism proceeds sequentially through primary and secondary pathways of catabolism, radiorespirometric experiments were conducted to estimate the contribution of each of the primary pathways operating in the marine isolate. Table 6 shows the radiochemical recoveries of specifically labeled carbon atoms of glucose. Normalizing these recoveries to a total of 100% in each case of the specifically labeled carbons (i.e., recovery of ^{14}C from CO_2 , cells, and medium totaled to 100%), it was calculated that $^{14}CO_2$ recovery of $C_1 = 53\%$, $C_2 = 46\%$, $C_3 = 67\%$, $C_{3,4} = 69\%$, and $C_6 = 42\%$; that of C_4 separately was determined to be 71%. Thus, the preferential oxidation of labeled carbons of glucose to $^{14}CO_2$ yielded a pattern: $C_4 > C_3 > C_1 > C_2 > C_6$ (Fig. 4). Since the respiratory recoveries of C_3 and C_4 are greater than those of the other carbons, the operation of the EM pathway in the marine isolate is very apparent. However, because C_4 recovery should not be greater than that of C_3 , and C_1 recovery should not exceed those of C_2 and C_6 , via this pathway, the slightly greater recovery of C_4 and C_3 and of C_1 than C_2 and C_6 seems to indicate some simultaneous participation of the PP and/or ED pathways. It is estimated, therefore, that as much as 80% of the administered glucose is oxidized via the EM pathway, 13% is oxidized via the ED pathway, and 7% is oxidized via the PP pathway. As shown in Table 6, the low yields of respiratory $^{14}CO_2$ from C_6 and C_1 as compared to those from C_4 and C_3 were compensated for by their proportionately higher cellular incorporations, and vice versa. As expected, there is a greater conservation of the methylcarbon atoms than the carboxyl carbons of pyruvates from glucose when oxidized via the tricarboxylic acid cycle for cellular biosynthesis. Contrary to the findings shown in Table 5, further radiorespirometric studies showed that the marine isolate utilized ^{14}C -labeled acetate, gluconate, and pyruvate, producing varying amounts of $^{14}CO_2$ (unpublished data). Similar studies of *M. flavus* and *M. aquaticus* reported previously elsewhere (18) showed that *M. flavus* oxidizes as much as 96% of the administered glucose via the EM pathway and only 4% via the PP pathway, whereas *M. aquaticus* catabolizes 81% of the glucose via the ED pathway and the remaining glucose via the PP and EM pathways. Both of these bacteria showed evidence of the tricarboxylic acid cycle as a secondary pathway for the oxidation of carbo-

TABLE 5. *Biochemical characteristics of the strains studied*^a

Characteristic	Marine isolate ATCC 25205	<i>M. flavus</i> ATCC 23276	<i>M. aquaticus</i> ATCC 25396
141. Agar digestion	—	—	—
142. Chitin digestion	—	—	—
143. Cellulose digestion	—	—	—
144. Gelatin liquefaction	—	—	—
145. Starch hydrolysis	—	—	—
146. Litmus milk reduction	+	—	+
147. Litmus milk acid	+	—	—
148. Litmus milk alkaline	—	—	+
149. Litmus milk hard curd	—	—	—
150. Litmus milk soft curd	—	—	+
151. Litmus milk peptonization	—	—	+
152. NO ₃ reduction to NO ₂	—	—	+
153. NO ₃ reduction to NH ₃	—	—	+
154. NO ₃ reduction to N ₂	—	—	—
155. NH ₃ from peptone water	+	+	+
156. Indole production	—	—	—
157. Methyl red reaction	—	—	—
158. Voges Proskauer reaction	—	—	—
159. H ₂ S from thiosulfate	—	—	—
160. Catalase	+	+	+
161. Oxidase	+	+	+
162. Reductase	—	—	—
163. Urease	—	—	—
164. Lipase	—	—	+
165. Hemolysins	—	—	—
166. Proteases (casein hydrolysis)	—	—	—
167. Guanine + cytosine content (mol%) of DNA base	38.3	51.0	67.3
<i>Utilization of various carbon sources:</i>			
168. Acetate	+	— ^b	+
169. Benzoate	—	—	—
170. Citrate	+	—	+
171. Formate	—	—	+
172. Fumarate	+	—	—
173. Gluconate	— ^b	— ^b	— ^b
174. Lactate	—	—	+
175. Malate	+	—	—
176. Malonate	+	—	+
177. Oxaloacetate	—	—	+
178. Pyruvate	+	— ^b	+
179. Succinate	+	—	+
180. Tartrate	+	—	—
<i>Utilization of various carbohydrates:</i>			
	(modified HL medium)	(HL medium)	(HL medium)
181. Adonitol	—	—	—
182. Arabinose	A	A	A
183. Dulcitol	—	—	—
184. Fructose	A	A	A
185. Galactose	A	A	A
186. Glucose	A	A	A
187. Glycerol	—	—	—
188. Inulin	A	A	A
189. Inositol	—	—	—
190. Lactose	A	A	—
191. Maltose	A	A	—
192. Mannitol	—	—	A
193. Mannose	A	A	A

Table 5—Continued

Characteristic	Marine isolate ATCC 25205	<i>M. flavus</i> ATCC 23276	<i>M. aquaticus</i> ATCC 25396
194. Melezitose	A	—	—
195. Melibiose	A	A	A
196. Raffinose	A	A	—
197. Rhamnose	A	—	—
198. Salicin	A	—	A
199. Sorbitol	—	—	A
200. Sucrose	A	A	A
201. Trehlose	A	A	—
202. Xylose	A	A	A
No. of total biochemical similarities with the marine isolate		49	40
No. of positive biochemical similarities with the marine isolate		16	19
No. of biochemical dissimilarities with the marine isolate		13	22

^a Symbols: A, definite acid produced oxidatively in about 2 to 3 weeks; +, positive reaction; —, no reaction even after 60 days of incubation. Duplicate tubes overlaid with sterile mineral oil for fermentative reaction showed no reaction.

^b Radiorespirometric studies showed that all the three organisms utilized ¹⁴C-labeled acetate, gluconate, and pyruvate individually in their respective growth media producing varying amounts of ¹⁴CO₂ (18; unpublished data).

FIG. 4. Aerobic radiorespirometric patterns of glucose catabolism by the marine isolate. Arrow indicates a relative time unit of 9 h required by the organism to consume all of the intact labeled substrate originally added.

TABLE 6. Utilization of [^{14}C]glucose by growing cells of the marine isolate

Substrate	Radiochemical inventory ^a			
	CO ₂	Cells	Medium	Total ^{14}C recovery
[1- ^{14}C]glucose	51	28	17	96
[2- ^{14}C]glucose	45	33	20	98
[3- ^{14}C]glucose	66	16	17	99
[3,4- ^{14}C]glucose	68	17	13	98
[6- ^{14}C]glucose	42	35	22	99

^a Determined at end of experiment. All radiochemical recoveries (CO₂, cells, and medium) were first calculated to a total of 100% in each case. Then, the normalized $^{14}\text{CO}_2$ recoveries were used to estimate the relative percent contribution of each catabolic pathway, as per the formulas shown in the text.

hydrates. Furthermore, both catabolized ^{14}C -labeled acetate, gluconate, and pyruvate, yielding respiratory $^{14}\text{CO}_2$.

DISCUSSION

In this study, the marine microcycus was extensively compared with strains of the currently recognized *Microcycus* species (Tables 1 to 5). A total of 202 characteristics was recorded for each organism: 10% of these were morphological, 20% were cultural, 40% were physiological, and 30% were biochemical features. On the basis of the characteristics shared under each category, the marine isolate was found to be similar 87 and 59% morphologically, 90 and 70% culturally, 60 and 70% physiologically, and 80 and 66% biochemically to *M. flavus* ATCC 23276 and *M. aquaticus* ATCC 25396, respectively. In all, the marine strain shared a total of 149 (60 positive and 89 negative) features with *M. flavus* ATCC 23276 and 131 (53 positive and 78 negative) features with *M. aquaticus* ATCC 25396. This accounts for the marine isolate having, at best, 74% cross-relation with *M. flavus* and 65% with *M. aquaticus*. Applying the Adansonian principles of numerical taxonomy and using Sneath's equation (31) for the similarity index (S-value) on the basis of only positive features shared by the marine isolate, the S-value with *M. flavus* is calculated to be about 53%; with *M. aquaticus* it is about 43%. Since these percent similarities are far below the minimal values expected of strains within the same species (20), it seems that the marine isolate does not belong to either of the two currently recognized *Microcycus* species. However, an S-value of 43 to 53% seems significant enough to justify separate-species status for the marine *Microcycus*. It was on the basis of the

same considerations that *M. flavus*, which, in agreement with a previous report (28), shows here only a 67% cross-relation having a 46% S-value with the type species, *M. aquaticus*, was proposed and recognized as a separate species (5, 29).

Thus, the marine isolate is here regarded as belonging to a new species, for which the name *Microcycus marinus* is proposed (ma.ri'nus. L. adj. *marinus* of the sea, marine). (It should be noted here that the marine isolate was tentatively named *Microcycus marinus* by the author in 1970 [29]; however, the name was not validly published at that time, and it therefore has no standing in nomenclature. The present proposal of the name *M. marinus* does meet the current requirements for the valid publication of a species name as given in the *Bacteriological Code* [19].) The type strain of this species has been deposited in the American Type Culture Collection, Rockville, Md., under the number 25205 and in the Deutsche Sammlung von Mikroorganismen Collection under the number 745. The description given above for the marine isolate serves as the description for both the species *M. marinus* and the type strain. Obviously, the species description will have to be expanded as additional strains of this organism are isolated and described.

The characters useful in differentiating *M. marinus* from *M. aquaticus* and *M. flavus* are given in Table 7. Similarly, *M. marinus* is readily differentiated from *M. major* by morphological, cultural, physiological, and biochemical characteristics; for example, compared to the small size of *M. marinus* (doughnut cells having 0.8 to 1.5 μm outer diameter and 0.3 to 0.7 μm width), *M. major* ring-like cells have a 5- to 10- μm outer diameter and 1- to 2- μm width. Also, whereas the former occurs occasionally as straight or curved rods 1 to 5 μm long, the latter shows the same forms 5 to 50 μm or more in length. Moreover, unlike *M. marinus*, *M. major* grows in some media as slightly bent rods with pointed ends and with age produces fiber-shaped (filamentous) structures several tens of micrometers in length and other bizarre shapes and involuted forms (7). Culturally, *M. marinus* grows on a seawater agar medium producing smooth, circular, mucoid, opaque, viscid, pink-pigmented, small-sized (<2 mm in diameter), convex colonies, whereas *M. major* grows on a freshwater agar medium giving rise to pale rose, round, shiny, transparent, large-sized (up to 0.5 cm in diameter), convex colonies. Also, unlike *M. marinus*, an older inoculum of *M. major* forms smaller (0.2 cm in diameter), dull, and orange-colored

TABLE 7. Characters useful in differentiating among the *Microcycilus* species studied

Character	<i>M. marinus</i>	<i>M. flavus</i>	<i>M. aquaticus</i>
Morpho- logical	Mostly rings and horseshoes (0.8–1.5 μm OD ^a); some coil or spiral forms	Mostly rings and horseshoes (1.5–3.0 μm OD); some coil or spiral forms	Mostly vibrioid cells (0.4–0.7 by 1.0–3.0 μm); rings not common. Coils, spirals, and long strings not seen
	Long sinuous strings in NaCl-deficient media	Long "spaghetti" strings in K ₂ HPO ₄ media	Gas vacuoles produced by all known strains; usually nonmotile but a flagellated variant has recently been found
	No gas vacuoles, flagella, or motility	No gas vacuoles, flagella, or motility	
Cultural	Small (2 mm in diameter), pinkish, mucoid, opaque, smooth, and convex colonies	Medium-sized (2–5 mm in diameter), yellowish, mucoid, opaque, smooth, and convex colonies	Medium-sized (2–5 mm in diameter), off-white, dull, mucoid, opaque, raised, and contoured colonies
	Even turbidity and viscid sediment but no pellicle in broth cultures	Even turbidity and viscid sediment but no pellicle in broth cultures	Pellicle, flocculent turbidity, and flaky sediment in broth cultures
Physio- logical	Seawater or NaCl required; can tolerate up to 15% NaCl. Susceptible to lincomycin and resistant to aueromycin, declomycin, dihydrostreptomycin, kanamycin, streptomycin, tetracycline, and vibramycin	NaCl not required for growth; can tolerate up to 3% NaCl. Susceptible to bacitracin, elkosin, gantrisin, sulfamerazine, sulfamethoxypyridazine, and thiosulfil	NaCl not required for growth; can tolerate up to 3% NaCl; susceptible to neomycin and resistant to chloramphenicol, oleandomycin, ristocetin, and sulfathiozole
Meta- bolic	Litmus milk, lipids, and NO ₃ not attacked	Litmus milk, lipids, and NO ₃ not attacked	Litmus milk rendered alkaline and peptonized; NO ₃ reduced to NO ₂ or NH ₃ ; Lipase present
	Malate or tartrate but not formate, lactate, or oxaloacetate utilized as sole carbon source	Formate, lactate, malate, oxaloacetate and tartrate not utilized as sole carbon source	Formate, lactate, and oxaloacetate but not malate or tartrate utilized as sole carbon source
	Oxidatively, acid produced from lactose, maltose, melezitose, raffinose, rhamnose, and salicin but not from mannitol or sorbitol	Oxidatively, acid produced from lactose, maltose, and raffinose but not from melezitose, mannitol, salicin, rhamnose, or sorbitol	Oxidatively, acid produced from mannitol, salicin, and sorbitol but not from lactose, maltose, melezitose, raffinose, or rhamnose
	Glucose catabolized mainly via the EM-tricarboxylic acid pathways with some via the PP and ED pathways	Glucose catabolized mainly via the EM-tricarboxylic acid pathways with a little via the PP pathway and none via the ED pathway	Glucose catabolized mainly via the ED-tricarboxylic acid pathways with some via the PP and EM pathways
G + C content	38.3 mol%	51.0 mol%	67.3 mol%

^a OD, Outer diameter.

colonies with wrinkled surface and wavy edge. In a seawater broth medium, *M. marinus* produces even turbidity and viscid sediment but never ring-growth around the container walls or pellicle over the broth surface. *M. major*, on the other hand, in a freshwater broth medium yields cottony-growth on the broth surface which easily sinks to the bottom and forms a flocculent precipitate (12). Although physiological as well as metabolic studies made of *M. major* to date are quite inadequate (12), it seems that, unlike *M. marinus*, the former hydrolyzes starch, reduces methylene blue, liquefies gelatin slightly, and fails to attack litmus milk, rhamnose, or xylose. Also, no thorough taxonomic studies have yet been made of *M. major*. Even though both *M. marinus* and *M. major* have been shown (8) to have similar G + C contents (38.3 to 39.7 mol%), *M. major*, because of its large size and pointed ends, its spiral forms, and other characters noted above, has been thought to be a nonmotile variant of a *Spirillum* sp. or possibly closely related to some blue-green algae which characteristically form spiral structures (7). Also, it has been suggested that this organism should be considered to be a flavobacterium (7).

It may be noted that the DNA base compositions of these organisms vary significantly (see Table 7). Using this as a sole parameter of genetic similarity, Claus et al. (8) proposed classifying *M. aquaticus*, *M. flavus* and *Spirosoma*, and *M. major* into three separate genera and suggested that ring formation (i.e., cell morphology) is not a sufficiently restrictive criterion to justify delineation of a genus. If, indeed, the characteristic morphology is overlooked and the DNA base composition is the only taxonomic determinant, then these bacteria could very well be included in the genus *Desulfovibrio*, *Flavobacterium*, *Pseudomonas*, *Spirillum*, or *Vibrio* (14). By the same token, if both gross morphology and DNA base composition were the only criteria for bacterial classification, then these *Microcycylus* organisms would belong to the genus *Spirillum* or *Vibrio* (10). The variation noted in the G + C contents of these *Microcycylus* species is not uncommon among species within the same genus and even among strains of the same species. For example, the DNA base composition of 28 strains of *Desulfovibrio desulfuricans* is shown to vary from 44 to 65 mol% (14).

DNA base homology has been used in the molecular taxonomy of a number of microbial groups (21). Recently, Konopka et al. (personal communication, 1975) studied the DNA heteroduplex formation as well as the G + C content of *Microcycylus* species and other ring-forming

bacteria. They found that the marine isolate (*M. marinus*) and *M. flavus* are genetically unrelated to *M. aquaticus* (duplex formation of 0 to 4% at 73°C) and, therefore, they proposed that these three bacteria should represent three different genera. Furthermore, these authors identified, mostly on the basis of G + C content, five gas-vacuolated isolates as belonging to the type species, *M. aquaticus*, and two *Spirosoma* isolates as belonging to *M. flavus*, even though the DNA base homology failed to show significant genetic relatedness expected of strains of one species; for example, under their experimental conditions, the heteroduplex formation was only 28 to 45% among the former group and 24 to 38% among the latter. In contrast, similar studies of 30 strains of *Escherichia coli* yielded a base-sequence complementarity of 78 to 100% (3). However, the degree of interspecific DNA reassociation and the stability of the heteroduplexes can vary greatly with different experimental conditions, such as differences in binding temperature and ionic strength (2, 3, 6). Another approach to classification based on genetic similarity and evolutionary relationships is the comparison of homologous proteins of closely related organisms, according to which it is expected that, among members of any one species, particular enzymes would display similar electrophoretic behavior; nevertheless, variations have been noted in the electrophoretic mobilities of the lactic dehydrogenases of otherwise closely related organisms (23). Similarly, various strains of *Leucothrix mucor* that were quite similar in their morphological, nutritional, and physiological characteristics as well as in DNA base composition were found to contain malate and isocitrate dehydrogenases with heterogeneous electrophoretic properties (16). There are several ways in which electrophoretic differences may not be due to changes in the genetic material (23); however, it is possible that a single mutational event causing a change in the amino acid composition can alter the electrophoretic mobility of a homologous protein.

It is reasonable to presume that the interactions observed at the molecular level, such as DNA reassociations, reflect genetic and phylogenetic relationships among related organisms. However, in the absence of any formal guidelines, any attempt to derive taxonomic conclusions from genetic and molecular data alone is considered premature and certainly arbitrary (2, 6). By the same token, it is realized that basing the relatedness on phenetic data alone suffers from certain taxonomic limitations (33). Sneath and Sokal (32) suggested the use of the term "phenon" per se in the numerical taxon-

omy of bacteria without necessarily equating phenons with species, genus, etc. Cowan (9), in fact, questioned the validity of the species concept in bacteria. However, studies of polyphasic taxonomy complementing numerical taxonomy with polynucleotide sequence relationships have been favored by some workers (2, 3, 6). Although the polyphasic approach has shown a high correlation between the two methods in most cases, it has failed to resolve a few baffling sets of data: for example, the case of *E. coli* 0128a versus *Shigella flexneri* 24570, with an S-value of only 59% but a DNA reassociation relationship of 80% at 75°C and, conversely, the case of *E. coli* 0128a versus *Salmonella typhimurium* LT2, with an S-value of 71.3% but a DNA duplex formation of only 12% at 75°C (3); similarly, the case of *Vibrio cholerae* versus *V. parahaemolyticus*, with interspecies S-values of 66 to 75%, G + C contents of 46 to 48 mol%, but a relative DNA reassociation of only 0 to 5% at 75°C. Although the low levels of genetic relatedness between the two *Vibrio* species apparently justify separate genus status for each, the retention of *V. parahaemolyticus* within the genus *Vibrio* was recommended as a cautious and more practical assessment of all the data on hand, and additional research was suggested (6). Similarly, although separate genus status could be justified for *M. marinus*, it seems more pragmatic at this time to recognize it as a separate species within the genus *Microcycylus*. This would be consistent with the recent judgement giving separate species status to *M. aquaticus*, *M. flavus*, and *M. major* (5). Clearly, an extensive comparative study, phenetic as well as genetic, is needed of all *Microcycylus* and related ring-forming organisms to better establish their taxonomic status.

ACKNOWLEDGMENTS

I thank Richard L. Weiss, The Biological Laboratories, Harvard University, Cambridge, Mass., for doing the electron microscopy of *M. marinus* and M. Mandel, Department of Molecular Biology, M. D. Anderson Hospital and Tumor Institute, University of Texas, Houston, Tex., for determining the DNA base compositions.

This research was supported by institutional faculty research grant no. 212-138 from the National Science Foundation.

LITERATURE CITED

1. Bayer, M. E. 1967. Response of cell walls of *Escherichia coli* to a sudden reduction of the environmental osmotic pressure. *J. Bacteriol.* 73:365-375.
2. Brenner, D. J., G. R. Fanning, K. E. Johnson, R. V. Citarella, and S. Falkow. 1969. Polynucleotide sequence relationships among members of *Enterobacteriaceae*. *J. Bacteriol.* 98:637-650.
3. Brenner, D. J., G. R. Fanning, F. J. Skerman, and S. Falkow. 1972. Polynucleotide sequence divergence among strains of *Escherichia coli* and closely related organisms. *J. Bacteriol.* 109:953-965.
4. Brock, T. D. 1964. Knots in *Leucothrix mucor*. *Science* 144:870-872.
5. Buchanan, R. E., and N. E. Gibbons. 1974. *Bergey's manual of determinative bacteriology*, 8th ed. The Williams and Wilkins Co., Baltimore.
6. Citarella, R. V., and R. R. Colwell. 1970. Polyphasic taxonomy of the genus *Vibrio*: polynucleotide sequence relationships among selected *Vibrio* species. *J. Bacteriol.* 104:434-442.
7. Claus, D. 1967. Taxonomy of some highly pleomorphic bacteria. Spisy Priorodoved. Fak. Univ. J. E. Purkyne Brno K40:254-257.
8. Claus, D., J. E. Bergendahl, and M. Mandel. 1968. DNA base composition of *Microcycylus* species and organisms of similar morphology. *Arch. Mikrobiol.* 63:26-28.
9. Cowan, S. T. 1962. The species—a macromyth. Microbial classification. Symp. Soc. Gen. Microbiol. 12:433-455.
10. De Ley, J. 1964. *Pseudomonas* and related genera. *Annu. Rev. Microbiol.* 18:17-46.
11. Gilleland, H. E., Jr., and R. G. E. Murray. 1975. Demonstration of cell division by septation in a variety of gram-negative rods. *J. Bacteriol.* 121:721-725.
12. Gromov, B. V. 1963. A new bacterium of the genus *Microcycylus*. Dokl. Akad. Nauk. SSSR 152:733-734.
13. Hallock, F. A. 1960. The life cycle of *Vibrio alternans* (sp. nov.). *Trans. Am. Microsc. Soc.* 79:404-411.
14. Hill, L. R. 1966. An index to deoxyribonucleic acid base composition of bacterial species. *J. Gen. Microbiol.* 44:419-437.
15. Hugh, R., and E. Leifson. 1953. The taxonomic significance of fermentative versus oxidative metabolism of carbohydrates by various gram-negative bacteria. *J. Bacteriol.* 66:24-26.
16. Kelly, M. T., and T. D. Brock. 1969. Molecular heterogeneity of isolates of marine bacterium *Leucothrix mucor*. *J. Bacteriol.* 100:14-21.
17. Konopka, A. E., J. T. Staley, and J. C. Lara. 1975. Gas vesicle assembly in *Microcycylus aquaticus*. *J. Bacteriol.* 122:1301-1309.
18. Kottel, R. H., and H. D. Raj. 1973. Pathways of carbohydrate metabolism in *Microcycylus* species. *J. Bacteriol.* 113:341-349.
19. Lapage, S. P., P. H. A. Sneath, E. F. Lessel, V. B. D. Skerman, H. P. R. Seeliger, and W. A. Clark (ed.) 1975. *Bacteriological code*. American Society for Microbiology, Washington, D.C.
20. Liston, J., W. J. Wiebe, and R. R. Colwell. 1963. Quantitative approach to the study of bacterial species. *J. Bacteriol.* 85:1061-1070.
21. McCarthy, B. J. 1967. Arrangement of base sequences in deoxyribonucleic acid. *Bacteriol. Rev.* 31:215-229.
22. Mandel, M. 1966. Deoxyribonucleic acid base composition in the genus *Pseudomonas*. *J. Gen. Microbiol.* 43:273-292.
23. Markert, C. L. 1968. Molecular basis for isozymes. *Ann. N. Y. Acad. Sci.* 151:14-40.
24. Migula, W. 1894. Über ein neues system der bakterien. *Arb. Bakteriolog. Inst. Karlsruhe* 1:235-238.
25. Migula, W. 1900. *System der Bakterien. Handbuch der Morphologie Entwicklungsgeschichte und Systematik der Bakterien* 2:955-960.
26. Nikitin, D. I. 1971. A new soil microorganism—*Renobacter vacuolatum*. Gen. et sp. n. Dokl. Akad. Nauk. SSSR 198:447-448.
27. Ørskov, J. 1928. Beschreibung eines neuen mikroben, *Microcycylus aquaticus*, mit eigentümlicher morphologie. *Zentralbl. Bakteriolog. Parasitenk. Infektionskr. Hyg. Abt. 1* 107:180-184.
28. Raj, H. D. 1967. Radiorespirometric studies of *Leucothrix mucor*. *J. Bacteriol.* 94:615-623.
29. Raj, H. D. 1970. A new species: *Microcycylus flavus*. *Int. J. Syst. Bacteriol.* 20:61-81.

30. Sieburth, J. M., H. L. Pratt, P. W. Johnson, and D. Scales. 1975. Microbial seascapes—a pictorial essay of marine microorganisms and their environments. University Park Press, Baltimore.
31. Sneath, P. H. A. 1957. The application of computers to taxonomy. *J. Gen. Microbiol.* 17:201–226.
32. Sneath, P. H. A., and R. R. Sokal. 1962. Numerical taxonomy. *Nature (London)* 193:855–860.
33. Sokal, R. R., and P. H. A. Sneath. 1963. Principles of numerical taxonomy. W. H. Freeman and Co., San Francisco.
34. Van Ert, M., and J. T. Staley. 1971. Gas vacuolated strains of *Microcycylus aquaticus*. *J. Bacteriol.* 108:236–240.
35. Weibel, E. 1887. Untersuchungen über vibrionen. *Zentralbl. Bakteriol. Parasitenk. Infektionskr. Hyg. Abt. 1* 16:465–472.
36. Wiebe, W. J., and G. B. Chapman. 1968. Fine structure of selected marine pseudomonads and achromobacters. *J. Bacteriol.* 95:1862–1873.