DNA Relatedness among the Pathovar Strains of *Pseudomonas syringae* subsp. *savastanoi* Janse (1982) and Proposal of *Pseudomonas savastanoi* sp. nov.

L. GARDAN,^{1*} C. BOLLET,² M. ABU GHORRAH,^{1†} F. GRIMONT,³ AND P. A. D. GRIMONT³

Institut National de la Recherche Agronomique, Station de Pathologie Végétale, F-49070 Beaucazoué,¹

Laboratoire de Bactériologie et d'Hygiène Hospitalière, Hôpital Salvator, F-13009 Marseille,²

and Unité des Entérobactéries, Institut National de la Santé et de la Recherche Médicale U199, Institut Pasteur, F-75724 Paris Cedex 15,³ France

We found that *Pseudomonas syringae* subsp. *savastanoi* strains belong to a DNA relatedness group that includes strains of *P. syringae* pv. *glycinea* and *P. syringae* pv. *phaseolicola*. This DNA group was distinct from *P. syringae* pv. *syringae* (including the type strain of *P. syringae*). The results of a numerical analysis were in accord with DNA hybridization data. Thus, *P. syringae* subsp. *savastanoi* (Janse) 1982 is elevated to species level as *Pseudomonas savastanoi* sp. nov., which includes *P. savastanoi* pv. *savastanoi*, *P. savastanoi* pv. *glycinea*, and *P. savastanoi* pv. *phaseolicola*.

In 1908, Smith (22) named "*Bacterium savastanoi*," the bacterium which causes knots on several plants belonging to the family Oleaceae. This species was later transferred to the genus *Pseudomonas* as "*Pseudomonas savastanoi*" by Stevens (24).

In 1978, Young et al. (27) proposed a new nomenclature and classification for plant-pathogenic bacteria and introduced the concept of pathovar, and all fluorescent oxidase-negative *Pseudomonas* species (except *Pseudomonas viridiflava*) were considered to be members of a single species, *Pseudomonas syringae*, which had a number of pathovars. Thus, "*P. savastanoi*" became *P. syringae* pv. *savastanoi* (27). The pathovar is an infrasubspecific subdivision which is not covered by the *International Code of Nomenclature of Bacteria* (14). Thus, the name *P. syringae* pv. *savastanoi* did not appear on the Approved Lists of Bacterial Names (21).

"*Bacterium savastanoi* var. *fraxini*" (2), which was isolated from excrescences on *Fraxinus excelsior* L., and "*Bacterium tonellianum*" (6), which was isolated by Ferraris from galls on *Nerium oleander* L. and which became "*P. savastanoi* var. *nerii*" (21), were included in *P. syringae* pv. *savastanoi* by Young et al. (27).

In 1982, Janse revived the epithet *savastanoi* in designating a new subspecies, *Pseudomonas syringae* subsp. *savastanoi* (17). The following three pathovars were recognized in this subspecies: *P. syringae* subsp. *savastanoi* pv. *oleae*, which causes parenchymatic galls on members of various genera of the Oleaceae; *P. syringae* subsp. *savastanoi* pv. *nerii*, which causes parenchymatic galls or wartlike excrescences on *N. oleander* L. and members of various genera of the Oleaceae; and *P. syringae* subsp. *savastanoi* pv. *fraxini*, which causes wartlike excrescences on *F. excelsior* L. and *Olea europea* L. (12). Although Janse did not specifically describe the taxon, his proposal in fact created *P. syringae* subsp. *syringae* (14).

Several authors have described physiological, nutritional,

and biochemical characteristics of *P. syringae* subsp. *savastanoi* strains isolated from different hosts and compared these organisms with some pathovars of *P. syringae* subsp. *syringae* (10, 11, 16, 25).

In a numerical taxonomy study, 34 strains of *P. syringae* subsp. *savastanoi* constituted a phenon that was closely related to *P. syringae* pv. *glycinea* and *P. syringae* pv. *phaseolicola* (20).

Previously, DNA-DNA hybridization data have shown that *P. syringae* is a heterogeneous species (18).

The purpose of this work was to determine the taxonomic position of *P. syringae* subsp. *savastanoi* and related pathovars by using numerical taxonomy and DNA-DNA hybridization. Our results indicated that *P. syringae* subsp. *savastanoi* should be elevated to species level as *Pseudomonas savastanoi* sp. nov.

MATERIALS AND METHODS

Bacterial strains. Three sets of strains were used. The first set comprised 143 *P. syringae* subsp. *savastanoi* strains that were isolated from various hosts (Table 1). The second set included 50 reference strains of *P. syringae* pathovars and the type strains of *Pseudomonas cichorii*, *P. viridiflava*, and *Pseudomonas amygdali*; the origins of these strains have been published previously (7). The third set contained 35 strains of miscellaneous *Pseudomonas* spp. (Table 2), *Agrobacterium tumefaciens* CFBP 2413^T (T = type strain), and *Xanthomonas campestris* pv. *campestris* CFBP 2350. All of the bacteria were cultured routinely on YBGA (yeast extract, 7 g; Bacto Peptone, 7 g; glucose, 7 g; agar, 15 g; distilled water, 1,000 ml; pH 7.2) incubated at 25°C.

Biochemical and physiological tests. The presence of oxidase, gelatinase, and arginine dihydrolase, fluorescent pigment production, levan formation, acid production from sucrose, sorbitol, erythritol, and mannitol, reduction of nitrate, hydrolysis of Tween 80 and esculin, and the hypersensitivity reaction on tobacco leaves were tested as described by Lelliott et al. (15). Pectolytic activity was tested by using the method of Prunier and Kaiser (19) and on

* Corresponding author.

† Present address: Department of Plant Protection, Faculty of Agronomy, University of Damas, Damas, Syria.

TABLE 1. Origins of the 143 *P. syringae* subsp. *savastanoi* strains tested

No. of strains	Host	Geographic origin (no. of strains)
58	Olive	Algeria (31), Tunisia (1), France (7), Portugal (1), United States (2), Yugoslavia (4), Greece (1), Italy (7), Syria (4)
33	Oleander	France (13), Algeria (4), Greece (5), Yugoslavia (1), Netherlands (1), United States (4), Italy (5)
39	Ash	France (25), Algeria (2), United Kingdom (2), United States (1), Netherlands (9)
6	Privet	Italy (6)
4	<i>Phillyrea</i> sp.	Algeria (4)
3	Jasmine	Greece (3)

pectate gels at pH 5 and 8 (9). Utilization of DL-lactate, L-(+)-tartrate, and D-(-)-tartrate was tested on a basal medium (NH₄H₂PO₄, 1 g; KCl, 0.2 g; MgSO₄ · 2H₂O, 0.2 g; agar, 3 g; bromothymol blue, 0.08 g; distilled water, 1,000 ml; pH 7.2) supplemented with 0.1% (wt/vol) organic acid

(sodium salt). The presence of DNase was tested on DNA agar (Diagnostics Pasteur, Marnes-la-Coquette, France).

Assimilation of 49 carbohydrates, 49 organic acids, and 49 amino acids was studied by using API 50CH, API 50AO, and API 50AA strips (BioMérieux, La Balme-les-Grottes, France) which were incubated at 24°C and examined for growth after 6 days.

Numerical analysis. A total of 167 characters were included in the numerical taxonomy analysis. The distance matrix was calculated by using the Jaccard coefficient (23). Cluster analysis was done by using the unweighted pair group method with averages (23).

DNA extraction. Previously described methods were used to extract and purify DNA (1).

DNA-DNA hybridization. Native DNAs were labeled in vitro by nick translation with tritium-labeled nucleotides (Amersham International, Amersham, England). The procedure used for the hybridization experiments (S1 nuclease-trichloroacetic acid method) has been described previously (3, 8). The reassociation temperature was 60°C. DNA-DNA hybridization tests were carried out by using labeled DNAs from *P. syringae* pv. *syringae* CFBP 1392^T and *P. syringae* subsp. *savastanoi* CFBP 1670^T.

Thermal stability of reassociated DNAs. The temperature at which 50% of reassociated DNA became hydrolyzable by the S1 enzyme (*T_m*) was determined by using the method of

TABLE 2. Origins of 33 fluorescent and nonfluorescent oxidase-positive *Pseudomonas* strains and *A. tumefaciens* and *X. campestris* pv. *campestris* strains

Taxon	CFBP no.	Host or origin and place and year of isolation
<i>Pseudomonas aeruginosa</i>	2466 ^T	Host, origin, and year unknown
<i>P. alcaligenes</i>	2437 ^T	Water, origin and year unknown
<i>P. caryophylli</i>	2429 ^T	<i>Dianthus caryophyllus</i> , United States, 1951
<i>P. cepacia</i>	2227 ^T	<i>Allium cepa</i> , United States, year unknown
<i>P. cepacia</i>	2234	Hospital, France, year unknown
<i>P. corrugata</i>	2431 ^T	<i>Lycopersicon esculentum</i> , United Kingdom, 1972
<i>P. corrugata</i>	145.41	<i>Lycopersicon esculentum</i> , France, 1982
<i>P. corrugata</i>	30.5	<i>Lycopersicon esculentum</i> , France, 1984
<i>P. corrugata</i>	22.4	<i>Lycopersicon esculentum</i> , France, 1983
<i>P. fluorescens</i>	2102 ^T	Water, United Kingdom, 1951
<i>P. fluorescens</i>	2123	Water, Netherlands, 1966
<i>P. fluorescens</i>	2125	Host, origin, and year unknown
<i>P. fluorescens</i>	2127	Egg, United States, year unknown
<i>P. fluorescens</i>	2129	Water, origin and year unknown
<i>P. fluorescens</i>	2130	Water, origin and year unknown
<i>P. fluorescens</i>	2299	<i>Beta vulgaris</i> (root), France, 1983
<i>P. fuscovaginae</i>	2065 ^T	<i>Oryza sativa</i> , Japan, 1976
<i>P. gladioli</i> pv. <i>gladioli</i>	2427 ^T	<i>Gladiolus</i> sp., origin unknown, 1966
<i>P. gladioli</i> pv. <i>alliicola</i>	2422	<i>Allium cepa</i> , United States, 1939
<i>P. marginalis</i> pv. <i>marginalis</i>	1387 ^T	<i>Cichorium intybus</i> , United States, 1949
<i>P. marginalis</i> pv. <i>pastinacae</i>	2038 ^T	<i>Pastinaca sativa</i> , United States, 1959
<i>P. marginalis</i> pv. <i>alfalfae</i>	2039 ^T	<i>Medicago sativa</i> , United States, 1971
<i>P. putida</i>	2066 ^T	Soil, United States, year unknown
<i>P. putida</i>	2298	<i>Malus sylvestris</i> (root), France, 1983
<i>P. pseudoalcaligenes</i>	2435 ^T	Sinus drainage, origin unknown, 1966
<i>P. rubrilineans</i>	1294	<i>Saccharum officinale</i> , Réunion, 1970
<i>P. rubrisubalbicans</i>	1296	<i>Saccharum officinale</i> , Réunion, 1970
<i>P. solanacearum</i>	2047 ^T	<i>Lycopersicon esculentum</i> , United States, 1953
<i>P. solanacearum</i> (race 3)	1420	<i>Solanum phureja</i> , Colombia, 1965
<i>P. solanacearum</i> (race 2)	1482	<i>Musa</i> sp., Panama, 1958
<i>P. stutzeri</i>	2443 ^T	Spinal fluid, United States, 1970
<i>P. tolaasii</i>	2068 ^T	<i>Agaricus bisporus</i> , United Kingdom, 1965
<i>P. tolaasii</i>	2152	<i>Agaricus bisporus</i> , France, 1975
<i>A. tumefaciens</i>	2413 ^T	Host plant and origin unknown, 1972
<i>X. campestris</i> pv. <i>campestris</i>	2350	<i>Brassica oleracea</i> , United Kingdom, 1957

FIG. 1. Dendrogram of the distances among 231 strains of *P. savastanoi*, *P. syringae* sensu lato, and related *Pseudomonas* spp.

TABLE 3. Distribution of species and pathovars among eight phenotypes

Phenon or group	Subphenon	No. of strains	Strain(s) or no. of strains
I		15	<i>P. corrugata</i> CFBP 2431 ^T , CFBP 145.41, CFBP 30.5, and CFBP 22.4, <i>P. tolaasii</i> CFBP 2068 ^T and CFBP 2152, <i>P. fluorescens</i> CFBP 2299, CFBP 2129, CFBP 2102 ^T , CFBP 2123, and CFBP 2125, <i>P. marginalis</i> pv. pastinacea CFBP 2038 ^T , <i>P. marginalis</i> pv. marginalis CFBP 1387 ^T , <i>P. marginalis</i> pv. alfalfae CFBP 2039 ^T , <i>P. putida</i> CFBP 2298
II		3	<i>P. fluorescens</i> CFBP 2127 and CFBP 2130, <i>P. aeruginosa</i> CFBP 2466 ^T
III		4	<i>P. gladioli</i> pv. gladioli CFBP 2427 ^T , <i>P. gladioli</i> pv. alliicola CFBP 2422, <i>P. cepacia</i> CFBP 2227 ^T , and CFBP 2234
IV	IVa	3	<i>P. syringae</i> pv. photiniae CFBP 2899, <i>P. syringae</i> pv. myricae CFBP 2897, <i>P. syringae</i> pv. aesculi CFBP 2894
	IVb	2	<i>P. syringae</i> pv. thea CFBP 2353, <i>P. syringae</i> pv. taetis CFBP 1694,
	IVc	11	<i>P. syringae</i> pv. japonica CFBP 2896, <i>P. syringae</i> pv. hibisci CFBP 2895, <i>P. syringae</i> pv. mellea CFBP 2344, <i>P. syringae</i> pv. tabaci CFBP 2106, <i>P. syringae</i> pv. dysoxyl CFBP 2356, <i>P. syringae</i> pv. aptata CFBP 1617, <i>P. syringae</i> pv. panici CFBP 2345, <i>P. syringae</i> pv. papulans CFBP 1754, <i>P. syringae</i> pv. atrofaciens CFBP 2213, <i>P. syringae</i> pv. primulae CFBP 1660, <i>P. syringae</i> pv. aceris CFBP 2339
	IVd	3	<i>P. syringae</i> pv. tomato CFBP 2212, <i>P. syringae</i> pv. maculicola CFBP 1657, <i>P. syringae</i> pv. apii CFBP 2103
	IVe	4	<i>P. syringae</i> pv. passiflorae CFBP 2346, <i>P. syringae</i> pv. delphinii CFBP 2215, <i>P. syringae</i> pv. coronafaciens CFBP 2216, <i>P. syringae</i> pv. striafaciens CFBP 1674
V	Va	12	Isolated phenotypes of <i>P. syringae</i> pv. helianthi CFBP 2067, <i>P. syringae</i> pv. pisi CFBP 2105, <i>P. syringae</i> pv. ribicola CFBP 2348; <i>P. syringae</i> pv. viridiflava CFBP 2107, <i>P. syringae</i> pv. cichorii CFBP 2101, <i>P. syringae</i> pv. morsprunorum CFBP 2351, <i>P. syringae</i> pv. eriobotryae CFBP 2343, <i>P. syringae</i> pv. lapsa CFBP 1731, <i>P. syringae</i> pv. syringae CFBP 1392, <i>P. syringae</i> pv. atropurpurea CFBP 2340, <i>P. syringae</i> pv. mori CFBP 1642, <i>P. syringae</i> pv. porri CFBP 1908
		4	<i>P. syringae</i> pv. philadelphia CFBP 2898, <i>P. syringae</i> pv. berberidis CFBP 1727, <i>P. syringae</i> pv. anthriscini CFBP 1620, <i>P. syringae</i> pv. viburni CFBP 1702
		4	<i>P. syringae</i> pv. persicae CFBP 1573, W24-1, W24-2, and W24-3
VI	Vb	1	Isolated phenotype of <i>P. syringae</i> pv. lacrymans CFBP 2440
		142	<i>P. syringae</i> subsp. <i>savastanoi</i> (57 strains), strains from oleander (33 strains), ash (39 strains), privet (6 strains), <i>Phillyrea</i> sp. (4 strains), and jasmine (3 strains)
VII		3	<i>P. syringae</i> pv. glycinea CFBP 2214, <i>P. syringae</i> pv. phaseolicola CFBP 1390, <i>P. syringae</i> pv. ulmi CFBP 1407
VIII		2	<i>P. amygdali</i> CFBP 2354 and CFBP W28-1
Isolated phenotypes		17	<i>P. fuscovaginae</i> CFBP 2065 ^T , <i>P. putida</i> CFBP 2066 ^T , <i>P. rubisubalbicans</i> CFBP 1296, <i>P. rubrilineans</i> CFBP 1294, <i>P. caryophylli</i> CFBP 2429 ^T , <i>A. tumefaciens</i> CFBP 2413 ^T , <i>P. syringae</i> pv. <i>syringae</i> CFBP 1574, <i>P. syringae</i> pv. <i>sesami</i> CFBP 1671, <i>P. syringae</i> subsp. <i>savastanoi</i> SO52, <i>P. syringae</i> pv. <i>cannabina</i> CFBP 2341, <i>P. stutzeri</i> CFBP 2443 ^T , <i>P. alcaligenes</i> CFBP 2437 ^T , <i>P. solanacearum</i> CFBP 2047 ^T , CFBP 1482, and CFBP 1420, <i>X. campestris</i> pv. <i>campestris</i> CFBP 2350, <i>P. pseudoalcaligenes</i> CFBP 2435 ^T

Crosa et al. (3). The ΔT_m was the difference between the T_m of the heteroduplex and the T_m of the homoduplex.

RESULTS

Numerical analysis. A dendrogram displaying the distance relationships among the 231 strains which we studied is shown in Fig. 1.

At a distance of 0.346, eight phenotypes and 17 isolated strains were observed. The distribution of species, subspecies, and pathovars in the eight major phenotypes is shown in Table 3.

Phenotype I and II contained fluorescent oxidase-positive *Pseudomonas* strains. Phenotype III contained only nonfluorescent phytopathogenic *Pseudomonas* strains.

Phenotype IV to VIII included reference strains of fluorescent and nonfluorescent (*P. amygdali*) phytopathogenic *Pseudomonas* spp. that were either oxidase negative or oxidase positive (*P. cichorii*), corresponding to groups I through III of Lelliott et al. (15).

Phenotype IV was subdivided into five subphenotypes (subphenotype

IVa to IVe), and phenotype V was subdivided into subphenotypes Va and Vb.

Phenotype VI contained *P. syringae* subsp. *savastanoi* strains. All but one strain of *P. syringae* subsp. *savastanoi* fell into this phenotype. All 39 strains isolated from ash were clustered in one subphenotype. The strains isolated from olive and oleander did not constitute a subphenotype within phenotype VI.

Only 4 of the 17 unclustered strains were identified as *P. syringae* strains (*P. syringae* subsp. *savastanoi* SO52, *P. syringae* pv. *cannabina* CFBP 2341, *P. syringae* pv. *syringae* CFBP 1574, and *P. syringae* pv. *sesami* CFBP 1671) (Table 3).

Phenotype I to III were clearly differentiated by biochemical tests from phenotype IV to VIII. Strains in phenotype IV to VIII could not utilize *N*-acetylglucosamine, acetate, *N*-valerate, arginine, spermine, tyrosine, 2-ketogluconate, ethanolamine, L-tryptophan, L-ornithine, DL-kynurenine, isobutyrate, citraconate, itaconate, mesaconate, levulinate, and

TABLE 4. Characteristics that differentiate phenon IV to VIII

Phenon or subphenon	Utilization of:										Levan production	Fluorescence
	meso-Tartrate	DL-Hydroxybutyrate	Erythritol	DNase	Esculin	L-(+)-Tartrate	DL-Lactate	Glucosamine	β -Alanine	D-(-)-Tartrate		
IVa	+	—	—	+	—	+	—	—	—	—	+	+
IVb	+	d	—	—	—	d	—	+	+	—	+	+
IVc	+	+	+	—	+	d	d	d	—	d	+	+
IVd	+	+	—	d	—	—	+	—	—	+	+	+
IVe	+	—	+	—	+	—	—	—	—	—	+	+
Va	+	—	—	d	d	—	—	—	—	+	+	+
Vb	—	d	—	+	—	—	—	—	—	+	+	+
VI	—	—	—	+	—	+	—	—	—	—	—	+
VII	d	—	d	d	—	+	—	—	—	d	+	+
VIII	—	—	—	—	—	—	—	—	—	—	+	—

^a +, 90 to 100% of the strains are positive; —, 0 to 10% of the strains are positive; d, 11 to 89% of the strains are positive.

p-aminobenzoate and could not reduce nitrate. All of the strains in phenon I to III gave the opposite reactions.

Characteristics that differentiate phenon IV to VIII are shown in Table 4. Identification of a strain as a phenon IV or V strain requires identification at the subphenon level (Table 4). The reactions exhibited by the phenon VI strains differed somewhat when the source of isolation was considered (Table 5).

DNA relatedness. Results of DNA relatedness experiments are shown in Table 6. *P. syringae* subsp. *savastanoi* strains isolated from six hosts were 75 to 100% related to type strain CFBP 1670. For the two lowest relatedness values (75 and 79%), the ΔT_m values were 1.5 and 3.0°C, respectively. The reference strains of *P. syringae* pv. *glycinea* and *P. syringae* pv. *phaseolicola* were 72 and 83% related to strain CFBP 1670^T, respectively. The ΔT_m value calculated for *P. syringae* pv. *glycinea* and *P. syringae* subsp. *savastanoi* hybridized DNAs was 2.5°C. Thus, *P. syringae* subsp. *savastanoi*, *P. syringae* pv. *glycinea*, and *P. syringae* pv. *phaseolicola* are members of a single DNA hybridization group.

The type strain of *P. syringae* pv. *syringae*, strain CFBP 1392, was 43% related to strain CFBP 1670^T when DNA from strain CFBP 1670^T was labeled; the level of relatedness was 54% when DNA from strain CFBP 1392^T was labeled. *P. syringae* pv. *persicae*, *P. syringae* pv. *tomato*, and the other *Pseudomonas* species which we studied were less than 51% related to either *P. syringae* subsp. *savastanoi* CFBP 1670^T or *P. syringae* pv. *syringae* CFBP 1392^T; thus, these strains are members of DNA groups other than the *P. syringae* pv. *syringae* or *P. syringae* subsp. *savastanoi* DNA group. In this study the *P. syringae* pv. *syringae* DNA group was represented only by the type strain.

DISCUSSION

Before 1978, a *Pseudomonas* strain that was isolated for the first time from a new diseased host was considered a species. Young et al. (27) and Dye et al. (5) considered the named, phytopathogenic, oxidase-negative fluorescent, *Pseudomonas* spp. to be pathovars of *P. syringae*. At the present time, 45 pathovars are recognized (5, 26).

Janse (11) found that strains of *P. syringae* pv. *savastanoi* that were isolated from different hosts had nearly identical biochemical and physiological characteristics. Variations were observed only in the production of levan, in the hydrolysis of pectate, and in the production of indoleacetic acid and cytokininlike compounds. Janse also observed a marked variation in the results of pathogenicity tests among the strains that were isolated from different hosts. For these reasons he named this group of strains *P. syringae* subsp. *savastanoi* and proposed the following three pathovars: *P. syringae* subsp. *savastanoi* pv. *oleae*, *P. syringae* subsp. *savastanoi* pv. *nerii*, and *P. syringae* subsp. *savastanoi* pv. *fraxini* (12, 17).

Cross-pathogenicity is currently being tested by one of us (L.G.) to determine whether pathogenicity characteristics match taxonomic grouping. Pathogenicity of olive strains on oleander has not been demonstrated (6a, 11). All strains of *P. syringae* subsp. *savastanoi* are pathogenic on ash. Ash strains are pathogenic only on ash, and 80% of oleander strains are pathogenic on olive. However, cross-pathogenicity data for different hosts are not complete.

On the basis of DNA-DNA hybridization, physiological and biochemical characteristics, we propose that *P. syringae* subsp. *savastanoi* should be elevated to species level as

TABLE 5. Reactions of phenon VI strains (*P. syringae* subsp. *savastanoi*) isolated from different hosts

Host	No. of isolates	% of strains positive										Hydrolysis of polypectate (pH 5)	Levan production
		Utilization of:											
		L-Serine	<i>n</i> -Caproate	DL-5-Amino-valerate	L-Leucine	Raffinose	2-Amino-benzoate	L-Tyrosine	D-Xylose	Trigo-nelline	L-Arabi-nose		
Olive	58	100	12	0	50	0	0	0	45	50	37	13	0
Ash	39	21	82	53	92	23	13	18	92	0	95	100	21
Oleander	33	100	18	0	76	0	0	0	3	0	79	97	0
Jasmine	3	100	66	0	100	0	0	0	0	0	100	100	0
<i>Phillyrea</i> sp.	4	100	0	0	100	0	0	0	0	0	100	100	0
Privet	6	100	0	0	16	0	0	0	0	0	100	16	16

TABLE 6. Levels of DNA relatedness among *Pseudomonas* strains

Unlabeled DNA from:		% of relative binding at 60°C with labeled DNA from:	
Taxon	Strain (host)	<i>P. syringae</i> CFBP 1392 ^T	<i>P. savastanoi</i> CFBP 1670 ^T
<i>P. savastanoi</i> pv. <i>savastanoi</i> (phenon VI)	CFBP 1670 ^T (olive)	54	100
	K124-4 (olive)	53	82
	T35-1 (olive)	49	87
	T12-6 (olive)	64	83
	K23-15 (olive)	49	93
	CFBP 2088 (oleander)	58	79 (3) ^a
	L145-2 (oleander)	52	88
	T37-6 (oleander)	49	90
	L36-7 (oleander)	50	84
	L86-1 (oleander)	53	81
	CFBP 1838 (ash)	58	86
	CFBP 2093 (ash)	58	97
	T36-3 (ash)	55	92
	T5-1 (ash)	47	75 (1.5)
	T12-7 (jasmine)	60	92
	CFBP 1751 (jasmine)	53	88
	T12-10 (jasmine)	54	83
	Phi (<i>Phillyrea</i> sp.)	54	100
	T51-3 (<i>Phillyrea</i> sp.)	62	93
	T51-1 (<i>Phillyrea</i> sp.)	50	81
	T37-11 (privet)	59	86
	T35-10 (privet)	57	86
	T37-15 (privet)	53	90
<i>P. savastanoi</i> pv. <i>phaseolicola</i> (phenon VII)	CFBP 1390 ^T	56	83
<i>P. savastanoi</i> pv. <i>glycinea</i> (phenon VII)	CFBP 2214 ^T	50	72 (2.5)
<i>P. syringae</i> pv. <i>syringae</i> (isolated phenotype)	CFBP 1392 ^T	100	43
<i>P. syringae</i> pv. <i>persicae</i> (phenon IVb)	CFBP 1573 ^T	46	41
<i>P. syringae</i> pv. <i>tomato</i> (phenon IVd)	CFBP 2212 ^T	51	47
<i>P. viridiflava</i> (isolated phenotype)	CFBP 2107 ^T	49	44
<i>P. cichorii</i> (isolated phenotype)	CFBP 2101 ^T	25	24
<i>P. marginalis</i> (phenon I)	CFBP 1387 ^T	21	22
<i>P. fluorescens</i> (phenon II)	CFBP 2102 ^T	20	21
<i>P. putida</i> (phenon I)	CFBP 2066 ^T	9	10

^a The values in parentheses are ΔT_m values (in degrees Celsius).

Pseudomonas savastanoi and that this species should include three pathovars, *P. savastanoi* pv. *savastanoi*, *P. savastanoi* pv. *glycinea*, and *P. savastanoi* pv. *phaseolicola*.

Description of *Pseudomonas savastanoi* (Smith). *Pseudomonas savastanoi* (sa.vas.ta'no.i. L.gen.n. *savastanoi* of Savastano, the first worker who studied olive knot disease). Gram-negative rods that are 0.4 to 0.8 by 1.0 to 3.0 μ m and motile by means of one to four polar flagella. Rather slow growing. Colonies are white or cream, smooth, flat, and glistening with entire or erose margins on YBGA. Produces a hypersensitive reaction on tobacco leaves. Metabolism is respiratory. Oxidase negative. Nitrates are not reduced. Blue fluorescent pigment is produced under UV light on King B medium. Arginine test (Thornley) negative. Esculin, gelatin, and starch are not hydrolyzed. Assimilates sucrose, L-arabinose, gluconate, caprylate, fumarate, DL-glycerate, L-malate, pyruvate, citrate, D- α -alanine, and L-proline. Does not assimilate lactose, L-xylose, adonitol, 2-aminobutyrate, DL-lactate, DL-3-hydroxybutyrate, D-(–)-tartrate, L-cysteine, L-methionine, and L-valine. Strains isolated from members of various genera of the Oleaceae and *N. oleander* do not produce levan, whereas strains isolated from *F. excelsior*

L. (21%), *Phaseolus vulgaris*, and *Glycine max* L. do produce levan. The G+C content of the DNA is 60 mol% (4).

P. savastanoi pv. *savastanoi* causes knots, galls, and cankers on members of the various genera of the Oleaceae and *N. oleander* L.; *P. savastanoi* pv. *glycinea* causes bacterial blight of soybean; and *P. savastanoi* pv. *phaseolicola* causes halo blight of bean.

The type strain is strain NCPPB 639 (= ATCC 13522 = ICMP 4352 = CFBP 1670).

Description of the type strain. Strain NCPPB 639^T was isolated from *O. europea* L. in Yugoslavia. This strain has physiological and biochemical characteristics that are typical of the species. In addition, it does not assimilate sorbitol and *trans*-aconitate.

REFERENCES

1. Brenner, D. J., A. C. McWorter, J. K. Leete Knutson, and A. G. Steigerwalt. 1982. *Escherichia vulneris*: a new species of *Enterobacteriaceae* associated with human wounds. J. Clin. Microbiol. 15:1133–1140.
2. Brown, N. A. 1932. Canker of ash trees produced by a variety of the olive tubercule organism, *Bacterium savastanoi*. J. Agric. Res. 44:701–722.

3. Crosa, J. M., D. J. Brenner, and S. Falkow. 1973. Use of a single-strand-specific nuclease for analysis of bacterial and plasmid deoxyribonucleic acid homo- and heteroduplexes. *J. Bacteriol.* 115:904-911.
4. De Ley, J. 1968. DNA base composition and hybridization in the taxonomy of phytopathogenic bacteria. *Annu. Rev. Phytopathol.* 6:63-90.
5. Dye, D. W., J. F. Bradbury, M. Goto, A. C. Hayward, R. A. Lelliott, and M. N. Schroth. 1980. International standards for naming pathovars of phytopathogenic bacteria and a list of pathovar names and pathotype strains. *Rev. Plant Pathol.* 59:153-168.
6. Ferraris, T. 1926. *Trattato di patologia e terapia vegetale*, vol. 1, 3rd ed. Hopli, Milan, Italy.
- 6a. Gardan, L. Unpublished data.
7. Gardan, L., S. Cottin, C. Bollet, and G. Hunault. 1991. Phenotypic heterogeneity of *Pseudomonas syringae*. *Res. Microbiol.* 142:995-1003.
8. Grimont, P. A. D., M. Y. Popoff, F. Grimont, C. Coynault, and M. Lemelin. 1980. Reproducibility and correlation study of three deoxyribonucleic acid hybridization procedures. *Curr. Microbiol.* 4:325-330.
9. Hildebrand, D. C. 1971. Pectate and pectin gels for differentiation of *Pseudomonas* sp. and other bacterial plant pathogens. *Phytopathology* 61:1430-1436.
10. Hildebrand, D. C., and M. N. Schroth. 1972. Identification of the fluorescent pseudomonads, p. 281-287. In H. P. Maas Geesteranus (ed.), *Proceedings of the Third International Conference on Plant Pathogenic Bacteria*. Pudoc, Wageningen, The Netherlands.
11. Janse, J. D. 1981. The bacterial disease of ash (*Fraxinus excelsior*) caused by *Pseudomonas syringae* subsp. *savastanoi* pv. *fraxini*. II. Etiology and taxonomic considerations. *Eur. J. For. Pathol.* 11:425-438.
12. Janse, J. D. 1982. *Pseudomonas syringae* subsp. *savastanoi* (ex Smith) subsp. nov., nom. rev., the bacterium causing excrescences on Oleaceae and *Nerium oleander* L. *Int. J. Syst. Bacteriol.* 32:166-169.
13. Lapage, S. P., S. Bascomb, W. R. Wilcox, and M. A. Curtis. 1973. Identification of bacteria by computer: general aspects and perspectives. *J. Gen. Microbiol.* 77:273-290.
14. Lapage, S. P., P. H. A. Sneath, E. F. Lessel, V. B. D. Skerman, H. P. R. Seeliger, and W. A. Clark (ed.). 1975. International code of nomenclature of bacteria. 1975 Revision. American Society for Microbiology, Washington, D.C.
15. Lelliott, R. A., E. Billing, and A. C. Hayward. 1966. A determinative scheme for the fluorescent plant pathogenic pseudomonads. *J. Appl. Bacteriol.* 29:470-489.
16. Misaghi, I., and R. G. Grogan. 1969. Nutritional and biochemical comparisons of plant pathogenic and saprophytic fluorescent pseudomonads. *Phytopathology* 59:1436-1451.
17. Moore, W. E. C., and L. V. H. Moore (ed.). 1989. Index of bacterial and yeast nomenclatural changes, p. 49. American Society for Microbiology, Washington, D.C.
18. Pecold, P. C., and R. G. Grogan. 1973. Deoxyribonucleic acid homology groups among phytopathogenic *Pseudomonas* species. *Int. J. Syst. Bacteriol.* 23:111-121.
19. Prunier, J. P., and P. Kaiser. 1964. Etude de l'activité pectinolytique chez des bactéries phytopathogènes et saprophytes des plantes. I. Recherche des enzymes pectinolytiques. *Ann. Epiphyt. (Paris)* 15:205-219.
20. Sands, D. C., M. N. Schroth, and D. C. Hildebrand. 1970. Taxonomy of phytopathogenic pseudomonads. *J. Bacteriol.* 101:9-23.
21. Skerman, V. B. D., V. McGowan, and P. H. A. Sneath (ed.). 1989. Approved lists of bacterial names. Amended edition. American Society for Microbiology, Washington, D.C.
22. Smith, E. D. 1908. Recent studies on the olive tubercule organism. *Bull. Bur. Plant Ind. U. S. Dep. Agric.* 131:25-43.
23. Sneath, P. H. A., and R. R. Sokal. 1973. Numerical taxonomy. The principles and practice of numerical classification. W. H. Freeman, San Francisco.
24. Stevens, F. L. 1913. The fungi which cause plant disease. Macmillan Co., New York.
25. Wilson, E. E., M. G. Heskett, M. L. Johnson, and T. Kosuge. 1972. Metabolic behaviour of *Pseudomonas savastanoi* isolates from olive and oleander on certain carbohydrates and amino substrates. *Phytopathology* 62:350-355.
26. Young, J. M., J. F. Bradbury, R. E. Davis, R. S. Dickey, and G. L. Ercolani. 1991. Nomenclatural revisions of plant pathogenic bacteria and list of names, 1980-1988. *Rev. Plant Pathol.* 70:221.
27. Young, J. M., D. W. Dye, J. F. Bradbury, C. G. Panagopoulos, and C. F. Robb. 1978. A proposed nomenclature and classification for plant pathogenic bacteria. *N. Z. J. Agric. Res.* 21:153-177.