Tuberculosis in seals caused by a novel member of the *Mycobacterium tuberculosis* complex: *Mycobacterium pinnipedii* sp. nov.

Debby V. Cousins,¹ Ricardo Bastida,² Angel Cataldi,³ Viviana Quse,⁴ Sharon Redrobe,⁵ Sue Dow,⁵ Padraig Duignan,⁶ Alan Murray,⁶ Christine Dupont,⁶ Niyaz Ahmed,⁷ Des M. Collins,⁸ W. Ray Butler,⁹ David Dawson,¹⁰ Diego Rodríguez,² Julio Loureiro,³ Maria Isabel Romano,³ A. Alito,³ M. Zumarraga³ and Amelia Bernardelli¹¹

Correspondence

Debby V. Cousins

dcousins@agric.wa.gov.au

¹Australian Reference Laboratory for Bovine Tuberculosis, Department of Agriculture Western Australia, 3 Baron-Hay Court, South Perth, WA 6151, Australia

²CONICET and Departamento de Ciencias Marinas, Facultad de Ciencias Exactas y Naturales, Universidad Nacional de Mar del Plata, Casilla de Correo 43, (7600) Mar del Plata, Argentina

³Instituto de Biotecnología, CICVyA, Instituto Nacional de Tecnología Agropecuaria, Los Reseros y Las Cabañas, (1712) Castelar, Argentina

⁴Fundación Mundo Marino, Avda Décima 157, (7105) San Clemente del Tuyú, Argentina

⁵Bristol Zoo Gardens, Bristol BS8 3HA, UK

⁶Pathobiology Group, Institute of Veterinary, Animal and Biomedical Sciences, Massey University, Private Bag 11-222, Palmerston North, New Zealand

⁷Centre for DNA Fingerprinting and Diagnostics (CDFD), Nacharam, Hyderabad 500 076, India

⁸AgResearch, Wallaceville Animal Research Centre, Upper Hutt, New Zealand

⁹Division of AIDS, STD and TB Laboratory Research, National Center for Infectious Diseases, Centers for Disease Control and Prevention, Atlanta, GA 30333, USA

¹⁰Mycobacterium Reference Laboratory, Queensland Health Pathology Services, Prince Charles Hospital, Chermside, Qld 4032, Australia

¹¹Departamento de Micobacterias, DILACOT, Servicio Nacional de Sanidad y Calidad Agroalimentaria (SENASA), Avda A Fleming 1653, (1640) Martínez, Argentina

A comparison of *Mycobacterium tuberculosis* complex isolates from seals (pinnipeds) in Australia, Argentina, Uruguay, Great Britain and New Zealand was undertaken to determine their relationships to each other and their taxonomic position within the complex. Isolates from 30 cases of tuberculosis in six species of pinniped and seven related isolates were compared to representative and standard strains of the *M. tuberculosis* complex. The seal isolates could be distinguished from other members of the *M. tuberculosis* complex, including the recently defined '*Mycobacterium canettii*' and '*Mycobacterium caprae*', on the basis of host preference and phenotypic and genetic tests. Pinnipeds appear to be the natural host for this 'seal bacillus', although the organism is also pathogenic in guinea pigs, rabbits, humans, Brazilian tapir (*Tapirus terrestris*) and, possibly, cattle. Infection caused by the seal bacillus is predominantly associated with granulomatous lesions in the peripheral lymph nodes, lungs, pleura, spleen and peritoneum. Cases of disseminated disease have been found. As with other members of the *M. tuberculosis* complex, aerosols are the most likely route of transmission. The name *Mycobacterium pinnipedii* sp. nov. is proposed for this novel member of the *M. tuberculosis* complex (the type strain is 6482^T=ATCC BAA-688^T=NCTC 13288^T).

Published online ahead of print on 24 January 2003 as DOI 10.1099/ijs.0.02401-0.

Abbreviations: BCG, Bacille Calmette–Guérin; FAFLP, fluorescent amplified fragment length polymorphism; PZA, pyrazinamide; SS, seal spoligotype; TCH, thiophen-2-carboxylic acid hydrazide.

The GenBank/EMBL/DDBJ accession number for the 16S rRNA gene sequence of strain 6482^T is AF502574.

INTRODUCTION

The *Mycobacterium tuberculosis* complex has traditionally consisted of four members: *M. tuberculosis* (*sensu stricto*), which primarily infects human and primates; *Mycobacterium bovis*, which predominantly causes tuberculosis in cattle (Karlson & Lessel, 1970), but can also cause disease in a wide variety of other animals, including man; *M. bovis* Bacille Calmette–Guérin (BCG), an attenuated strain that is used for vaccination; *Mycobacterium africanum*, a heterogeneous group of isolates responsible for human tuberculosis in Africa, which appears to be intermediate between *M. tuberculosis* and *M. bovis* (Castets *et al.*, 1969); and *Mycobacterium microti*, a less frequently isolated pathogen that traditionally causes tuberculosis in voles (Wells & Oxen, 1937; Wells & Robb-Smith, 1946), but has been identified more recently as a cause of disease in immunocompromised humans (van Soolingen *et al.*, 1998). Each member of the *M. tuberculosis* complex is associated with a specific primary host, although infection is known to occur in various alternative hosts. Although all of these strains effectively share the same 16S rRNA gene sequence (Rogall *et al.*, 1990b) and high DNA–DNA homology (from hybridization studies), they can be separated by some phenotypic characteristics (Grange & Yates, 1994) and, as they have different primary hosts, they have been regarded as separate species (Collins *et al.*, 1982). More recently, two novel strains have been described: ‘*Mycobacterium canettii*’, a novel smooth variant of *M. tuberculosis* that was first isolated from a Somali-born patient (van Soolingen *et al.*, 1997) and subsequently from a Swiss patient exposed in Africa (Pfyffer *et al.*, 1998); and ‘*Mycobacterium caprae*’ (basonym: *M. tuberculosis* subsp. *caprae*), a strain that occurs primarily in Spanish goats and has recently been elevated to species level (Aranaz *et al.*, 1999; Niemann *et al.*, 2002; Aranaz *et al.*, 2003). *M. tuberculosis*, *M. africanum*, *M. microti* and *M. bovis* were accepted as separate species by using a combination of phenotypic characteristics and apparent host specificity. ‘*M. canettii*’ and ‘*M. caprae*’ were accepted by virtue of host preference and phenotypic and genetic characteristics.

Previous logical arguments have suggested that all members of the *M. tuberculosis* complex are sufficiently similar to belong to the species *M. tuberculosis* (Wayne, 1984; Tsukamura *et al.*, 1985; van Soolingen *et al.*, 1997). Acceptance of such an approach would cause currently accepted species (*M. bovis*, *M. microti* and *M. africanum*) to warrant classification at the subspecies level.

Between 1986 and 1995, *M. tuberculosis* complex organisms were isolated from cases of tuberculosis in captive or wild Australian sea lions (*Neophoca cinerea*), New Zealand fur seals (*Arctocephalus forsteri*), an Australian fur seal (*Arctocephalus pusillus doriferus*) and a seal trainer who worked with the affected colony in Australia (Forshaw & Phelps, 1991; Thompson *et al.*, 1993; Cousins, 1995; Woods *et al.*, 1995). Similar organisms were recovered from captive Southern sea lions (*Otaria flavescens*), wild South American

fur seals (*Arctocephalus australis*) and a wild Subantarctic fur seal (*Arctocephalus tropicalis*) in Uruguay or Argentina during the period 1989–2000 (Bernardelli *et al.*, 1994, 1996; Castro Ramos *et al.*, 1998; Bastida *et al.*, 1999). Between 1996 and 1998, *M. tuberculosis* complex organisms were recovered from two South American fur seals in a zoological collection in Great Britain and a Brazilian tapir (*Tapirus terrestris*) housed in an adjacent enclosure, and from two New Zealand fur seals in New Zealand in 1997 and 1998 (Hunter *et al.*, 1998).

Many of the isolates obtained from cases of tuberculosis in Australia, Uruguay and Argentina have been well-characterized (Cousins *et al.*, 1993; Bernardelli *et al.*, 1996; Cousins, 1996; Alito *et al.*, 1999; Zumárraga *et al.*, 1999) and this information, together with preliminary tests on the seal isolates from Great Britain and New Zealand, suggested that the seal bacillus (Cousins *et al.*, 1993), isolated from pinnipeds from all four continents, may be a unique member of the *M. tuberculosis* complex. A recent study of four Australian and six Argentinian seal isolates by fluorescent amplified fragment length polymorphism (FAFLP) has further substantiated the hypothesis that the seal bacillus occupies a unique taxonomic position within the *M. tuberculosis* complex (Ahmed *et al.*, 2003). This report consolidates the results of tests performed previously and provides additional information, resulting in a comprehensive comparison of isolates available from pinniped-related cases of tuberculosis, and indicates that the seal bacillus should be considered as a novel species of the *M. tuberculosis* complex.

METHODS

Bacterial strains. *M. tuberculosis* complex isolates recovered from 30 pinnipeds and a seal trainer from 1985 to 2000 were available for study (Table 1). Isolates recovered from two guinea pigs and two rabbits after pathogenicity experiments in Australia, a bovine in New Zealand (this isolate had a restriction endonuclease analysis pattern similar to those of the Australian seal isolates) and a Brazilian tapir were also included. Where appropriate, isolates were compared to representative (and reference) strains of *M. tuberculosis* (H37Rv or Mt14323), *M. africanum* (TMC3), ‘*M. caprae*’ (CIP 105776^T), *M. microti* (NCTC 8710^T), *M. bovis* (AN5) and *M. bovis* BCG (P3) (Table 2).

Phenotypic characteristics. Isolates were examined for growth and phenotypic characteristics according to standard procedures (Vestal, 1975). *In vitro* susceptibility patterns to isoniazid, rifampicin, streptomycin and ethambutol were determined for three isolates from Australia, three from Argentina, one from Uruguay, two from Great Britain and two from New Zealand by using the Mycobacterial Growth Indicator Tube (MGIT; Becton Dickinson) system (Bernardelli *et al.*, 1999; Morcillo *et al.*, 2000).

Pathogenicity studies in guinea pigs and rabbits. Isolates from two Australian seals (Au-1 and Au-2) were each injected into a guinea pig and a rabbit, and three isolates from Argentina (Ar-1, Ar-2 and Ar-3) and the isolate from Uruguay (U-1) were inoculated into guinea pigs, to examine the pathogenicity of the seal isolates.

Table 1. *M. tuberculosis* complex isolates recovered from pinnipeds or related cases from various countries, 1985–2000

Isolate (reference no.)	Source	Year
Australia		
Au-1 (6481)	Captive <i>Neophoca cinerea</i> (Australian sea lion)	1985
(4823F)	Guinea pig LN ex 6481	
(4524D)	Rabbit lung ex 6481	
Au-2 (6482 ^T)	Captive <i>Arctocephalus forsteri</i> (New Zealand fur seal)	1986
(4821D)	Guinea pig LN ex 6482	
(1676)	Rabbit lung ex 6482	
Au-3 (6510)	Captive <i>Neophoca cinerea</i>	1986
Au-4 (6954)	Captive <i>Neophoca cinerea</i>	1986
Au-5 (6866)	Captive <i>Arctocephalus forsteri</i>	1986
Au-6 (6884)	Captive <i>Neophoca cinerea</i>	1986
Au-7 (146-D)	Human (seal trainer)	1988
Au-8 (14109)	Wild <i>Neophoca cinerea</i> 1	1991
Au-9 (14126)	Wild <i>Neophoca cinerea</i> 2	1992
Au-10 (92/1161T)	Wild <i>Arctocephalus pusillus doriferus</i> (Australian fur seal)	1992
Au 11 (A95-127)	Wild <i>Arctocephalus forsteri</i>	1995
Uruguay		
U-1 (1337)	Captive <i>Otaria flavescens</i> (Southern sea lion)	1987
Argentina		
Ar-1 (1203-5)	Wild <i>Arctocephalus australis</i> (South American fur seal)	1989
Ar-2 (M-17-92, 1489-50)	Wild <i>Arctocephalus australis</i>	1992
Ar-3 (M-31-92, 1855-8)	Wild <i>Arctocephalus australis</i>	1992
Ar-4 (M-38-92, 1862-4)	Wild <i>Arctocephalus australis</i>	1992
Ar-5 (M-47-92, 1866-7)	Wild <i>Otaria flavescens</i>	1992
Ar-6 (M-11-91, 1868-9)	Wild <i>Arctocephalus australis</i>	1995
Ar-7 (M-05-95, 1920-1)	Wild <i>Arctocephalus australis</i>	1995
Ar-8 (M-35-95, 1981-2)	Wild <i>Arctocephalus australis</i>	1995
Ar-9 (M-08-96, 2003 & 5)	Wild <i>Arctocephalus australis</i>	1996
Ar-10 (M-55-95, 2007-9)	Wild <i>Arctocephalus tropicalis</i> (Subantarctic fur seal)	1996
Ar-11 (M-10-96, 2027 & 9)	Wild <i>Arctocephalus australis</i>	1996
Ar-12 (M-33-96, 2050-3)	Wild <i>Arctocephalus australis</i>	1996
Ar-13 (M-02-99, 2186)	Wild <i>Arctocephalus australis</i>	1999
Ar-14 (M-30-00, 2192)	Wild <i>Arctocephalus australis</i>	2000*
Ar-15 (M-33-00, 2225-6 & 9)	Wild <i>Arctocephalus australis</i>	2000*
United Kingdom		
UK-1 (623/971757)	Captive <i>Arctocephalus australis</i>	1996
UK-2 (624/97)	Captive <i>Tapirus terrestris</i> (Brazilian tapir)	1996*
UK-3 (2281)	Captive <i>Arctocephalus australis</i>	1998
New Zealand		
NZ-1	Bovine	1991*
NZ-1	Wild <i>Arctocephalus forsteri</i>	1997
NZ-2	Wild <i>Arctocephalus forsteri</i>	1998

*Unpublished.

Mycolic acid analysis. Mycolic acid profiles of two representative isolates (Au-1 and Au-2) were examined by HPLC, according to previously published procedures (Butler *et al.*, 1996, 1999).

Tests for MPB70. All isolates were tested for the presence of the MPB70 antigen either by using the immunoperoxidase test (Corner *et al.*, 1988; Veerman *et al.*, 1990; Liébana *et al.*, 1996) and/or by performing SDS-PAGE on antigen preparations (Cousins *et al.*, 1993; Alito *et al.*, 1999).

16S rDNA sequence determination and PCR-based tests for genetic markers. PCR-mediated amplification of 16S rDNA was performed by using procedures described previously (Edwards *et al.*, 1989; Kirschner *et al.*, 1993). The nucleotide sequences obtained were compared to all known 16S rDNA mycobacterial sequences in GenBank and the *M. bovis* sequence (available at the Sanger website, http://www.sanger.ac.uk/Projects/M_bovis/) by using the FastA application. Sequences were aligned with the program PILEUP from the Genetics Computer Group (GCG) version 9 UNIX software.

Table 2. *M. tuberculosis* complex isolates that were tested by spoligotyping and used to prepare the dendrogram (Fig. 2)

AFS, Australian fur seal; ASL, Australian sea lion; Bj, Beijing strain; COB, country of birth; NZFS, New Zealand fur seal; SAFS, South American fur seal; TMC, Trudeau Mycobacterium Collection.

Species/strain	Source of isolate	Country of origin/provided by
<i>M. africanum</i>		
1. TMC 3	Reference strain, human origin	
2. TMC 12	Reference strain, human origin	
3. TMC 54	Reference strain, human origin	
4. 19884 (4163/69)	Clinical isolate, human origin	Australia
5. 19887 (5166/88)	Clinical isolate, human origin	Australia
6. 19890 (486/93)	Clinical isolate, human origin	Australia
7. 22054	Clinical isolate, human origin	Australia
<i>M. bovis/M. bovis</i> BCG		
8. BCG Japanese	Vaccine strain	Richard Wallace, USA
9. BCG Russian	Vaccine strain	Richard Wallace, USA
10. BCG Pasteur	Vaccine strain	RIVM, Netherlands
11. AN5	Reference strain, cattle origin	CSIRO, Australia
12. 3958	Cattle	Australia (Western Australia)
13. 6205	Cattle	Australia (Western Australia)
14. 11487	Cattle	Australia (Western Australia)
15. 14457	Cattle	Australia (Western Australia)
16. 14899	Cattle	Australia (Western Australia)
17. 15145	Cattle	Australia (Western Australia)
18. 17319	Cattle	Australia (Queensland)
19. 17898	Cattle	Australia (Queensland)
20. 20007	Red deer	Canada
21. 22950	Goat	Spain
<i>M. microti</i>		
22. NCTC 8710 ^T	Reference strain, vole origin	
23. 3377	Vole	T. Jenkins, UK
24. 3381	Vole	T. Jenkins, UK
<i>M. tuberculosis</i>		
25. H37Rv	Reference strain, human origin	RIVM
26. 14323	Reference strain, human origin	RIVM
27. 26079	Clinical isolate, human origin	Diagnosed in Australia, COB Indonesia
28. 26141	Clinical isolate, human origin	Diagnosed in Australia, COB Indonesia
29. 26142	Clinical isolate, human origin	Diagnosed in Australia, COB Vietnam
30. 26152	Clinical isolate, human origin	Diagnosed in Australia, COB Vietnam
31. 27204	Clinical isolate, human origin, Bj	Diagnosed in Australia, COB India
32. 27206	Clinical isolate, human origin	Diagnosed in Australia, COB Yugoslavia
33. 27214	Clinical isolate, human origin, Bj	Diagnosed in Australia, COB Vietnam
34. 27222	Clinical isolate, human origin	Diagnosed in Australia, COB Australia
35. 27230	Clinical isolate, human origin	Diagnosed in Australia, COB Afghanistan
' <i>M. canettii</i> '		
36. So93	Clinical isolate, human origin	D. van Soolingen, Netherlands
' <i>M. caprae</i> '		
37. CIP 105776 ^T	Goat	Spain
38. 4/21	Goat	Spain
39. CB27	Goat	Spain
Seal bacilli		
40. SS-1, 146-D	Clinical isolate, seal trainer	Australia
41. SS-1, 6482 ^T	Captive NZFS	Australia
42. SS-1, 92/1162/T	Wild AFS	Australia
43. SS-2	Wild SAFS	Argentina
44. SS-3, 24890	Captive SAFS	Great Britain
45. SS-4, 25878	Wild NZFS	New Zealand

package. Phylogenetic analyses of the sequence data were done with programs from the Phylogeny Inference Package (PHYLIP) as described previously (Floyd *et al.*, 1996). The pairwise comparison program GAP, also from the GCG package, was used to determine the position of consensus strand nucleotides, relative to those of *Escherichia coli* (GenBank number J01859). The 16S rDNA sequence of strain 6482^T was deposited in GenBank under accession number AF502574.

PCR-based tests for known genetic markers. All isolates were tested by PCR for the presence of mycobacterial 16S rDNA, the gene that encodes the MPB70 antigen, the IS6110, IS1081 and *mtp40* sequences (Del Portillo *et al.*, 1991) and the P_{AN} promoter region that is present in pathogenic mycobacteria (Gormley *et al.*, 1997) by using previously published methods (Liébana *et al.*, 1996; Zumárraga *et al.*, 1999). Representative isolates were tested for *katG* and *gyrA* gene sequence polymorphisms at codons 463 and 95, respectively, by using methods described previously (Zumárraga *et al.*, 1999). Allele-specific polymorphisms were examined at nt 285 of the *oxyR* gene (Sreevatsan *et al.*, 1996), which differentiates *M. bovis* and '*M. caprae*' (adenine) from other members of the *M. tuberculosis* complex (guanidine), and in codon 57 (nt 169) of the *pncA* gene, which is responsible for pyrazinamide (PZA) resistance (Espínosa de los Monteros *et al.*, 1998), which is consistent with *M. bovis*.

Spoligotyping. All but two isolates were tested for known spacers between direct repeats in the DR allele by using the spoligotyping method developed by Kamerbeek *et al.* (1997) and performed as described previously (Aranaz *et al.*, 1996; Zumárraga *et al.*, 1999). Spoligotyping results were analysed by electronic scanning of images and converting and analysing them by using GelCompar version 1.3, as described previously (Romano *et al.*, 1995; Cousins *et al.*, 1998a, b). The patterns obtained from the South American isolates were compared to a database that consisted mostly of *M. bovis* isolates from Argentina. In addition, the patterns obtained from all seal isolates were compared with a large database of patterns that contained approximately 700 *M. tuberculosis* complex isolates, including approximately 500 *M. bovis* isolates from cattle, buffalo, deer, wild animals and humans from Australia and other countries (Cousins *et al.*, 1998a); they were also compared to the CDC database (Jack

Crawford, personal communication). The seal isolates and representative and reference strains of the *M. tuberculosis* complex (Table 2) were included in a dendrogram of spoligotype patterns that were generated by using Dice UPGMA analysis (GelCompar, version 3.1; Applied Maths) to examine the clonal relationships between them.

FAFLP. Heat-killed cells of isolates from three Australian sea lions, one Australian fur seal and six South American fur seals were digested by using *EcoRI*/*MseI* and analysed by FAFLP, using methods described previously (Ahmed *et al.*, 2002, 2003). Analyses were based on the differential amplification of 131 genomic loci. Standard genomic DNA from *M. tuberculosis* H37Rv, *M. bovis* AN5, *M. africanum* and *M. microti* (NCTC 8710^T) was used for comparative FAFLP analysis.

RESULTS AND DISCUSSION

Morphology, growth and phenotypic characteristics of seal isolates

A description of the physiological characteristics of the taxon can be found in the formal description. Biochemical testing clearly confirmed that the seal isolates belonged to the *M. tuberculosis* complex. The negative reactions in the nitrate reduction and niacin accumulation tests were consistent with *M. bovis* (Grange & Yates, 1994) (Table 3), a fact that led to their initial identification as such in Australia (Forshaw & Phelps, 1991), Argentina (Bernardelli *et al.*, 1996) and Great Britain. In some cases, varying amounts of niacin were produced, which is similar to results reported for *M. africanum* (Grange & Yates, 1994). In most cases, the seal isolates grew preferentially on media that contained sodium pyruvate, although some (including NZ-2 and NZ-3) also grew on Löwenstein–Jensen medium that contained glycerol. Slight differences from typical *M. bovis* isolates were noted in Australia and Argentina, in that the

Table 3. Phenotypic properties of the seal bacillus, compared to other members of the *M. tuberculosis* complex

Species: 1, *M. tuberculosis* (classic); 2, *M. tuberculosis* (Asian); 3, *M. africanum* (type I); 4, *M. africanum* (type II); 5, *M. microti*; 6, seal bacillus; 7, *M. bovis*; 8, '*M. caprae*'; 9, *M. bovis* BCG. Data were taken from references cited in the text. Abbreviations: +, positive; –, negative; v, variable; NA, not applicable.

Characteristic	1	2	3	4	5	6	7	8	9
Nitrate reduction	+	+	–	+	–	–	–	–	–
Niacin accumulation	+	+	v	v	+	–*	–	–	–
Pyruvate preference	–	–	–	–	–	+	+	+	+
Stimulated by glycerol	+	–	–	–	–	–	–	–	+
MPB70 antigen	–	–	–	–	–	–	+	?	+
Resistance to:									
TCH	+	–	–	+	–	–†	–	–‡	–
PZA	–	–	–	–	–	–	+	–	+
Pathogenicity in:									
Guinea pig	++		++		–	++	++	NA	–
Rabbit	–		+/-		–	+++	++	NA	–

*Occasional strains, including the isolates from New Zealand, gave weak or positive reactions in the niacin accumulation test.

†New Zealand strains were resistant to 1 µg TCH ml⁻¹, but sensitive to 10 µg TCH ml⁻¹.

‡Resistant to 1 and 2 µg TCH ml⁻¹, but sensitive to 5 and 10 µg TCH ml⁻¹.

cord formation observed after Ziehl–Neelsen staining was loose; further investigations uncovered differences between the seal isolates and *M. bovis*, including its susceptibility to PZA.

Pathogenicity and potential host range

Isolates inoculated into guinea pigs produced significant lesions or death within 6 weeks and those inoculated into rabbits caused death within 6 weeks, confirming that the isolates were fully virulent in both laboratory animals. The finding of a bovine isolate in New Zealand with characteristics indistinguishable from those of isolates from fur seals in New Zealand waters suggests that the seal bacillus is also capable of infecting cattle. This fact, combined with knowledge of its ability to cause disease in humans (Thompson *et al.*, 1993) and tapirs, suggests that the seal bacillus has the potential for a host range that extends beyond those of *M. tuberculosis*, *M. africanum* and *M. microti*.

Mycolic acid analysis

HPLC chromatograms of isolates Au-1 and Au-2 demonstrated a single cluster pattern that was consistent with species of the *M. tuberculosis* complex (data not shown) as reported previously (Butler *et al.*, 1991), providing additional evidence that these organisms belonged to the *M. tuberculosis* complex.

Tests for the MPB70 antigen

All seal isolates were negative when tested for the MPB70 antigen, despite containing the *mpb70* gene. The MPB70 antigen is considered to be characteristic of *M. bovis* (Corner *et al.*, 1988; Liébana *et al.*, 1996) and can be demonstrated by dot-blot immunoperoxidase (Liébana

et al., 1996) or SDS-PAGE (Cousins, 1996; Alito *et al.*, 1999). In a previous study, >97 % of *M. tuberculosis* isolates, 100 % of *M. microti* isolates and 90 % of *M. africanum* isolates were negative for the MPB70 antigen, whereas >99 % of *M. bovis* isolates were positive (Liébana *et al.*, 1996). In this regard, the seal bacillus was more like other members of the *M. tuberculosis* complex than *M. bovis*. The reported presence of the MPB70 antigen in a single isolate of *M. microti* from an alpaca (Alito *et al.*, 1999) was contrary to the findings of Liébana *et al.* (1996), who tested seven isolates of *M. microti* (including the reference strain).

16S rDNA sequence determination

16S rDNA sequencing is an accepted method of confirming the species designation of mycobacterial isolates (Böddinghaus *et al.*, 1990; Rogall *et al.*, 1990a, b). The 16S rDNA consensus strand (1400 nt) from the seal isolates demonstrated 99·9 % similarity to those of *M. tuberculosis* (GenBank number X58890) and *M. bovis* (available from the Sanger website at http://www.sanger.ac.uk/Projects/M_bovis/). A single nucleotide substitution (C→T) in the consensus strand occurred at *E. coli* position 1256 (data not shown). Phylogenetic analysis demonstrated that the consensus sequence was on the same branch as that of *M. tuberculosis*. The 16S rDNA regions of the isolates that were sequenced (1030 bp) were consistent with the sequence of the *M. tuberculosis* complex.

PCR-based testing of genetic markers

The *gyrA* and *katG* gene sequences of all seal isolates were identical (Table 4). These genetic markers are accepted methods of confirming that isolates belong to the *M. tuberculosis* complex (Thierry *et al.*, 1990, 1993;

Table 4. Genetic properties of the seal bacillus, compared to other members of the *M. tuberculosis* complex

Species: 1, *M. tuberculosis* (classic); 2, *M. tuberculosis* (Asian); 3, *M. africanum* (type I); 4, *M. africanum* (type II); 5, *M. microti*; 6, seal bacillus; 7, *M. bovis*; 8, '*M. caprae*'; 9, *M. bovis* BCG. All species contain IS6110 and IS1081, although some Asian strains of *M. tuberculosis* lack IS6110. Present (1–5), between one and five of the 3' spacers (39–43) are present; NIL, none of the five 3' spacers (39–43) are present.

Locus	1	2	3	4	5	6	7	8	9
<i>mtp40</i>	+*	+	+*	+*	–†	+	–	–	–
<i>pncA</i> C57	CAC (His)	CAC	CAC	CAC	CAC	CAC	GAC (Asp)	CAC	GAC
<i>katG</i> C463	CTG (Leu), CGG (Arg)	CTG/ CGG	CTG	CTG	CTG	CTG	CTG	CTG	CTG
<i>oxyR</i> nt 285	G	G	G	G	G	G	A	A	A
<i>gyrA</i> C95	AGC (Ser), ACC (Thr)	–	ACC	ACC	ACC	ACC	ACC	ACC	ACC
Spoligotyping: spacers 39–43	Present (1–5)		Present (1–5)		NIL	NIL	NIL	NIL	NIL

*Very occasionally, members of these species lack the *mtp40* gene (Liébana *et al.*, 1996).
†Seven of seven isolates (100 %) were negative for *mtp40* (Liébana *et al.*, 1996), whereas one isolate tested by Bernardelli *et al.* (1996) was reported as positive.

[illegible]

Collins & Stephens, 1991; Cousins *et al.*, 1991; Groenen *et al.*, 1993; Liébana *et al.*, 1996; Gormley *et al.*, 1997; Sreevatsan *et al.*, 1997). Results from sequencing of the *mtp40*, *pncA* and *oxyR* genes clearly demonstrated that the seal isolates were genetically more consistent with *M. tuberculosis* and *M. africanum* than with *M. bovis*.

Four different spoligotypes were identified in the seal isolates; all lacked the spacers 39–43, which are known to be characteristic of *M. bovis* (Fig. 1). All of the isolates from Australia and all but one of the Argentinian isolates had a unique but identical pattern, designated seal spoligotype 1 (SS-1). The remaining Argentinian isolate was designated SS-2. The three isolates from Great Britain had identical spoligotypes (SS-3) that differed by one spacer from the other seal spoligotypes. The seal isolates from New Zealand and the isolate from a New Zealand bovine had identical spoligotypes (SS-4) that lacked six spacers that were present in all other seal isolates. When compared to reference (and representative) strains of *M. tuberculosis*, *M. africanum*, *M. microti*, *M. bovis*, '*M. canettii*' and '*M. caprae*', the seal isolates formed a distinct cluster within the *M. tuberculosis* complex (Fig. 2). Spoligotyping confirmed that the seal isolates from Australia, Argentina, Uruguay and Great Britain were closely related. The finding of three spoligotypes with only minor differences from 29 isolates that originated from these diverse geographical regions indicated a clonal relationship between these isolates, which in turn suggests that the infection may have originated from a single source as a relatively recent event. Considering that these cases were diagnosed over a period of more than 15 years and that many of these populations inhabit geographically separate territories, a more likely explanation is that the DR locus exhibits considerable genetic stability in the seal bacillus. The spoligotype identified in the New Zealand isolates clustered with those of the other seal isolates but was genetically further removed, confirming a closer relationship to the other seal isolates than to other members of the *M. tuberculosis* complex. Similar small differences in

et al., 1999; Anh *et al.*, 2000) and has been used to trace the global spread of this strain. It has also been proved to be useful in defining populations of *M. microti* (van Soolingen *et al.*, 1998) and, in this study, demonstrated its usefulness in defining the limited genetic diversity of the seal bacillus.

Dendrograms constructed by using GelCompar indicated there was a close relationship between all seal-related isolates. Other methods of typing these isolates, including RFLP analysis with IS1081, IS6110, DR, PGRS and pUCD and VNTR (variable number of tandem repeats) typing, confirmed these findings (data not shown). Many investigators accept that members of the *M. tuberculosis* complex may be represented along a continuum with major peaks that correspond to each of the designated species. It is probable that the seal bacillus has evolved from another *M. tuberculosis* complex organism and has found a unique niche in this marine host. A study by Behr & Small (1999) that identified deletion events in *M. bovis* BCG has elicited information on the evolution of BCG strains. Similar evolutionary insights into the origin of the seal bacillus have been gained by using comparative genomic technologies that were described previously (Brosch *et al.*, 2002; Mostowy *et al.*, 2002). In both studies, the seal strain was separated from classical *M. bovis* by at least six deletions. The seal bacillus has a similar number of deletions to *M. microti* and Brosch *et al.* (2002) suggest that, along with *M. microti* and '*M. canettii*', the seal bacillus contains a unique deletion. These deletion studies provide further evidence that the seal bacillus should be designated as a separate species within the *M. tuberculosis* complex.

FAFLP

All 10 seal isolates produced indistinguishable results. When compared to the published sequences of *M. tuberculosis* strains CDC1551 and H37Rv and *M. bovis* strain AN5, up to 18 highly polymorphic FAFLP markers for the rapid identification of the seal bacillus were identified. In these studies, three loci appeared to be unique to the seal bacillus, 12 were shared with *M. bovis* and three were shared with *M. tuberculosis*. Further studies that include some of these loci may result in the identification of species-specific markers that are potentially useful for the development of PCR-based diagnostics for the seal bacillus. The identical genotype of all seal isolates that were tested by FAFLP confirmed their close clonal relationship, which had been identified by spoligotyping. It also substantiated previous studies that used FAFLP, which suggested that this technique may play a role in discriminating between mycobacterial species, including members of the *M. tuberculosis* complex (Goulding *et al.*, 2000; Huys *et al.*, 2000).

Description of *Mycobacterium pinnipedii* sp. nov.

Mycobacterium pinnipedii (pin.ni.pe'di.i. N.L. gen. neut. n. *pinnipedii* of a pinniped, referring to the host animal from which the organism was first isolated).

Isolates can be recovered from the lung and associated lymph nodes of tuberculous pinnipeds, and occasionally from mesenteric lymph nodes and organs such as the liver. Acid/alcohol-fast, non-spore-forming, non-motile bacilli with loose cord formation. Growth is generally enhanced by sodium pyruvate and usually occurs within 3–6 weeks of incubation on egg-based media at 36–37 °C. Colonies are dysgonic, rough, flat and non-photochromogenic. Isolates are negative for nitrate reduction and generally negative for niacin accumulation; some isolates demonstrate low-to-medium reactions for niacin. Susceptible to 50 µg PZA ml⁻¹ and 1 µg thiophen-2-carboxylic acid hydrazide (TCH) ml⁻¹ (isolates have occasionally demonstrated resistance to 1 µg TCH ml⁻¹, but are susceptible to 10 µg ml⁻¹). Pathogenic in guinea pigs and rabbits; the apparent incidental infection of a human, bovine and tapir indicates that they may have a wide host range. All isolates contain the sequences IS6110, IS1081, *mpb70* and *mtp40*, yet fail to produce detectable MPB70 antigen. The *pncA* gene contains CAC (His) at codon 57 and the *oxyR* gene shows G at nt 285, similar to *M. tuberculosis*, *M. microti* and *M. africanum*. The seal isolate spoligotypes form a cluster that is clearly different from those of all other members of the *M. tuberculosis* complex. The isolates are susceptible to isoniazid, rifampicin, streptomycin, ethambutol and paraminosalicylic acid.

The type strain is 6482^T (=ATCC BAA-688^T=NCTC 13288^T).

ACKNOWLEDGEMENTS

This work received financial support from the Brucellosis and Tuberculosis Eradication Campaign and the Tuberculosis Freedom Assurance Program within Australia. We are grateful to Fundación Mundo Marino, San Clemente del Tuyú, Argentina, for their support and to Frank Haverkort for his assistance with susceptibility testing and supplying *M. tuberculosis* isolates. We thank Dr Hans G. Trüper for advice on the Latin name of the organism.

REFERENCES

- Ahmed, N., Bal, A., Khan, A. A. & 9 other authors (2002). Whole genome fingerprinting and genotyping of multiple drug resistant (MDR) isolates of *Pseudomonas aeruginosa* from endophthalmitis patients in India. *Infect Genet Evol* 1, 237–242.
- Ahmed, N., Alam, M., Abdul Maajeed, A., Asad Rahman, S., Cataldi, A., Cousins, D. & Hasnain, S. E. (2003). Genome sequence based, comparative analysis of the fluorescent amplified fragment length polymorphisms (FAFLP) of tubercle bacilli from seals provides molecular evidence for a new species within the *Mycobacterium tuberculosis* complex. *Infect Genet Evol* 2, 193–199.
- Alito, A., Romano, M. I., Bigi, F., Zumarraga, M. & Cataldi, A. (1999). Antigenic characterization of mycobacteria from South American wild seals. *Vet Microbiol* 68, 293–299.
- Anh, D. D., Borgdorff, M. W., Van, L. N., Lan, N. T., van Gorkom, T., Kremer, K. & van Soolingen, D. (2000). *Mycobacterium tuberculosis* Beijing genotype emerging in Vietnam. *Emerg Infect Dis* 6, 302–305.

- Aranaz, A., Liébana, E., Mateos, A. & 8 other authors (1996). Spacer oligonucleotide typing of *Mycobacterium bovis* strains from cattle and other animals: a tool for studying epidemiology of tuberculosis. *J Clin Microbiol* **34**, 2734–2740.
- Aranaz, A., Liébana, E., Gómez-Mampaso, E. & 8 other authors (1999). *Mycobacterium tuberculosis* subsp. *caprae* subsp. nov.: a taxonomic study of a new member of the *Mycobacterium tuberculosis* complex isolated from goats in Spain. *Int J Syst Bacteriol* **49**, 1263–1273.
- Aranaz, A., Cousins, D., Mateos, A. & Domínguez, L. (2003). Elevation of *Mycobacterium tuberculosis* subsp. *caprae* Aranaz *et al.* 1999 to species rank as *Mycobacterium caprae* comb. nov., sp. nov. *Int J Syst Evol Microbiol* (in press).
- Bastida, R., Loureiro, J., Quse, V., Bernardelli, A., Rodriguez, D. & Costa, E. (1999). Tuberculosis in a wild subantarctic fur seal from Argentina. *J Wildl Dis* **35**, 796–798.
- Behr, M. A. & Small, P. M. (1999). A historical and molecular phylogeny of BCG strains. *Vaccine* **17**, 915–922.
- Bernardelli, A., Loureiro, J., Costa, E., Cataldi, A., Bastida, R. & Michelis, H. (1994). Tuberculosis in fur seals and sea lions of the south western Atlantic coast. In *IUATLD Newsletter*, May 1994. Paris: International Union Against Tuberculosis and Lung Disease.
- Bernardelli, A., Bastida, R., Loureiro, J., Michelis, H., Romano, M. I., Cataldi, A. & Costa, E. (1996). Tuberculosis in sea lions and fur seals from the south-western Atlantic coast. *Rev Sci Tech* **15**, 985–1005.
- Bernardelli, A., Trovero, A., Loureiro, J., Quse, V. & Morcillo, N. (1999). Sensibilidad in vitro a tuberculostáticos de cepas del complejo *Mycobacterium tuberculosis* aisladas en mamíferos marinos. In *Abstracts of the XX Congresso Brasileiro de Microbiologia*, p. 427, abstract SM-057. October 1999, Salvador, Brazil (in Portuguese).
- Böddinghaus, B., Rogall, T., Flohr, T., Blöcker, H. & Böttger, E. C. (1990). Detection and identification of mycobacteria by amplification of rRNA. *J Clin Microbiol* **28**, 1751–1759.
- Brosch, R., Gordon, S. V., Marmiesse, M. & 12 other authors (2002). A new evolutionary scenario for the *Mycobacterium tuberculosis* complex. *Proc Natl Acad Sci U S A* **99**, 3684–3689.
- Butler, W. R., Jost, K. C., Jr & Kilburn, J. O. (1991). Identification of mycobacteria by high-performance liquid chromatography. *J Clin Microbiol* **29**, 2468–2472.
- Butler, W. R., Floyd, M. M., Silcox, V. & 9 other editors (1996). *Standardized Method for HPLC Identification of Mycobacteria*, pp. 3–22. Atlanta, GA: Centers for Disease Control and Prevention.
- Butler, W. R., Floyd, M. M., Silcox, V. & 9 other editors (1999). *Mycolic Acid Pattern Standards for HPLC Identification of Mycobacteria*, pp. 3–10. Atlanta, GA: Centers for Disease Control and Prevention.
- Castets, M., Rist, N. & Boisvert, H. (1969). La variété africaine du bacille tuberculeux humain. *Med Afr Noire* **16**, 321–322 (in French).
- Castro Ramos, M., Ayala, M., Errico, F. & Silvera, F. V. (1998). Aislamiento de *Mycobacterium bovis* en Pinnípedos *Otaria byronia* (*Lobo marino común*) en Uruguay. *Rev Med Vet* **79**, 197–200 (in Portuguese).
- Collins, D. M. & Stephens, D. M. (1991). Identification of an insertion sequence, IS1081, in *Mycobacterium bovis*. *FEMS Microbiol Lett* **83**, 11–15.
- Collins, C. H., Yates, M. D. & Grange, J. M. (1982). Subdivision of *M. tuberculosis* in five variants for epidemiological purposes: methods and nomenclature. *J Hyg Camb* **89**, 235–242.
- Corner, L. A., John, M., Bundesen, P. G. & Wood, P. R. (1988). Identification of *Mycobacterium bovis* isolates using a monoclonal antibody. *Vet Microbiol* **18**, 191–196.
- Cousins, D. V. (1995). Tuberculosis in seals in Australia. In *Proceedings of the Australian Veterinary Association Conference*, pp. 51–57. Melbourne, Australia: Australian Association of Veterinary Conservation Biologists (AAVCB).
- Cousins, D. V. (1996). *Molecular epidemiology and diagnosis of Mycobacterium bovis and M. bovis-like organisms causing tuberculosis*, pp. 1–317. PhD thesis, University of Western Australia.
- Cousins, D. V., Wilton, S. D. & Francis, B. R. (1991). Use of DNA amplification for the rapid identification of *Mycobacterium bovis*. *Vet Microbiol* **27**, 187–195.
- Cousins, D. V., Williams, S. N., Reuter, R., Forshaw, D., Chadwick, B., Coughran, D., Collins, P. & Gales, N. (1993). Tuberculosis in wild seals and characterisation of the seal bacillus. *Aust Vet J* **70**, 92–97.
- Cousins, D., Williams, S., Liébana, E., Aranaz, A., Bunschoten, A. E., Van Embden, J. & Ellis, T. (1998a). Evaluation of four DNA typing techniques in epidemiological investigations of bovine tuberculosis. *J Clin Microbiol* **36**, 168–178.
- Cousins, D. V., Skuce, R. A., Kazwala, R. R. & van Embden, J. D. A. (1998b). Towards a standardized approach to DNA fingerprinting of *Mycobacterium bovis*. International Union Against Tuberculosis and Lung Disease, Tuberculosis in Animals Subsection. *Int J Tuberc Lung Dis* **2**, 471–478.
- Del Portillo, P., Murillo, L. A. & Patarroyo, M. E. (1991). Amplification of a species-specific DNA fragment of *Mycobacterium tuberculosis* and its possible use in diagnosis. *J Clin Microbiol* **29**, 2163–2168.
- Edwards, U., Rogall, T., Blöcker, H., Emde, M. & Böttger, E. C. (1989). Isolation and direct complete nucleotide determination of entire genes. Characterization of a gene coding for 16S ribosomal RNA. *Nucleic Acids Res* **17**, 7843–7853.
- Espinosa de los Monteros, L. E., Galan, J. C., Gutierrez, M. & 8 other authors (1998). Allele-specific PCR method based on *pncA* and *oxyR* sequences for distinguishing *Mycobacterium bovis* from *Mycobacterium tuberculosis*: intraspecies *M. bovis pncA* sequence polymorphism. *J Clin Microbiol* **36**, 239–242.
- Floyd, M. M., Guthertz, L. S., Silcox, V. A., Duffey, P. S., Jang, Y., Desmond, E. P., Crawford, J. T. & Butler, W. R. (1996). Characterization of an SAV organism and proposal of *Mycobacterium triplex* sp. nov. *J Clin Microbiol* **34**, 2963–2967.
- Forshaw, D. & Phelps, G. R. (1991). Tuberculosis in a captive colony of pinnipeds. *J Wildl Dis* **27**, 288–295.
- Gormley, E., Sandall, L., Hong, C., Lawton, D. & Murray, A. (1997). Identification and differentiation of mycobacteria using the *P_{AN}* promoter sequence from *Mycobacterium paratuberculosis* as a DNA probe. *FEMS Microbiol Lett* **147**, 63–68.
- Goulding, J. N., Stanley, J., Saunders, N. & Arnold, C. (2000). Genome-sequence-based fluorescent amplified-fragment length polymorphism analysis of *Mycobacterium tuberculosis*. *J Clin Microbiol* **38**, 1121–1126.
- Grange, J. M. & Yates, M. D. (1994). *Guidelines for Speciation within the Mycobacterium tuberculosis Complex*. WHO/Zoon/94.174. Geneva: World Health Organization Veterinary Public Health Unit.
- Groenen, P. M. A., Bunschoten, A. E., van Soolingen, D. & van Embden, J. D. A. (1993). Nature of DNA polymorphism in the direct repeat cluster of *Mycobacterium tuberculosis*; application for strain differentiation by a novel typing method. *Mol Microbiol* **10**, 1057–1065.
- Hunter, J. E., Duignan, P. J., Dupont, C., Fray, L., Fenwick, S. G. & Murray, A. (1998). First report of potentially zoonotic tuberculosis in fur seals in New Zealand. *N Z Med J* **111**, 130–131.
- Huys, G., Rigouts, L., Chemlal, K., Portaels, F. & Swings, J. (2000). Evaluation of amplified fragment length polymorphism analysis

- for inter- and intraspecific differentiation of *Mycobacterium bovis*, *M. tuberculosis*, and *M. ulcerans*. *J Clin Microbiol* **38**, 3675–3680.
- Kamerbeek, J., Schouls, L., Kolk, A. & 8 other authors (1997). Simultaneous detection and strain differentiation of *Mycobacterium tuberculosis* for diagnosis and epidemiology. *J Clin Microbiol* **35**, 907–914.
- Karlson, A. G. & Lessel, E. F. (1970). *Mycobacterium bovis* nom. nov. *Int J Syst Bacteriol* **20**, 273–282.
- Kirschner, P., Springer, B., Vogel, U., Meier, A., Wrede, A., Kiekenbeck, M., Bange, F.-C. & Böttger, E. C. (1993). Genotypic identification of mycobacteria by nucleic acid sequence determination: report of a 2-year experience in a clinical laboratory. *J Clin Microbiol* **31**, 2882–2889.
- Liébana, E., Aranaz, A., Francis, B. & Cousins, D. (1996). Assessment of genetic markers for species differentiation within the *Mycobacterium tuberculosis* complex. *J Clin Microbiol* **34**, 933–938.
- Morcillo, N., Bernardelli, A., Chirico, M. C. & Dolmann, A. L. (2000). Colorimetric assay in comparison with MGIT system and the agar proportion method for determining MICs antimicrobial agents for *Mycobacterium tuberculosis* complex. In *Abstracts of the World Congress on Lung Health and 10th ERS Annual Congress*. September 2000, Florence, Italy.
- Mostowy, S., Cousins, D., Brinkman, J., Aranaz, A. & Behr, M. A. (2002). Genomic deletions suggest a phylogeny for the *Mycobacterium tuberculosis* complex. *J Infect Dis* **186**, 74–80.
- Niemann, S., Richter, E. & Rüscher-Gerdes, S. (2002). Biochemical and genetic evidence for the transfer of *Mycobacterium tuberculosis* subsp. *caprae* Aranaz *et al.* 1999 to the species *Mycobacterium bovis* Karlson and Lessel 1970 (Approved Lists 1980) as *Mycobacterium bovis* subsp. *caprae* comb. nov. *Int J Syst Evol Microbiol* **52**, 433–436.
- Pfyffer, G. E., Auckenthaler, R., van Embden, J. D. & van Soolingen, D. (1998). *Mycobacterium canettii*, the smooth variant of *M. tuberculosis*, isolated from a Swiss patient exposed in Africa. *Emerg Infect Dis* **4**, 631–634.
- Qian, L., Van Embden, J. D. A., van der Zanden, A. G. M., Weltevreden, E. F., Duanmu, H. & Douglas, J. T. (1999). Retrospective analysis of the Beijing family of *Mycobacterium tuberculosis* in preserved lung tissues. *J Clin Microbiol* **37**, 471–474.
- Rogall, T., Flohr, T. & Böttger, E. C. (1990a). Differentiation of *Mycobacterium* species by direct sequencing of amplified DNA. *J Gen Microbiol* **136**, 1915–1920.
- Rogall, T., Wolters, J., Flohr, T. & Böttger, E. C. (1990b). Towards a phylogeny and definition of species at the molecular level within the genus *Mycobacterium*. *Int J Syst Bacteriol* **40**, 323–330.
- Romano, M. I., Alito, A., Bigi, F., Fisanotti, J. C. & Cataldi, A. (1995). Genetic characterization of mycobacteria from South American wild seals. *Vet Microbiol* **47**, 89–98.
- Sreevatsan, S., Escalante, P., Pan, X. & 11 other authors (1996). Identification of a polymorphic nucleotide in *oxyR* specific for *Mycobacterium bovis*. *J Clin Microbiol* **34**, 2007–2010.
- Sreevatsan, S., Pan, X., Stockbauer, K. E., Connel, N. D., Kreiswirth, B. N., Whittam, T. S. & Musser, J. M. (1997). Restricted structural gene polymorphism in the *Mycobacterium tuberculosis* complex indicates evolutionarily recent global dissemination. *Proc Natl Acad Sci U S A* **94**, 9869–9874.
- Thierry, D., Brisson-Noël, A., Vincent-Levy-Frébault, V., Nguyen, S., Guesdon, J. L. & Gicquel, B. (1990). Characterization of a *Mycobacterium tuberculosis* insertion sequence, IS6110, and its application in diagnosis. *J Clin Microbiol* **28**, 2668–2673.
- Thierry, D., Matsiota-Bernard, P., Pitsouni, E., Costopoulos, C. & Guesdon, J. L. (1993). Use of the insertion element IS6110 for DNA fingerprinting of *Mycobacterium tuberculosis* isolates presenting various profiles of drug susceptibility. *FEMS Immunol Med Microbiol* **6**, 287–297.
- Thompson, P. J., Cousins, D. V., Gow, B. L., Collins, D. M., Williamson, B. W. & Dagnia, H. T. (1993). Seals, seal trainers, and mycobacterial infection. *Am Rev Respir Dis* **147**, 164–167.
- Tsukamura, M., Mizuno, S. & Toyama, H. (1985). Taxonomic studies on the *Mycobacterium tuberculosis* series. *Microbiol Immunol* **29**, 285–299.
- van Soolingen, D., Qian, L., de Haas, P. E. & 7 other authors (1995). Predominance of a single genotype of *Mycobacterium tuberculosis* in countries of east Asia. *J Clin Microbiol* **33**, 3234–3238.
- van Soolingen, D., Hoogenboezem, T., de Haas, P. E. & 9 other authors (1997). A novel pathogenic taxon of the *Mycobacterium tuberculosis* complex, Canetti: characterization of an exceptional isolate from Africa. *Int J Syst Bacteriol* **47**, 1236–1245.
- van Soolingen, D., van der Zanden, A. G. M., de Haas, P. E. W. & 7 other authors (1998). Diagnosis of *Mycobacterium microti* infections among humans by using novel genetic markers. *J Clin Microbiol* **36**, 1840–1845.
- Veerman, G. M., Kelman, R., Colley, J. & Pike, J. G. (1990). Rapid confirmatory identification of *Mycobacterium bovis* using a dot blotting immunodetection technique. *Vet Microbiol* **22**, 335–340.
- Vestal, A. L. (1975). *Procedures for the Isolation and Identification of Mycobacteria*, PHS no. 1547. Atlanta, GA: Centers for Disease Control.
- Wayne, L. G. (1984). Mycobacterial speciation. In *The Mycobacteria: a Sourcebook*, pp. 42–43. Edited by G. P. Kubica & L. G. Wayne. New York: Marcel Dekker.
- Wells, A. Q. & Oxen, D. M. (1937). Tuberculosis in wild voles. *Lancet* **i**, 1221.
- Wells, A. Q. & Robb-Smith, A. H. T. (1946). *The Murine Type of Tubercle Bacillus (the Vole Acid-fast Bacillus) with Notes on the Morphology of Infection by the Vole Acid-fast Bacillus*. London, UK: Medical Research Council.
- Woods, R., Cousins, D. V., Kirkwood, R. & Obendorf, D. L. (1995). Tuberculosis in a wild Australian fur seal (*Arctocephalus pusillus doriferus*) from Tasmania. *J Wildl Dis* **31**, 83–86.
- Zumárraga, M. J., Bernardelli, A., Bastida, R. & 10 other authors (1999). Molecular characterization of mycobacteria isolated from seals. *Microbiology* **145**, 2519–2526.