Erratum

Description of *Enterococcus canis* sp. nov. from dogs and reclassification of *Enterococcus porcinus* Teixeira et al. 2001 as a junior synonym of *Enterococcus villorum* Vancanneyt et al. 2001

E. M. De Graef, L. A. Devriese, M. Vancanneyt, M. Baele, M. D. Collins, K. Lefebvre, J. Swings and F. Haesebrouck

International Journal of Systematic and Evolutionary Microbiology (2003) vol. 53, part 4, pp. 1069–1074

On p. 1073, towards the end of the description of *Enterococcus canis* sp. nov., the text should read as follows. 'Strains are negative in API tests for hippurate, β -glucuronidase, β -galactosidase and alkaline phosphatase, and positive in Crystal tests for hydrolysis of L-valine-AMC, 4MU-phosphate and L-isoleucine-AMC. Urease and 4MU- β -D-glucuronide are variable in Crystal tests. Acid is not produced from erythritol, L-xylose, adonitol, L-sorbose, rhamnose, dulcitol, sorbitol, α -methyl-D-mannoside, melibiose, inulin, melezitose, D-raffinose, glycogen, xylitol, D-turanose, D-lyxose, D- and L-fucose, D-tagatose, D- and L-arabitol, and 5-ketogluconate.'