Vibrio crassostreae sp. nov., isolated from the haemolymph of oysters (*Crassostrea gigas*)

N. Faury,¹ D. Saulnier,¹ F. L. Thompson,² M. Gay,¹ J. Swings² and F. Le Roux¹

Correspondence

F. Le Roux

fleroux@ifremer.fr

¹Laboratoire de Génétique et Pathologie, Institut français de recherche pour l'exploitation de la mer (Ifremer), 17390 La Tremblade, France

²Laboratory for Microbiology and BCCMTM/LMG Bacteria Collection, Laboratory for Microbiology, Ghent University, K.L. Ledeganckstraat 35, Ghent 9000, Belgium

Polyphasic analysis of five new *Vibrio* isolates originating from the haemolymph of diseased cultured oysters is described. The new isolates were closely related to *Vibrio splendidus*, having 98 % 16S rRNA gene sequence similarity. *gyrB* phylogenetic analysis, fluorescent amplified-fragment length polymorphism (FAFLP) fingerprinting and DNA–DNA hybridization experiments clearly showed that the new isolates form a tight genomic group that is different from the currently known *Vibrio* species. It is proposed to accommodate these isolates in a novel species, *Vibrio crassostreae* sp. nov. (type strain LGP 7^T = LMG 22240^T = CIP 108327^T). Phenotypic and chemotaxonomic features that differentiate *V. crassostreae* from other known *Vibrio* species include arginine dihydrolase, utilization and fermentation of various carbon sources, β -galactosidase activity, NO₂ production and the presence of the fatty acids 14 : 0 iso and 16 : 0 iso.

Vibrio splendidus-related species have been associated with mortality of molluscs and fish (Nicolas *et al.*, 1996; Sugumar *et al.*, 1998; Gatesoupe *et al.*, 1999; Lacoste *et al.*, 2001; Waechter *et al.*, 2002; Farto *et al.*, 2003). Epidemiological studies of *V. splendidus* strains associated with oyster mortality outbreaks have demonstrated a high genetic diversity within this group and suggested its polyphyletic nature (Le Roux *et al.*, 2002, 2004). Six species, *Vibrio lentus*, *Vibrio kanaloae*, *Vibrio pomeroyi*, *Vibrio tasmaniensis*, *Vibrio chagasii* and *V. splendidus*, have been described within this group so far, but there exist only a limited number of diagnostic biochemical features that allow clear species discrimination within this group (Macián *et al.*, 2001; Thompson *et al.*, 2003a, b). In a previous study we investigated a collection of *V. splendidus*-related isolates originated from the haemolymph of oysters that are potentially pathogenic for the oyster *Crassostrea gigas* (Gay *et al.*, 2004). These strains were characterized by DNA gyrase subunit B (*gyrB*) gene sequencing (Le Roux

et al., 2004). Several strains clustered together but could not be assigned to any known *Vibrio* species. In the present study we present a detailed polyphasic analysis of a group of five *Vibrio* isolates, including 16S rRNA and *gyrB* gene sequencing, fluorescent amplified-fragment length polymorphism (FAFLP) fingerprinting, DNA–DNA hybridizations and biochemical tests. Collectively, the results suggest that the five isolates belong to a novel species, for which we propose the name *Vibrio crassostreae* sp. nov.

The strains used in this study were purchased from national collections (*V. splendidus* LMG 4042^T, *V. tasmaniensis* LMG 20012^T, *V. kanaloae* LMG 20539^T, *V. pomeroyi* LMG 20537^T, *Vibrio cyclitrophicus* LMG 21359^T, *V. chagasii* LMG 21353^T, *V. lentus* CIP 107166^T) or isolated from the haemolymph of cultured *C. gigas* in La Tremblade (France) [LMG 22240^T (=LGP 7^T = CIP 108327^T), LMG 22241 (=LGP 8 = CIP 108328), LMG 22248 (=LGP 15 = CIP 108329), LMG 22249 (=LGP 107 = CIP 108330), LMG 22242 (=LGP 17), LMG 22243 (=LGP 1) and LGP 108] and deposited in the BCCM/LMG Bacteria Collection (Gent, Belgium) and in the Institut Pasteur Bacteria Collection (CIP; Paris, France). All strains were cultured on tryptone soy agar (TSA; Oxoid) supplemented with 2 % (w/v) NaCl for 48 h at 20 °C.

PCR amplification, cloning and sequencing of the 16S rRNA and *gyrB* gene fragments were done as described previously (Yamamoto & Harayama, 1995; Lambert *et al.*, 1998; Le Roux *et al.*, 2004). Sequences were aligned and phylogenetic analyses were performed with SEAVIEW and

Published online ahead of print on 21 May 2004 as DOI 10.1099/ij.s.0.63232-0.

Abbreviation: FAFLP, fluorescent amplified-fragment length polymorphism.

The GenBank/EMBL/DDBJ accession numbers for the 16S rRNA and *gyrB* gene sequences of LGP 7^T are AJ582808 and AJ582799, respectively.

Tables of supplementary DNA–DNA hybridization, phenotypic and fatty acid data are available in IJSEM Online.

PHYLO_WIN software (Galtier *et al.*, 1996). Phylogenetic trees were constructed using neighbour-joining, maximum-likelihood and maximum-parsimony. For neighbour-joining analysis, distance matrices were calculated by using Kimura's 2-parameter distances (Gascuel, 1997). Reliability of topologies was assessed by the bootstrap method with 1000 replicates. FAFLP analysis was carried out as described previously (Thompson *et al.*, 2001).

For DNA–DNA hybridization experiments, *in vitro* labelling of the DNA with tritium-labelled nucleotides was performed by the random primer method (Megaprime labelling kit; Amersham) and hybridization was carried out at 60 °C by the S1-nuclease method (Crosa *et al.*, 1973; Grimont *et al.*, 1980) with adsorption of S1-resistant DNA onto Whatman DE81 filters.

Phenotypic characterization of the strains was done using the following commercially available kits: the Gram kit (bioMérieux), oxidase (Bactident oxidase; Merck), respiratory activity (meat liver medium; Diagnostic Pasteur), glucose metabolism (MEVAG; Diagnostic Pasteur), API 20E and API 50CH (bioMérieux) with the modification suggested by MacDonell *et al.* (1982), namely 2 % NaCl was added to the bacterial suspension. Motility, NaCl requirement and tolerance (0, 2, 4, 6, 8 and 10 %, w/v) and temperature tolerance (4, 20, 35 and 40 °C) were tested in 1.5 % (w/v) peptone broth (Diagnostic Pasteur). Numerical analysis of phenotypic features was performed using simple matching coefficients (Sneath, 1972) and the unweighted pair group method (Sneath & Sokal, 1973). Sensitivity to O/129 (150 µg per disc) was determined with Oxoid discs. Fatty acid methyl ester analysis was carried out as described by Huys *et al.* (1994).

The phylogenetic tree based on the *gyrB* nucleotide sequences (1064 gap-free sites long) confirmed the clustering of *V. crassostreae* sp. nov. strains LGP 7^T, LGP 8, LGP 15, LGP 107 and LGP 108 with a bootstrap value of 100 % and their distinction from their closest phylogenetic neighbours *V. cyclitrophicus*, *V. lentus*, *V. pomeroyi*, *V. kanaloae*, *V. tasmaniensis* and *V. splendidus* (Fig. 1). The phylogenetic tree based on almost-complete sequences of the 16S rRNA gene does not allow clear differentiation of the two representative isolates (LGP 7^T and LGP 8) from other species phenotypically related to *V. splendidus* (Fig. 2). Results are in accordance with previous studies, showing that the 16S rRNA gene sequences of *V. splendidus*-related strains are very similar (Macián *et al.*, 2001; Le Roux *et al.*, 2002, 2004; Thompson *et al.*, 2003a, b). Similar results were obtained by maximum-parsimony and maximum-likelihood analyses (data not shown).

The FAFLP patterns of four representative *V. crassostreae* strains consisted of 125 bands (± 9 SD). The mutual FAFLP pattern similarity among these strains was at least 86 %. Strains of *V. crassostreae* sp. nov. were clearly differentiated from all the other currently known species of the *Vibrionaceae*, *V. splendidus* and *V. kanaloae* being the most closely

Fig. 1. Phylogenetic tree of partial *gyrB* sequences. The *Vibrio campbellii* homologue was used as the outgroup; 1064 gap-free sites were compared. Horizontal branch lengths are proportional to evolutionary divergence. Bootstrap percentages from 1000 replicates appear next to the corresponding branch.

related, with 54 % similarity (Fig. 3). The FAFLP data suggest that the isolates indeed belong to a novel *Vibrio* species.

DNA–DNA hybridization experiments confirmed the grouping found with FAFLP and *gyrB*. *V. crassostreae* strains had at least 78 % DNA–DNA relatedness, but at maximum 61 % towards five other *V. splendidus*-related species (Supplementary Table A in IJSEM Online).

Strains of *V. crassostreae* sp. nov. could be differentiated from their closest phylogenetic neighbours by 17 phenotypic characters analysed in this study (Supplementary Table B). Those tests were coded as 1 (positive result) or 0 (negative result) and numerical analysis was performed using simple matching coefficients (Fig. 4). *V. crassostreae* sp. nov. has the main fatty acid traits of vibrios, i.e. 16:1 ω 7c and/or 15:0 iso 2-OH, 18:1 ω 7c, 12:0, 14:0, 12:0 3-OH (Supplementary Table C). The fatty acids 16:0 iso and

Fig. 2. Phylogenetic tree of partial 16S rRNA gene sequences; 1200 gap-free sites were compared. Other features as in Fig. 1.

Fig. 3. Dendrogram (Dice, Ward) of the FAFLP band patterns of *V. crassostreae* sp. nov. and all other known *V. splendidus*-related species.

14:0 iso seem to be useful for discriminating among *V. splendidus*-related species.

The present study illustrates the use of the *gyrB*-based phylogenetic structure in an interim period to cluster strains before validation of species affiliation by DNA–DNA hybridization and description of phenotypic features. However, in the case of *V. pomeroyi*, *V. kanaloae* and *V. tasmaniensis*, our *gyrB*-based analysis appears to be less discriminatory than DNA–DNA hybridization or FAFLP fingerprinting (Thompson *et al.*, 2001). Sequence analyses of other loci, including *rpoD* (Yamamoto & Harayama, 1998) and *hsp60* sequences (Kwok *et al.*, 2002), are in progress in order to assess the usefulness of such genes to discriminate *V. splendidus*-related species.

Description of *Vibrio crassostreae* sp. nov.

Vibrio crassostreae (cra.sso.stre'ae. N.L. gen. n. *crassostreae* of *Crassostrea gigas*, the oyster species from which the strains were isolated).

Cells are Gram-negative, curved, 1 µm wide and 2–3 µm

Fig. 4. Cluster analysis of phenetic data for *Vibrio* species using simple matching-similarity coefficient and unweighted pair group method (expressed as a percentage on the scale bar) for 12 strains related to *V. splendidus*.

long. Cells are motile by at least one polar flagellum. Forms translucent, non-swarming, rounded colonies with entire margins on TSA. Strains form yellow, translucent, 5 mm colonies on thiosulfate-citrate-bile salts-sucrose (TCBS) agar. Cells grow at 4 °C. None of the strains grows at 0 or 8 % NaCl. All strains are β -galactosidase-negative, arginine dihydrolase- and gelatinase-positive. Oxidase- and catalase-positive and urease-negative. Facultatively anaerobic and produces NO_2 . The following compounds are utilized as sole carbon sources: glucose, sucrose, melibiose, amygdalin, glycerol, ribose, galactose, D-mannose, mannitol, *N*-acetylglucosamine, aesculin, cellobiose, starch, glycogen and L-fucose. None of the strains utilize inositol, rhamnose, arabinose, erythritol, D- or L-arabinose, D- or L-xylose, adonitol, methyl β -D-xyloside, L-sorbose, dulcitol, sorbitol, methyl α -D-mannoside, methyl α -D-glucoside, amygdalin, arbutin, salicin, lactose, inulin, melezitose, D-raffinose, xylitol, β -gentiobiose, D-turanose, D-lyxose, D-tagatose, D-fucose, D- or L-arabitol, 2-oxoglutarate or 5-oxoglutarate. All strains are sensitive to O/129. Major fatty acids are summed feature 3 ($39.4 \pm 1\%$, comprising 16:1 ω 7c and/or 15:0 iso 2-OH), 16:0 ($17.3 \pm 1.3\%$), 16:0 iso ($8.7 \pm 0.2\%$), 18:1 ω 7c ($7 \pm 1.3\%$), 12:0 ($5.5 \pm 0.3\%$), 14:0 ($5.4 \pm 1\%$), 12:0 3-OH ($3.3 \pm 0.2\%$), summed feature 2 ($2.6 \pm 0.1\%$, comprising 14:0 3-OH and/or 16:1 iso 1 and/or unidentified fatty acid with equivalent chain-length value of 10.928 and/or 12:0 ALDE), 14:0 iso (1.6%), 17:0 (1.4%), 17:1 ω 8c (1.3%) and 14:0 iso 3-OH (1.2%).

Type strain LGP 7^T (=LMG 22240^T=CIP 108327^T) was isolated from a diseased oyster (*Crassostrea gigas*) at the laboratoire de genetique et pathologie (Ifremer, France). Reference strains are LMG 22241 (=LGP 8=CIP 108328), LMG 22248 (=LGP 15=CIP 108329), LMG 22249 (=LGP 107=CIP 108330) and LGP 108. The GenBank/EMBL/DBJ accession numbers for the 16S and *gyrB* gene sequences of LGP 7^T are AJ582808 and AJ582799, respectively.

Acknowledgements

This study was carried out with financial assistance from the French Research and Technological Ministry (CRB: Collection de Ressources Biologique) and the European Community Reference Laboratory for Bivalve Mollusc Diseases. F. L. T. acknowledges a postdoctoral grant from the BCCMTM/LMG Bacteria Collection, Belgium. J. S. acknowledges grants from the Fund for Scientific Research (FWO), Belgium.

References

- Crosa, J. H., Brenner, D. J. & Falkow, S. (1973). Use of a single-strand specific nuclease for analysis of bacterial and plasmid deoxyribonucleic acid homo- and heteroduplexes. *J Bacteriol* **115**, 904–911.
- Farto, R., Armada, S. P., Montes, M., Guisande, J. A., Perez, M. J. & Nieto, T. P. (2003). *Vibrio lentus* associated with diseased wild octopus (*Octopus vulgaris*). *J Invertebr Pathol* **83**, 149–156.

- Galtier, N., Gouy, M. & Gautier, C. (1996). SEAVIEW and PHYLO_WIN: two graphic tools for sequence alignment and molecular phylogeny. *Comput Appl Biosci* **12**, 543–548.
- Gascuel, O. (1997). BIONJ: an improved version of the NJ algorithm based on a simple model of sequence data. *Mol Biol Evol* **14**, 685–695.
- Gatesoupe, F. J., Lambert, C. & Nicolas, J. L. (1999). Pathogenicity of *Vibrio splendidus* strains associated with turbot larvae, *Scophthalmus maximus*. *J Appl Microbiol* **87**, 757–763.
- Gay, M., Berthe, F. C. J. & Le Roux, F. (2004). Screening of *Vibrio* isolates to develop an experimental infection model in the Pacific oyster *Crassostrea gigas*. *Dis Aquat Organ* **59**, 49–56.
- Grimont, P. A. D., Popoff, M. Y., Grimont, F., Coynault, C. & Lemelin, M. (1980). Reproducibility and correlation study of three deoxyribonucleic acid hybridisation procedures. *Curr Microbiol* **4**, 325–330.
- Huys, G., Vancanneyt, M., Coopman, R., Janssen, P., Falsen, E., Altwegg, M. & Kersters, K. (1994). Cellular fatty acid composition as a chemotaxonomic marker for the differentiation of phenospecies and hybridization groups in the genus *Aeromonas*. *Int J Syst Bacteriol* **44**, 651–658.
- Kwok, A. Y., Wilson, J. T., Coulthart, M., Ng, L. K., Mutharia, L. & Chow, A. W. (2002). Phylogenetic study and identification of human pathogenic *Vibrio* species based on partial *hsp60* gene sequences. *Can J Microbiol* **48**, 903–910.
- Lacoste, A., Jalabert, F., Malham, S., Cueff, A., Gélébart, F., Cordevant, C., Lange, M. & Poulet, S. A. (2001). A *Vibrio splendidus* strain is associated with summer mortality of juvenile oysters *Crassostrea gigas* in the Bay of Morlaix (North Brittany, France). *Dis Aquat Organ* **46**, 139–145.
- Lambert, C., Nicolas, J. L., Cilia, V. & Corre, S. (1998). *Vibrio pectenica* sp. nov., a pathogen of scallop (*Pecten maximus*) larvae. *Int J Syst Bacteriol* **48**, 481–487.
- Le Roux, F., Gay, M., Lambert, C., Waechter, M., Poubalanne, S., Chollet, B., Nicolas, J. L. & Berthe, F. C. J. (2002). Comparative analysis of *Vibrio splendidus*-related strains isolated during *Crassostrea gigas* mortality events. *Aquat Living Resour* **15**, 251–258.
- Le Roux, F., Gay, M., Lambert, C., Nicolas, J. L., Gouy, M. & Berthe, F. C. J. (2004). Phylogenetic study and identification of *Vibrio splendidus*-related strains based on *gyrB* gene sequences. *Dis Aquat Organ* **58**, 143–150.
- MacDonell, M. T., Singleton, F. L. & Hood, M. A. (1982). Diluent composition for use of API 20E in characterizing marine and estuarine bacteria. *Appl Environ Microbiol* **44**, 423–427.
- Macián, M. C., Ludwig, W., Aznar, R., Grimont, P. A. D., Schleifer, K. H., Garay, E. & Pujalte, M. J. (2001). *Vibrio lentus* sp. nov., isolated from Mediterranean oysters. *Int J Syst Evol Microbiol* **51**, 1449–1456.
- Nicolas, J. L., Corre, S., Gauthier, G., Robert, R. & Ansquer, D. (1996). Bacterial problems associated with scallop *Pecten maximus* larval culture. *Dis Aquat Organ* **27**, 67–76.
- Sneath, P. H. A. (1972). Computer taxonomy. *Methods Microbiol* **7**, 29–98.
- Sneath, P. H. A. & Sokal, R. R. (1973). *Numerical Taxonomy: the Principles and Practice of Numerical Classification*. San Francisco: W. H. Freeman.
- Sugumar, G., Nakai, T., Hirata, Y., Matsubara, D. & Muroga, K. (1998). *Vibrio splendidus* biovar II as the causative agent of bacillary necrosis of Japanese oyster *Crassostrea gigas* larvae. *Dis Aquat Organ* **33**, 111–118.
- Thompson, F. L., Hoste, B., Vandemeulebroecke, K. & Swings, J. (2001). Genomic diversity amongst *Vibrio* isolates from different sources determined by fluorescent amplified fragment length polymorphism. *Syst Appl Microbiol* **24**, 520–538.
- Thompson, F. L., Thompson, C. C., Li, Y., Gomez-Gil, B., Vandenberghe, J., Hoste, B. & Swings, J. (2003a). *Vibrio kanaloae* sp. nov., *Vibrio pomeroyi* sp. nov. and *Vibrio chagasii* sp. nov., from sea water and marine animals. *Int J Syst Evol Microbiol* **53**, 753–759.
- Thompson, F. L., Thompson, C. C. & Swings, J. (2003b). *Vibrio tasmaniensis* sp. nov., isolated from Atlantic salmon (*Salmo salar* L.). *Syst Appl Microbiol* **26**, 65–69.
- Waechter, M., Le Roux, F., Nicolas, J. L., Marissal, E. & Berthe, F. (2002). Characterization of pathogenic bacteria of the cupped oyster *Crassostrea gigas*. *C R Biol* **325**, 231–238 (in French).
- Yamamoto, S. & Harayama, S. (1995). PCR amplification and direct sequencing of *gyrB* genes with universal primers and their application to the detection and taxonomic analysis of *Pseudomonas putida* strains. *Appl Environ Microbiol* **61**, 1104–1109.
- Yamamoto, S. & Harayama, S. (1998). Phylogenetic relationships of *Pseudomonas putida* strains deduced from the nucleotide sequences of *gyrB*, *rpoD* and 16S rRNA genes. *Int J Syst Bacteriol* **48**, 813–819.