Sphingomonas dokdonensis sp. nov., isolated from soil

Jung-Hoon Yoon, Mi-Hwa Lee, So-Jung Kang, Soo-Young Lee and Tae-Kwang Oh

Correspondence

Jung-Hoon Yoon
jhyoon@kribb.re.kr

Tae-Kwang Oh
otk@kribb.re.kr

Korea Research Institute of Bioscience and Biotechnology (KRIBB), PO Box 115, Yusong, Taejeon, Korea

A Gram-negative, rod-shaped, *Sphingomonas*-like bacterial strain, DS-4^T, was isolated from soil of Dokdo, Korea, and its taxonomic position was investigated using a polyphasic approach. Strain DS-4^T grew optimally on trypticase soy agar medium without NaCl at pH 6.0–6.5 and 25 °C. It contained Q-10 as the predominant ubiquinone and C_{18:1}ω7c, C_{16:0}, C_{14:0} 2-OH and C_{16:1}ω7c and/or iso-C_{15:0} 2-OH as the major fatty acids. Sphingoglycolipid, phosphatidylcholine, phosphatidylglycerol, diphosphatidylglycerol, phosphatidylethanolamine and unidentified phospholipid were the major polar lipids. The DNA G + C content was 66.9 mol%. Phylogenetic analysis based on 16S rRNA gene sequences showed that strain DS-4^T fell within the evolutionary radiation comprising *Sphingomonas* species. Levels of 16S rRNA gene sequence similarity between strain DS-4^T and the type strains of *Sphingomonas* species ranged from 93.0 to 97.6%. DNA–DNA relatedness data and differential phenotypic properties, together with the phylogenetic distinctiveness, demonstrated that strain DS-4^T differs from the recognized *Sphingomonas* species. On the basis of phenotypic, phylogenetic and genetic data, this strain represents a novel species of the genus *Sphingomonas*, for which the name *Sphingomonas dokdonensis* sp. nov. is proposed, with DS-4^T (=KCTC 12541^T = CIP 108841^T) as the type strain.

Dokdo is an island located at the edge of the East Sea, Korea, and restriction of access to the island for a long time may have resulted in an interesting microbial diversity, which led us to investigate the microbial community of soils on the island. In an attempt to investigate the microbial community of Dokdo, many bacterial strains have been isolated and characterized taxonomically (Yoon *et al.*, 2005, 2006). In this study, we report on the detailed taxonomic characterization of a yellow-pigmented *Sphingomonas*-like bacterial strain, DS-4^T, which was considered to belong to the class *Proteobacteria* from the results of 16S rRNA gene sequence analysis.

Soil samples collected in Dokdo (37° 14' 12" N 131° 52' 07" E) provided the source for isolation of bacterial strains. Strain DS-4^T was isolated by the standard dilution plating technique at 25 °C on nutrient agar (Difco). The type strains of six *Sphingomonas* species were used as reference strains for DNA–DNA hybridization. *Sphingomonas aquatilis* KCTC 2881^T, *Sphingomonas asaccharolytica* KCTC 2825^T, *Sphingomonas koreensis* KCTC 2882^T, *Sphingomonas mali* KCTC 2826^T and *Sphingomonas pruni* KCTC 2824^T were obtained from the Korean Collection for Type Cultures (KCTC), Taejeon, Korea. *Sphingomonas melonis* DSM

14444^T was obtained from the Deutsche Sammlung von Mikroorganismen und Zellkulturen (DSMZ), Braunschweig, Germany.

Morphological, physiological and biochemical characteristics of strain DS-4^T were investigated using routine cultivation at 25 °C on trypticase soy agar (TSA; Difco) with the pH adjusted to 6.5. Cell morphology was examined by light microscopy (Nikon E600) and transmission electron microscopy. The presence of flagella was determined by transmission electron microscopy using cells from exponentially growing cultures. For transmission electron microscopic observation, cells were negatively stained with 1% (w/v) phosphotungstic acid and after air-drying the grids were examined with a Philips CM-20 transmission electron microscope. The Gram reaction was determined using the bioMérieux Gram stain kit according to the manufacturer's instructions. Growth at various temperatures (4–40 °C) was measured on TSA (pH 6.5). Growth in the absence of NaCl and growth at various NaCl concentrations [0.5% (w/v) and 1.0–10.0% (w/v) at intervals of 1.0%] were investigated in trypticase soy broth prepared according to the formula of the Difco medium except that no NaCl was used. The pH range for growth was determined in nutrient broth (Difco) adjusted to various pH values (pH 4.5–10.5 at intervals of 0.5). The pH was adjusted to various levels by the addition of HCl or Na₂CO₃ prior to sterilization. Growth under

The GenBank/EMBL/DDBJ accession number for the 16S rRNA gene sequence of strain DS-4^T is DQ178975.

anaerobic conditions was determined after incubation in an anaerobic chamber on TSA (pH 6.5) and on TSA (pH 6.5) supplemented with nitrate, both of which had been prepared anaerobically using nitrogen. Catalase and oxidase activities and hydrolysis of casein, gelatin, hypoxanthine, starch, Tween 20, Tween 40, Tween 60 and Tween 80, tyrosine, urea and xanthine were determined as described by Cowan & Steel (1965). Hydrolysis of aesculin and nitrate reduction were studied as described by Lányi (1987). Utilization of substrates as sole carbon and energy sources was tested according to the method of Yurkov *et al.* (1994) and by using the API 20NE system (bioMérieux). Sensitivity to antibiotics was tested on TSA (pH 6.5) plates using antibiotic discs containing the following: polymyxin B, 100 U; streptomycin, 50 µg; penicillin G, 20 U; chloramphenicol, 100 µg; ampicillin, 10 µg; cephalothin, 30 µg; gentamicin, 30 µg; novobiocin, 5 µg; tetracycline, 30 µg; kanamycin, 30 µg; lincomycin, 15 µg; oleandomycin, 15 µg; neomycin, 30 µg; carbenicillin, 100 µg. Enzyme activity and other physiological properties were tested using the API ZYM and API 20E systems (bioMérieux).

Cell biomass for DNA extraction and for isoprenoid quinone and polar lipid analyses was obtained from cultures grown in trypticase soy broth (Difco) at 25 °C. Chromosomal DNA was extracted and purified according to the method described by Yoon *et al.* (1996), with the exception that RNase T1 was used in combination with RNase A to minimize contamination with RNA. The 16S rRNA gene was amplified by PCR using two universal primers as described previously (Yoon *et al.*, 1998). Sequencing of the amplified 16S rRNA gene and phylogenetic analysis were performed as described by Yoon *et al.* (2003). The DNA G+C content was determined by the method of Tamaoka & Komagata (1984) with the modification that DNA was hydrolysed and the resultant nucleotides were analysed by reversed-phase HPLC. For fatty acid methyl ester analysis, the cell mass of strain DS-4^T was harvested from TSA (pH 6.5) plates after incubation for 4 days at 25 °C. Isoprenoid quinones were extracted according to the method of Komagata & Suzuki (1987) and analysed using reversed-phase HPLC and a YMC ODS-A (250 × 4.6 mm) column. Polar lipids were extracted according to the procedures described by Minnikin *et al.* (1984) and identified by two-dimensional TLC followed by spraying with appropriate detection reagents (Minnikin *et al.*, 1984; Komagata & Suzuki, 1987). The presence of phosphatidylcholine was identified by spraying with Dragendorff's reagent. DNA–DNA hybridization was performed fluorometrically by the method of Ezaki *et al.* (1989) using photobiotin-labelled DNA probes and microdilution wells. Hybridization was performed with five replications for each sample. The highest and lowest values obtained in each sample were excluded and the means of the remaining three values were quoted as DNA–DNA relatedness values.

The morphological, cultural, physiological and biochemical characteristics of strain DS-4^T are given in the species

description (see later) and are shown in Table 1. The almost complete 16S rRNA gene sequence (1441 nt) of strain DS-4^T was determined in this study. In the phylogenetic tree based on the neighbour-joining algorithm, strain DS-4^T fell within the radiation of the cluster comprising *Sphingomonas* species (Fig. 1). The phylogenetic affiliation of strain DS-4^T to the genus *Sphingomonas* was also recovered in trees based on the maximum-likelihood and maximum-parsimony algorithms (data not shown). Levels of 16S rRNA gene sequence similarity between strain DS-4^T and the type strains of *Sphingomonas* species ranged from 93.0% (*Sphingomonas taejonensis* JSS54^T) to 97.6% (*Sphingomonas asaccharolytica* IFO 15499^T).

The results obtained from chemotaxonomic analysis were consistent with the result of 16S rRNA gene sequence analysis. The predominant respiratory lipoquinone detected in strain DS-4^T was ubiquinone-10 (Q-10) at a peak area ratio of approximately 93%; minor amounts of Q-8 and Q-9 were detected. This predominant ubiquinone type was the same as those of *Sphingomonas* species (Yabuuchi *et al.*, 1990; Takeuchi *et al.*, 1995, 2001; Lee *et al.*, 2001; Busse *et al.*, 2003; Ohta *et al.*, 2004). The fatty acid profile of strain DS-4^T comprised unsaturated fatty acids C_{18:1}ω7c (45.7%) and C_{16:1}ω5c (1.3%), straight-chain fatty acids C_{16:0} (19.0%) and C_{14:0} (0.8%), summed feature 3 comprising C_{16:1}ω7c and/or iso-C_{15:0} 2-OH (14.8%), hydroxy fatty acid C_{14:0} 2-OH (12.1%) and 11-methyl C_{18:1}ω7c (6.3%). 3-Hydroxy fatty acids were not detected in strain DS-4^T. This fatty acid profile was similar to those of *Sphingomonas* species (Yabuuchi *et al.*, 1990; Takeuchi *et al.*, 1995, 2001; Lee *et al.*, 2001; Busse *et al.*, 2003; Ohta *et al.*, 2004). The major polar lipids were sphingoglycolipid, phosphatidylcholine, phosphatidylglycerol, diphosphatidylglycerol, phosphatidylethanolamine and unidentified phospholipid (Fig. 2). Phosphatidylmonomethylethanolamine and phosphatidyl-dimethylethanolamine were not detected in strain DS-4^T (Fig. 2). The polar lipid profile of strain DS-4^T was distinguishable from those of *Sphingomonas assaccharolytica*, *Sphingomonas mali* and *Sphingomonas pruni* in that phosphatidylcholine was present as a major polar lipid and phosphatidylmonomethylethanolamine and phosphatidyl-dimethylethanolamine were absent (Kämpfer *et al.*, 1997). The DNA G+C content of strain DS-4^T was 66.9 mol%.

Strain DS-4^T exhibited mean DNA–DNA relatedness levels of 9–17% with the type strains of six *Sphingomonas* species that showed 16S rRNA gene sequence similarity values of greater than 97% to strain DS-4^T: *S. aquatilis* KCTC 2881^T (10%), *S. asaccharolytica* KCTC 2825^T (15%), *S. koreensis* KCTC 2882^T (9%), *S. mali* KCTC 2826^T (17%), *S. pruni* KCTC 2824^T (10%) and *S. melonis* DSM 14444^T (17%). Strain DS-4^T differed from six phylogenetically related *Sphingomonas* species by several phenotypic characteristics (Table 1). The phylogenetic distinctiveness, together with DNA–DNA relatedness data and differential phenotypic properties, are sufficient to categorize strain DS-4^T as a member of a species that is distinct from the recognized

Table 1. Differential phenotypic characteristics of *Sphingomonas dokdonensis* sp. nov. DS-4^T and some phylogenetically related *Sphingomonas* species

Species: 1, *S. dokdonensis* sp. nov.; 2, *S. asaccharolytica* (data from Takeuchi *et al.*, 1995); 3, *S. mali* (data from Takeuchi *et al.*, 1995); 4, *S. pruni* (data from Takeuchi *et al.*, 1995); 5, *S. aquatilis* (data from Lee *et al.*, 2001); 6, *S. koreensis* (data from Lee *et al.*, 2001); 7, *S. melonis* (data from Buonauro *et al.*, 2002). +, Positive reaction; –, negative reaction; w, weakly positive reaction; ND, not determined. All species are rod-shaped and negative for urease, nitrate reduction, indole production, arginine dihydrolase and utilization of D-glucose (weakly positive for *S. asaccharolytica*) and citrate.

Characteristic	1	2	3	4	5	6	7
Colony colour	Yellow	Light-yellow	Light-yellow	Light-yellow	Yellow	Yellow	Deep-yellow
Motility	+	+	+	+	+	+	–
β -Galactosidase	–	+	+	+	+	+	+
Hydrolysis of:							
Aesculin	+	+	+	+	–	–	+
Gelatin	+	–	–	–	–	–	–
Utilization of:							
L-Arabinose	–	w	w	+	+	–	+
D-Cellobiose	+	–	+	+	ND	ND	+
D-Fructose	–	–	w	+	ND	ND	+
D-Galactose	–	–	+	+	ND	ND	+
Maltose	–	+	+	+	+	+	+
D-Mannose	+	w	+	+	–	–	+
Salicin	+	–	+	+	+	+	No growth or –
Sucrose	–	–	+	+	+	+	+
D-Trehalose	+	–	+	+	ND	ND	+
D-Xylose	–	+	+	+	ND	ND	+
Acetate	+	–	–	w	ND	ND	ND
N-Acetylglucosamine	–	+	+	+	+	+	+
Gluconate	–	–	+	+	–	–	–
Caprate	–	–	–	–	+	–	–
Adipate	+	–	+	–	–	–	–
L-Malate	–	–	–	w	–	+	+
DNA G+C content (mol%)	66.9	64.8	65.4–65.9	65.4	63	66	65.0

Sphingomonas species (Wayne *et al.*, 1987; Stackebrandt & Goebel, 1994). Therefore, on the basis of the data presented, strain DS-4^T should be placed in the genus *Sphingomonas* as a novel species, for which the name *Sphingomonas dokdonensis* sp. nov. is proposed.

Description of *Sphingomonas dokdonensis* sp. nov.

Sphingomonas dokdonensis (dok.do.nen'sis. N.L. fem. adj. *dokdonensis* of Dokdo, Korea, where the organism was first isolated).

Cells are Gram-negative rods (0.4–0.6 × 0.8–3.5 µm). Motile by means of a single polar flagellum. Colonies on TSA (pH 6.5) are circular, convex, smooth, glistening, yellow in colour and 0.8–1.0 mm in diameter after 4 days of incubation at 25 °C. Optimal temperature for growth is 25 °C. Growth occurs at 10 and 34 °C, but not at 4 and 35 °C. Optimal pH for growth is between 6.0 and 6.5; growth occurs at pH 5.0 and 9.5, but not at pH 4.5 and 10.0. Growth occurs in the presence of 0–5 % (w/v) NaCl; optimal growth occurs without NaCl. Anaerobic growth does not

occur on TSA (pH 6.5) or on TSA (pH 6.5) supplemented with nitrate. Catalase- and oxidase-positive. Casein, starch and Tween 20, Tween 40, Tween 60 and Tween 80 are hydrolysed, but hypoxanthine, xanthine and L-tyrosine are not. H₂S is not produced. Lysine decarboxylase, ornithine decarboxylase and tryptophan deaminase are absent. Pyruvate is utilized as a sole carbon and energy source, but D-mannitol, benzoate, formate, phenylacetate, succinate and L-glutamate are not. In assays with the API ZYM system, alkaline phosphatase, esterase (C4), esterase lipase (C8), leucine arylamidase, valine arylamidase, acid phosphatase and naphthol-AS-BI-phosphohydrolase are present and cystine arylamidase is weakly present, but lipase (C14), trypsin, α -chymotrypsin, α -galactosidase, β -glucuronidase, α -glucosidase, β -glucosidase, N-acetyl- β -glucosaminidase, α -mannosidase and α -fucosidase are absent. Susceptible to chloramphenicol, gentamicin, tetracycline, kanamycin, neomycin and oleandomycin, but not to polymyxin B, streptomycin, penicillin G, ampicillin, cephalothin, novobiocin, carbenicillin or lincomycin. The predominant ubiquinone is Q-10. The major fatty acids (>10 % of total fatty acids) are C_{18:1}ω7c, C_{16:0}, C_{14:0} 2-OH and

Fig. 1. Neighbour-joining phylogenetic tree based on 16S rRNA gene sequences showing the positions of *Sphingomonas dokdonensis* sp. nov. DS-4^T, *Sphingomonas* species and some other related taxa. Bootstrap values (expressed as percentages of 1000 replicates) of >50% are shown at branch points. *Rhodospirillum rubrum* ATCC 11170^T was used as an outgroup. Bar, 0.01 substitutions per nucleotide position.

C_{16:1}ω7c and/or iso-C_{15:0} 2-OH. Major polar lipids are sphingoglycolipid, phosphatidylcholine, phosphatidylglycerol, diphosphatidylglycerol, phosphatidylethanolamine and unidentified phospholipid. The DNA G+C content

is 66.9 mol% (determined by HPLC). Other phenotypic characteristics are given in Table 1.

The type strain, DS-4^T (=KCTC 12541^T=CIP 108841^T), was isolated from soil.

Acknowledgements

This work was supported by the 21C Frontier program of Microbial Genomics and Applications (grant MG05-0401-2-0) from the Ministry of Science and Technology (MOST) of the Republic of Korea. We are grateful to the Ulleung County Administration and the Cultural Heritage Administration of the Republic of Korea for aiding access to Dokdo.

References

- Buonaurio, R., Stravato, V. M., Kosako, Y., Fujiwara, N., Naka, T., Kobayashi, K., Cappelli, C. & Yabuuchi, E. (2002). *Sphingomonas melonis* sp. nov., a novel pathogen that causes brown spots on yellow Spanish melon fruits. *Int J Syst Evol Microbiol* 52, 2081–2087.
- Busse, H.-J., Denner, E. B. M., Buczolits, S., Salkinoja-Salonen, M., Bennisar, A. & Kämpfer, P. (2003). *Sphingomonas aurantiaca* sp. nov., *Sphingomonas aerolata* sp. nov. and *Sphingomonas faeni* sp. nov., air- and dustborne and Antarctic, orange-pigmented, psychrotolerant bacteria, and emended description of the genus *Sphingomonas*. *Int J Syst Evol Microbiol* 53, 1253–1260.
- Cowan, S. T. & Steel, K. J. (1965). *Manual for the Identification of Medical Bacteria*. London: Cambridge University Press.

Fig. 2. Two-dimensional thin-layer chromatogram of polar lipids of *Sphingomonas dokdonensis* sp. nov. DS-4^T. SGL, sphingoglycolipid; PC, phosphatidylcholine; PG, phosphatidylglycerol; DPG, diphosphatidylglycerol; PE, phosphatidylethanolamine; PL, unidentified phospholipid.

- Ezaki, T., Hashimoto, Y. & Yabuuchi, E. (1989). Fluorometric deoxyribonucleic acid–deoxyribonucleic acid hybridization in micro-dilution wells as an alternative to membrane filter hybridization in which radioisotopes are used to determine genetic relatedness among bacterial strains. *Int J Syst Bacteriol* **39**, 224–229.
- Kämpfer, P., Denner, E. B. M., Meyer, S., Moore, E. R. B. & Busse, H.-J. (1997). Classification of “*Pseudomonas azotocolligans*” Anderson 1955, 132, in the genus *Sphingomonas* as *Sphingomonas trueperi* sp. nov. *Int J Syst Bacteriol* **47**, 577–583.
- Komagata, K. & Suzuki, K. (1987). Lipids and cell-wall analysis in bacterial systematics. *Methods Microbiol* **19**, 161–203.
- Lányi, B. (1987). Classical and rapid identification methods for medically important bacteria. *Methods Microbiol* **19**, 1–67.
- Lee, J.-S., Shin, Y. K., Yoon, J.-H., Takeuchi, M., Pyun, Y.-R. & Park, Y.-H. (2001). *Sphingomonas aquatilis* sp. nov., *Sphingomonas koreensis* sp. nov. and *Sphingomonas taejonensis* sp. nov., yellow-pigmented bacteria isolated from natural mineral water. *Int J Syst Evol Microbiol* **51**, 1491–1498.
- Minnikin, D. E., O'Donnell, A. G., Goodfellow, M., Anderson, G., Athalye, M., Schaal, A. & Parlett, J. H. (1984). An integrated procedure for the extraction of bacterial isoprenoid quinones and polar lipids. *J Microbiol Methods* **2**, 233–241.
- Ohta, H., Hattori, R., Ushiba, Y., Mitsui, H., Ito, M., Watanabe, H., Tonosaki, A. & Hattori, T. (2004). *Sphingomonas oligophenolica* sp. nov., a halo- and organo-sensitive oligotrophic bacterium from paddy soil that degrades phenolic acids at low concentrations. *Int J Syst Evol Microbiol* **54**, 2185–2190.
- Stackebrandt, E. & Goebel, B. M. (1994). Taxonomic note: a place for DNA–DNA reassociation and 16S rRNA sequence analysis in the present species definition in bacteriology. *Int J Syst Bacteriol* **44**, 846–849.
- Takeuchi, M., Sakane, T., Yanagi, M., Yamasato, K., Hamana, K. & Yokota, A. (1995). Taxonomic study of bacteria isolated from plants: proposal of *Sphingomonas rosa* sp. nov., *Sphingomonas pruni* sp. nov., *Sphingomonas asaccharolytica* sp. nov., and *Sphingomonas mali* sp. nov. *Int J Syst Bacteriol* **45**, 334–341.
- Takeuchi, M., Hamana, K. & Hiraishi, A. (2001). Proposal of the genus *Sphingomonas sensu stricto* and three new genera, *Sphingobium*, *Novosphingobium* and *Sphingopyxis*, on the basis of phylogenetic and chemotaxonomic analyses. *Int J Syst Evol Microbiol* **51**, 1405–1417.
- Tamaoka, J. & Komagata, K. (1984). Determination of DNA base composition by reverse-phase high-performance liquid chromatography. *FEMS Microbiol Lett* **25**, 125–128.
- Wayne, L. G., Brenner, D. J., Colwell, R. R. & 9 other authors (1987). International Committee on Systematic Bacteriology. Report of the ad hoc committee on reconciliation of approaches to bacterial systematics. *Int J Syst Bacteriol* **37**, 463–464.
- Yabuuchi, E., Yano, I., Oyaizu, H., Hashimoto, Y., Ezaki, T. & Yamamoto, H. (1990). Proposals of *Sphingomonas paucimobilis* gen. nov. and comb. nov., *Sphingomonas parapaucimobilis* sp. nov., *Sphingomonas yanoikuyae* sp. nov., *Sphingomonas adhaesiva* sp. nov., *Sphingomonas capsulate* comb. nov., and two genospecies of the genus *Sphingomonas*. *Microbiol Immunol* **34**, 99–119.
- Yoon, J.-H., Kim, H., Kim, S.-B., Kim, H.-J., Kim, W. Y., Lee, S. T., Goodfellow, M. & Park, Y.-H. (1996). Identification of *Saccharomonospora* strains by the use of genomic DNA fragments and rRNA gene probes. *Int J Syst Bacteriol* **46**, 502–505.
- Yoon, J.-H., Lee, S. T. & Park, Y.-H. (1998). Inter- and intraspecific phylogenetic analysis of the genus *Nocardioides* and related taxa based on 16S rRNA gene sequences. *Int J Syst Bacteriol* **48**, 187–194.
- Yoon, J.-H., Kang, K. H. & Park, Y.-H. (2003). *Psychrobacter jeotgali* sp. nov., isolated from jeotgal, a traditional Korean fermented seafood. *Int J Syst Evol Microbiol* **53**, 449–454.
- Yoon, J.-H., Kang, S.-J., Lee, S.-Y., Lee, M.-H. & Oh, T.-K. (2005). *Virgibacillus dokdonensis* sp. nov., isolated from a Korean island, Dokdo, located at the edge of the East Sea in Korea. *Int J Syst Evol Microbiol* **55**, 1833–1837.
- Yoon, J.-H., Kang, S.-J. & Oh, T.-K. (2006). *Dokdonella koreensis* gen. nov., sp. nov., isolated from soil. *Int J Syst Evol Microbiol* **56**, 145–150.
- Yurkov, V., Stackebrandt, E., Holmes, A. & 7 other authors (1994). Phylogenetic positions of novel aerobic, bacteriochlorophyll a-containing bacteria and description of *Roseococcus thiosulfatophilus* gen. nov., sp. nov., *Erythromicrobium ramosum* gen. nov., sp. nov., and *Erythrobacter litoralis* sp. nov. *Int J Syst Bacteriol* **44**, 427–434.