Eudoraea adriatica gen. nov., sp. nov., a novel marine bacterium of the family *Flavobacteriaceae*

Karine Alain,† Laurent Intertaglia, Philippe Catala and Philippe Lebaron

UPMC Univ Paris 06, UMR7621, Observatoire Océanologique, F-66650 Banyuls-sur-Mer, France

Correspondence

Philippe Lebaron

philippe.lebaron@obs-banyuls.fr

A novel aerobic, non-motile, Gram-negative bacterium, designated AS06/20a^T, was isolated from coastal waters of the Adriatic Sea and subjected to polyphasic taxonomic analysis. Cells were rod-shaped and formed non-pigmented punctiform colonies on agar plates. The novel isolate grew heterotrophically on a variety of carbon compounds, including organic acids, carbohydrates, polyols, amino acids and complex organic substrates. Cells grew at 15–33 °C (optimum 30 °C), pH 6.5–8.5 (optimum pH 7.5–8.0) and between 2 and 6% (w/v) NaCl (optimum 2%). The dominant fatty acids (>5%) detected in strain AS06/20a^T were iso-C_{15:1} G, iso-C_{15:0}, iso-C_{15:0} 3-OH, iso-C_{17:1} ω9c and iso-C_{17:0} 3-OH. The major respiratory quinone was MK-6. The G+C content of the genomic DNA was 38.9 mol%. Phylogenetic analyses based on 16S rRNA gene sequences showed that strain AS06/20a^T fell within the family *Flavobacteriaceae* in the phylum *Bacteroidetes*. Comparative 16S rRNA gene sequence analysis indicated that strain AS06/20a^T was most closely related to members of the genera *Zeaxanthinibacter* and *Robiginitalea*, sharing 92–93% 16S rRNA gene sequence similarity with representatives of these genera. It is proposed, from the results of the polyphasic taxonomic analysis, that the strain should be placed in a new genus, *Eudoraea* gen. nov., with *Eudoraea adriatica* sp. nov. as the type species. The type strain is AS06/20a^T (=DSM 19308^T=CIP 109577^T=OBB 358^T).

Bacteria belonging to the phylum *Bacteroidetes* are widely distributed in nature and have been found in almost every habitat of the biosphere, the lithosphere and the hydrosphere (Bernardet *et al.*, 2002; Kirchman, 2002). They seem particularly common and abundant in marine surface waters, as indicated by fluorescence *in situ* hybridization analyses (Glöckner *et al.*, 1999; Cottrell & Kirchman, 2000a; Simon *et al.*, 1999; Alonso & Pernthaler, 2005). Cultured isolates of *Bacteroidetes* are all heterotrophs and especially efficient in degrading complex biomacromolecules such as protein, chitin, pectin, agar, starch and cellulose.

In marine ecosystems, the heterotrophic prokaryotes are the main consumers of dissolved organic material (DOM) and represent a very dynamic compartment in global

biogeochemical cycles. Uptake of DOM is the first step in the microbial loop that ultimately mineralizes over half of the organic matter formed by photosynthetic bacteria and algae (Cole *et al.*, 1988). Although the importance of DOM uptake is well recognized, the relative contributions of the different phylogenetic groups to DOM uptake remain to be elucidated. Because of their abundance and heterotrophic metabolism, *Bacteroidetes* are thought to occupy a special place in the carbon cycling of oceans. They are obviously involved in the degradation and uptake of DOM and probably contribute to the degradation of the high-molecular-mass (HMM; >1 kDa) fraction of the DOM (Kirchman, 2002). Indeed, the HMM fraction is composed of biopolymers; hydrolysis of these macromolecules requires specific extracellular enzymes, which are produced by numerous known members of the *Bacteroidetes*. Using a method combining microautoradiography and hybridization of fluorescent rRNA-targeted oligonucleotide probes to whole cells from natural marine assemblages (MicroFISH), Cottrell & Kirchman (2000b) demonstrated that *Bacteroidetes* represented, for the considered habitats, a large fraction of the community that was efficient in degrading chitin, *N*-acetylglucosamine and protein, part of the HMM fraction of the DOM. All these data together suggest strongly that, in marine environments, *Bacteroidetes* are key players in the degradation of the HMM fraction of the DOM, which has been demonstrated

†Present address: UMR6197, Laboratoire de Microbiologie des Environnements Extrêmes, IUEM, Technopôle Brest-Iroise, F-29280 Plouzané, France.

Abbreviations: ASW, artificial seawater; DOM, dissolved organic material; HMM, high molecular mass.

The GenBank/EMBL/DDBJ accession number for the 16S rRNA gene sequence of *Eudoraea adriatica* AS06/20a^T is AM745437.

A scanning electron micrograph of cells of strain AS06/20a^T and graphs showing the effects of temperature, pH and salinity on the maximum growth rate of the novel isolate are available as supplementary material with the online version of this paper.

to be more bioreactive and more bioavailable than the low-molecular-mass fraction (Amon & Benner, 1996). As relevant work on cultured representatives from marine origin is lacking, there is great interest in isolating and describing phenotypic and genotypic characteristics of marine *Bacteroidetes* known only by their 16S rRNA gene sequences. For example, there are notably very few sequence data on enzymes catalysing biopolymer hydrolysis (Kirchman, 2002). The isolation of strains is still the easiest way to access their physiological and genomic capital. This is probably the reason why the number of strains of *Bacteroidetes* from marine ecosystems described in the last five years has increased exponentially.

In this study, a novel marine bacterium belonging to the phylum *Bacteroidetes* is described. Based on the results of a polyphasic taxonomic analysis, strain AS06/20a^T represents a novel species in a new genus, *Eudoraea adriatica* gen. nov., sp. nov.

In April 2006, marine surface waters were collected on the coast of the Adriatic Sea, Italy (44.690° N 12.520° E). One subsample collected at a depth of 10 m was spread on a marine agar 2216 (MA; Difco) plate and then incubated at 25 °C. After 2 weeks, a small non-pigmented colony was picked and purified by repeated streaking on MA plates. It was referenced as strain AS06/20a^T. Stock cultures were stored at -80 °C in marine broth 2216 (MB; Difco) supplemented with 5 % (v/v) DMSO or 35 % (v/v) glycerol until characterization.

The almost complete 16S rRNA gene (1393 bp) of the strain was a double-strand sequenced from one single colony, as described elsewhere (Agogue *et al.*, 2005). This sequence was compared to those in available databases by using the program BLAST (Altschul *et al.*, 1990) and then aligned to its nearest neighbours using the program CLUSTAL_X (Thompson *et al.*, 1997). Alignments were refined manually using the program SEAVIEW (Galtier *et al.*, 1996). Phylogenetic trees were constructed using the PHYLIP version 3.63 software (<http://evolution.genetics.washington.edu/phylip/getme.html>) on the basis of evolutionary distance (neighbour-joining method with Jukes and Cantor corrections) (Saitou & Nei, 1987) and maximum-likelihood (Felsenstein, 1981). The robustness of the inferred topologies was assessed by bootstrap analyses based on 1000 bootstrap resamplings for the neighbour-joining and 100 replications for the maximum-likelihood method (Felsenstein, 1985). The 16S rRNA gene-based analysis placed strain AS06/20a^T within the phylum *Bacteroidetes*. The novel isolate was affiliated to the family *Flavobacteriaceae*, one of the main phyletic lines of the phylum *Bacteroidetes* (Reichenbach, 1989; Bernardet *et al.*, 1996, 2002) (Fig. 1). The results of different phylogenetic reconstructions performed with different treeing algorithms were in accordance with each other. The novel isolate formed a distinct lineage together with *Zeaxanthinibacter enoshimensis* TD-ZE3^T and *Robiginitalea biformata* HTCC2501^T, but this lineage did not cluster

robustly with any of the recognized genera in the family. Within this lineage, the novel isolate showed only a distant relatedness to its nearest phylogenetic neighbours, the most closely related genus being *Zeaxanthinibacter* (93 % 16S rRNA gene sequence similarity), followed by *Robiginitalea* (92 %). These low 16S rRNA gene sequence similarities between strain AS06/20a^T and the genera described so far suggest that the novel isolate represents a member of a new genus within the family *Flavobacteriaceae*.

The DNA G+C content was determined by the Identification Service of the DSMZ (Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH, Braunschweig, Germany) by HPLC analysis of deoxyribonucleosides as described by Mesbah *et al.* (1989). The G+C content of strain AS06/20a^T was 38.9 mol%.

Colonies on MA were non-pigmented and cream in colour. They were circular, punctiform, opaque with an entire edge and possessed a smooth surface. After 1 week incubation, colonies were less than 1 mm in diameter. Morphological characteristics of the cells were determined by light microscopy (Olympus AX70) and by transmission electron microscopy (Hitachi H-7500) after negative staining with uranyl acetate (Raguénès *et al.*, 1997). Gram-staining was determined using standard procedures. Gliding motility was determined using the hanging drop method on cells grown at 30 °C in low nutrient medium [0.1 % (w/v) MB solidified with 1 % agar], as described by Bernardet *et al.* (2002). In summary, cells of strain AS06/20a^T were Gram-negative, non-motile, straight rods of 1.48–2.99 µm in length (mean 2.22 ± 0.54 µm; *n*=8) and 0.6–0.78 µm in width (mean 0.70 ± 0.07 µm; *n*=8) in the mid-exponential phase of growth (see Supplementary Fig. S1 in IJSEM Online). Gliding motility was not observed under the conditions tested. Cells swelled into balloon-like shapes during the late stationary growth phase.

In order to analyse respiratory quinones and polar lipids, strain AS06/20a^T was grown for 5 days on MB medium at 30 °C and checked for purity. Analyses of isoprenoid quinones and polar lipids were carried out by the Identification Service of the DSMZ (B. Tindall, DSMZ Culture Collection). As described for the other members of the family *Flavobacteriaceae*, the main respiratory quinone was menaquinone-6 (MK-6; 83 %) (Bernardet *et al.*, 2002); MK-7 was also detected (17 %). Phosphatidylethanolamine was the only major phospholipid identified in strain AS06/20a^T. Three unidentified polar lipids, two containing amino groups, were also present as major components.

Determination of the whole-cell fatty acid composition was performed on cultures grown at 30 °C for 72 h on MA. This analysis was carried out at the DSMZ according to the standard protocol of the Microbial Identification System (MIDI, 2001). Extracts were analysed using a Hewlett Packard model HP6890A GC equipped with a flame-ionization detector as described by Kämpfer & Kroppenstedt (1996). Results are summarized in Table 1. The dominant fatty acids in strain AS06/20a^T were

Fig. 1. Phylogenetic tree based on 16S rRNA gene sequences of strain AS06/20a^T and representative members of related genera within the family Flavobacteriaceae. Sequence data of reference strains were obtained from the GenBank/EMBL and/or RDP databases. Accession numbers are indicated in parentheses. The topology shown corresponds to an unrooted tree obtained by the maximum-likelihood algorithm, established using the PHYLIP package. Bootstrap values (from 100 replicates) are indicated at branch nodes. The positioning of the novel isolate was confirmed using the neighbour-joining method. Bar, 2.0 nt substitutions per 100 nt.

iso-C_{15:1} G, iso-C_{15:0}, iso-C_{15:0} 3-OH, iso-C_{17:1} ω9c, iso-C_{17:0} 3-OH and the fatty acids in summed feature 3.

Unless stated otherwise, physiological characterization was carried out aerobically in MB in triplicate and incubation was done in the dark and under agitation. Growth was routinely monitored by measuring the increase in optical density at 600 nm using a spectrophotometer. Cell numbers were determined by flow cytometry in order to calculate calibration curves [cell number = f(OD₆₀₀)]. Growth rates were calculated using linear regression analysis of four to nine points along the logarithmic portions of the resulting growth curves. Growth temperatures were tested over the range 9–44 °C (i.e. 9, 15, 20, 25, 30, 33, 37 and 44 °C). The temperature range for growth was 15–33 °C, with optimum growth rate at 30 °C (see Supplementary Fig. S2a in IJSEM Online). No growth was observed at 9 or 37 °C. The pH range for growth was tested at 30 °C in MB medium, buffered and adjusted to the required pH as described elsewhere (Alain *et al.*, 2002). The pH range for growth was rather narrow (see Supplementary Fig. S2b in IJSEM Online); growth was observed at pH 6.5–8.5, the optimum being around pH 7.5–8.0. Salt tolerance was tested at 30 °C in MB medium prepared with various concentrations of NaCl (0.02, 0.5, 1, 2, 3, 4, 5, 6, 7 and 9%, w/v). The isolate

required NaCl for growth (see Supplementary Fig. S2c in IJSEM Online). This is congruent with the fact that this strain was isolated from brackish waters of the North Adriatic Sea. Growth was observed at salt concentrations ranging from 2 to 6% (w/v) NaCl, the optimum salinity being around 2%. No growth was observed at 1 or 7% (w/v) NaCl.

Strain AS06/20a^T was aerobic. Conventional phenotypic tests, including those for oxidase, catalase, Tween esterase and nitrate reductase, were performed according to standard methods (Smibert & Krieg, 1994). The results are given in Table 1. Biochemical tests were performed at 30 °C using API ZYM (bioMérieux) and Biolog GN2 microplates (Oxoid). These tests were inoculated with cells grown on MA plates, swabbed from the surface of the agar plates and then suspended in half-strength artificial seawater (ASW) to the density specified by the manufacturer. Supplementary biochemical tests were also performed using API 20NE strips (bioMérieux), following the manufacturer's instructions. The data obtained are given in Table 1. Testing for oxidation of carbon sources with Biolog GN2 plates indicated that the strain was able to oxidize a range of sugars, organic acids and amino acids. To confirm these results and to test for the capability of the strain to catabolize different substrates as sole carbon and

Table 1. Phenotypic and genotypic characteristics of strain AS06/20a^T (*Eudoraea adriatica* gen. nov., sp. nov.)

Characteristic	Strain AS06/20a ^T
Temperature range for growth (°C) (optimum)	15–33 (30)
NaCl range for growth (% w/v) (optimum)	2–6 (2)
pH range for growth (optimum)	6.5–8.5 (7.5–8.0)
Biochemical properties (API 20NE)	
Catalase, oxidase, aesculin hydrolysis, β -galactosidase	+
Denitrification, indole production, glucose fermentation, arginine dihydrolase, urease, gelatin hydrolysis	–
Enzymic activities (API ZYM)	
Alkaline phosphatase, esterase (C4), esterase lipase (C8), valine arylamidase, leucine arylamidase, α -chymotrypsin, acid phosphatase, naphthol-AS-BI-phosphohydrolase, β -galactosidase, β -glucosidase, <i>N</i> -acetyl- β -glucosaminidase, α -mannosidase	+
Lipase (C14), cystine arylamidase, trypsin, α -galactosidase, α -glucosidase, α -fucosidase	–
Substrate assimilation	
D-Glucose, L-arabinose, D-mannose, D-mannitol, <i>N</i> -acetylglucosamine, potassium gluconate, <i>m</i> -inositol, acetic acid, citric acid, pyruvic acid, propionic acid, capric acid, adipic acid, malic acid, phenylacetic acid, L-arginine, yeast extract	+
Formic acid, Tween 40, Tween 80, maltose	–
Degradation of macromolecules	
Aesculin	+
Agar, cellulose (filter paper), gelatin, starch, urea	–
Susceptibility to antibiotics	
Vancomycin, rifampicin, tetracycline, polymyxin B	+
Oxacillin, penicillin, ciprofloxacin	–
Fatty acid composition (%) [*]	iso-C _{15:1} G (15.97 %), iso-C _{15:0} (29.37 %), iso-C _{15:0} 3-OH (7.27 %), iso-C _{17:1} ω 9c (8.96 %), iso-C _{17:0} 3-OH (14.13 %) and summed feature 3 (11.57 %)
Polar lipids [†]	PE, L, AL1, AL2
Quinones	MK-6 (83 %), MK-7 (17 %)
DNA G + C content (mol%)	38.9

^{*}Fatty acids comprising more than 5 % of the total content are listed. The following fatty acids were present in lesser amounts: iso-C_{13:0} (0.48 %), C_{14:0} (0.17 %), anteiso-C_{15:0} (0.72 %), C_{15:1} ω 6c (1.06 %), iso-C_{16:0} (0.25 %), C_{16:0} (0.67 %), C_{15:0} 2-OH (0.30 %), C_{15:0} 3-OH (1.33 %), iso-C_{17:0} (0.17 %), C_{17:1} ω 8c (0.48 %), C_{17:1} ω 6c (0.83 %), iso-C_{16:0} 3-OH (1.38 %), C_{16:0} 3-OH (0.37 %), C_{18:1} ω 5c (0.70 %), C_{17:0} 3-OH (0.20 %), and three unknown fatty acids of equivalent chain-length 11.54 (0.21 %), 13.56 (2.26 %) and 16.58 (1.15 %). Summed feature 3 comprises iso-C_{15:0} 2-OH and/or C_{16:1} ω 7c.

[†]PE, Phosphatidylethanolamine; L, unidentified polar lipid; AL, unidentified aminolipid.

energy sources, with oxygen as a terminal electron acceptor, the strain was grown aerobically in the dark on a mineral medium supplemented with one substrate. The defined medium had the following composition (l⁻¹): phosphate buffer, 30 mM; NaCl, 20 g; MgCl₂ · 6H₂O, 3 g; CaCl₂ · 2H₂O, 1.0 g; NH₄Cl, 0.3 g; KCl, 0.5 g; Na₂SO₄, 3 g; NaNO₃, 1 g; trace element solution, 1 ml; selenite-tungstate solution, 1 ml; vitamin solution, 1 ml. The strain grew chemo-organoheterotrophically on a variety of carbon compounds, including pentoses, hexoses, polyols, organic acids, amino acids and complex organic substrates (Table 1).

Antibiotic sensitivity tests were performed by using susceptibility discs (Bio-Rad) or filter-paper discs impreg-

nated with different antibiotics. Discs were placed on MA plates spread with a culture of the isolate and were then incubated at 30 °C for 1 week. Susceptibility was scored as positive at zone diameters above 10 mm. The results are summarized in Table 1.

The phenotypic and genotypic characteristics of the novel isolate generally met the minimal standards for the family *Flavobacteriaceae* (Bernardet *et al.*, 2002). Nevertheless, strain AS06/20a^T differed from members of other genera of the *Flavobacteriaceae* with validly published names in a number of phylogenetic, genotypic, chemotaxonomic, morphological and physiological features (Table 2). In brief, in addition to the phylogenetic distance, the novel taxon can be distinguished from its closest relatives by

Table 2. Differential characteristics of strain AS06/20a^T (*Eudoraea adriatica* gen. nov., sp. nov.) and members of related genera of the family *Flavobacteriaceae*

Strains: 1, AS06/20a^T (data from this study); 2, *Zeaxanthinibacter enoshimensis* TD-ZE3^T (Asker *et al.*, 2007); 3, *Robiginitalea biformata* HTCC2501^T (Cho & Giovannoni, 2004); 4, *Pibocella ponti* KMM 6031^T (Nedashkovskaya *et al.*, 2005); 5, *Maribacter sedimenticola* KMM 3903^T (Nedashkovskaya *et al.*, 2004); 6, *Zobellia galactanivorans* Dsij^T (Barbeyron *et al.*, 2001); 7, *Arenibacter latericius* KMM 426^T (Ivanova *et al.*, 2001); 8, *Sediminicola luteus* CNI-3^T (Khan *et al.*, 2006). +, Positive; –, negative; ND, not determined; w, weak.

Characteristic	1	2	3	4	5	6	7	8
Oxidase/catalase	+/+	+/+	+/+	+/+	+/+	+/+	+/+	+/w
Pigmentation*	–	Y	O	Y	Y	Y	DO	Y
Flexirubin pigments	–	–	–	–	–	+	–	–
Gliding motility	–	+	–	+	+	+	–	–
Growth in/at:								
8 % NaCl	–	–	+	+	+	–	–	–
7 % NaCl	–	+	+	+	–	–	+	+
1 % NaCl	–	ND	+	+	+	+	+	+
4 °C	–	–	–	+	+	–	–	–
10 °C	–	–	+	+	+	–	+	+
42 °C	–	–	+	–	–	+	–	–
Hydrolysis of:								
Agar	–	–	ND	–	+	+	–	ND
Gelatin	–	+	–	+	+	+	–	+
Starch	–	+	+	+	–	+	–	+
Cellulose	–	–	–	–	–	ND	–	–
Aesculin	+	+	+	ND	ND	+	–	ND
Urease	–	–	–	–	–	–	+	–
β-Galactosidase	+	+	–	ND	–	+	+	ND
Nitrate reduction	–	–	–	–	–	+	+	+
DNA G + C content (mol%)	38.9	46.5	56.4	35.5	37	43	37–38	38
16S rRNA gene sequence similarity (%)†	100	93	92	90	91	91	91	90

*Y, Yellow; O, orange; DO, dark orange.

†Similarities calculated in reference to the 16S rRNA gene sequence of strain AS06/20a^T.

some of its phenotypic features, including its absence of colonial pigmentation, the narrow NaCl and pH ranges for growth, and its restricted degrading capabilities regarding the tested macromolecules. In addition to these phenotypic differences, the novel isolate is unambiguously distinct from its two closest neighbours by its genomic DNA G + C content, which is more than 5 mol% lower than those of *Zeaxanthinibacter enoshimensis* and *Robiginitalea biformata*. In conclusion, in view of all the above-mentioned distinctive features, it is proposed that isolate AS06/20a^T represents a novel species in a new genus, *Eudoraea adriatica* gen. nov., sp. nov.

Description of *Eudoraea* gen. nov.

Eudoraea (Eu.do.ra'e.a. N.L. fem. n. *Eudora* a sea goddess in Greek mythology; N.L. fem. n. *Eudoraea* named after *Eudora*).

Cells are Gram-negative, non-spore-forming, non-motile, non-gliding rods. Pigments are not produced. Aerobic and chemo-organoheterotrophic. Catalase- and oxidase-positive. Mesophilic. Neutrophilic. Require NaCl for growth. The

predominant quinone is MK-6. Polar lipids include phosphatidylethanolamine. Major fatty acids are iso-C_{15:1}G, iso-C_{15:0}, iso-C_{15:0} 3-OH, iso-C_{17:1}ω9c, iso-C_{17:0} 3-OH and summed feature 3. The G + C content of the DNA is close to 39 mol%. Phylogenetically, the genus *Eudoraea* belongs to the family *Flavobacteriaceae*, phylum *Bacteroidetes*, showing a distant relatedness to the marine genera *Zeaxanthinibacter* and *Robiginitalea*. The type species is *Eudoraea adriatica*.

Description of *Eudoraea adriatica* sp. nov.

Eudoraea adriatica (a.dri.a'ti.ca. L. fem. adj. *adriatica* of the Adriatic Sea, where the type strain was isolated).

In addition to the characters described for the genus, the following properties apply. Colonies on MA medium are punctiform, cream in colour, opaque and smooth. Optimal growth occurs at 30 °C; grows at 15–33 °C. pH and NaCl ranges for growth are 6.5–8.5 (optimum 7.5–8.0) and 2–6 % (w/v) (optimum 2 %, w/v), respectively. Aesculin is hydrolysed, but agar, cellulose, gelatin, starch and urea are not. β-Galactosidase-positive. Glucose is not fermented.

Nitrate is not reduced. Various carbon compounds are used as sole carbon sources, including pentoses, hexoses, polyols, organic acids, amino acids and complex substrates (Table 1).

The type strain, AS06/20a^T (=DSM 19308^T=CIP 109577^T=OOB 358^T), was isolated from coastal waters of the Adriatic Sea, Italy (44.690° N 12.520° E). The DNA G+C content of strain AS06/20a^T is 38.9 mol%.

Acknowledgements

We thank Marie-Line Escande for assistance with the transmission electron microscopy. We acknowledge Professor J. P. Euzéby for support in the Latin etymologies of the genus and species names. This work was supported financially by the Equipe Mixte de Recherche linking the University Pierre et Marie Curie and the Centre National de la Recherche Scientifique to the Pierre Fabre Laboratories. The project was also carried out in the frame of the MarBEF Network of Excellence 'Marine Biodiversity and Ecosystem Functioning', which is funded by the Sustainable Development, Global Change and Ecosystems Program of the European Community's Sixth Framework Program (contract no. GOCE-CT-2003-505446). This publication is contribution number MPS-07060 of MarBEF. It was also partly funded by the French program 'Bio-diversité et Changement Global – project: development of a coastal microbial observatory' from the 'Institut Français de la Biodiversité'.

References

- Agogue, H., Casamayor, E. O., Bourrain, M., Obernosterer, I., Joux, F., Herndl, G. J. & Lebaron, P. (2005). A survey on bacteria inhabiting the sea surface microlayer of coastal ecosystems. *FEMS Microbiol Ecol* **54**, 269–280.
- Alain, K., Querellou, J., Lesongeur, F., Pignet, P., Crassous, P., Raguénès, G., Cueff, V. & Cambon-Bonavita, M.-A. (2002). *Caminibacter hydrogenophilus* gen. nov., sp. nov., a novel thermophilic, hydrogen-oxidizing bacterium isolated from an East Pacific Rise hydrothermal vent. *Int J Syst Evol Microbiol* **52**, 1317–1323.
- Alonso, C. & Pernthaler, J. (2005). Incorporation of glucose under anoxic conditions by bacterioplankton from coastal North Sea surface waters. *Appl Environ Microbiol* **71**, 1709–1716.
- Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. (1990). Basic local alignment search tool. *J Mol Biol* **215**, 403–410.
- Amon, R. M. W. & Benner, R. (1996). Bacterial utilization of different size classes of dissolved organic matter. *Limnol Oceanogr* **41**, 41–51.
- Asker, D., Beppu, T. & Ueda, K. (2007). *Zeaxanthinibacter enoshimensis* gen. nov., sp. nov., a novel zeaxanthin-producing marine bacterium of the family *Flavobacteriaceae*, isolated from seawater off Enosima Island, Japan. *Int J Syst Evol Microbiol* **57**, 837–843.
- Barbeyron, T., L'Haridon, S., Corre, E., Kloareg, B. & Potin, P. (2001). *Zobellia galactanovorans* gen. nov., sp. nov., a marine species of *Flavobacteriaceae* isolated from a red alga, and classification of [*Cytophaga*] *uliginosa* (Zobell and Upham 1944) Reichenbach 1989 as *Zobellia uliginosa* gen. nov., comb. nov. *Int J Syst Evol Microbiol* **51**, 985–997.
- Bernardet, J.-F., Segers, P., Vancanneyt, M., Berthe, F., Kersters, K. & Vandamme, P. (1996). Cutting a Gordian knot: emended classification and description of the genus *Flavobacterium*, emended description of the family *Flavobacteriaceae*, and proposal of *Flavobacterium hydatis* nom. nov. (basonym, *Cytophaga aquatilis* Strohl and Tait 1978). *Int J Syst Bacteriol* **46**, 128–148.
- Bernardet, J.-F., Nakagawa, Y. & Holmes, B., for the Subcommittee on the taxonomy of *Flavobacterium* and *Cytophaga*-like bacteria of the International Committee on Systematics of Prokaryotes (2002). Proposed minimal standards for describing new taxa of the family *Flavobacteriaceae* and emended description of the family. *Int J Syst Evol Microbiol* **52**, 1049–1070.
- Cho, J.-C. & Giovannoni, S. J. (2004). *Robiginitalea biformata* gen. nov., sp. nov., a novel marine bacterium in the family *Flavobacteriaceae* with a higher G+C content. *Int J Syst Evol Microbiol* **54**, 1101–1106.
- Cole, J. J., Findlay, S. & Pace, M. L. (1988). Bacterioplanktonic production in fresh and salt-water ecosystems: a crossover view. *Mar Ecol Prog Ser* **43**, 1–10.
- Cottrell, M. T. & Kirchman, D. L. (2000a). Community composition of marine bacterioplankton determined by 16S rRNA gene clone libraries and fluorescence in situ hybridization. *Appl Environ Microbiol* **66**, 5116–5122.
- Cottrell, M. T. & Kirchman, D. L. (2000b). Natural assemblages of marine proteobacteria and members of the *Cytophaga-Flavobacter* cluster consuming low- and high-molecular-weight dissolved organic matter. *Appl Environ Microbiol* **66**, 1692–1697.
- Felsenstein, J. (1981). Evolutionary trees from DNA sequences: a maximum likelihood approach. *J Mol Evol* **17**, 368–376.
- Felsenstein, J. (1985). Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* **39**, 783–791.
- Galtier, N., Gouy, M. & Gautier, C. (1996). SEAVIEW and PHYLO_WIN: two graphic tools for sequence alignment and molecular phylogeny. *Comput Appl Biosci* **12**, 543–548.
- Glöckner, F. O., Fuchs, B. M. & Amann, R. (1999). Bacterioplankton compositions of lakes and oceans: a first comparison based on fluorescence in situ hybridization. *Appl Environ Microbiol* **65**, 3721–3726.
- Ivanova, E. P., Nedashkovskaya, O. I., Chun, J., Lysenko, A. M., Frolova, G. M., Svetashev, V. I., Vysotskii, M. V., Mikhailov, V. V., Huq, A. & Colwell, R. R. (2001). *Arenibacter* gen. nov., a new genus of the family *Flavobacteriaceae* and description of a new species, *Arenibacter latericius* sp. nov. *Int J Syst Evol Microbiol* **51**, 1987–1995.
- Kämpfer, P. & Kroppenstedt, R. M. (1996). Numerical analysis of fatty acid patterns of coryneform bacteria and related taxa. *Can J Microbiol* **42**, 989–1005.
- Khan, S. T., Nakagawa, Y. & Harayama, S. (2006). *Sediminicola luteus* gen. nov., sp. nov., a novel member of the family *Flavobacteriaceae*. *Int J Syst Evol Microbiol* **56**, 841–845.
- Kirchman, D. L. (2002). The ecology of *Cytophaga-Flavobacteria* in aquatic environments. *FEMS Microbiol Ecol* **39**, 91–100.
- Mesbah, M., Premachandran, U. & Whitman, W. B. (1989). Precise measurement of the G+C content of deoxyribonucleic acid by high-performance liquid chromatography. *Int J Syst Bacteriol* **39**, 159–167.
- MIDI (2001). *Sherlock Microbial Identification System*. Newark, DE: MIDI.
- Nedashkovskaya, O. I., Kim, S. B., Han, S. K., Lysenko, A. M., Rohde, M., Rhee, M.-S., Frolova, G. M., Falsen, E., Mikhailov, V. V. & Bae, K. S. (2004). *Maribacter* gen. nov., a new member of the family *Flavobacteriaceae*, isolated from marine habitats, containing the species *Maribacter sedimenticola* sp. nov., *Maribacter aquivivus* sp. nov., *Maribacter orientalis* sp. nov. and *Maribacter ulvicola* sp. nov. *Int J Syst Evol Microbiol* **54**, 1017–1023.
- Nedashkovskaya, O. I., Kim, S. B., Lee, K. H., Bae, K. S., Frolova, G. M., Mikhailov, V. V. & Kim, I. S. (2005). *Pibocella ponti* gen. nov., sp. nov., a novel marine bacterium of the family *Flavobacteriaceae* isolated from the green alga *Acrosiphonia sonderi*. *Int J Syst Evol Microbiol* **55**, 177–181.

Raguénès, G., Christen, R., Guezennec, J., Pignet, P. & Barbier, G. (1997). *Vibrio diabolicus* sp. nov., a new polysaccharide-secreting organism isolated from a deep-sea hydrothermal vent polychaete annelid, *Alvinella pompejana*. *Int J Syst Bacteriol* **47**, 989–995.

Reichenbach, H. (1989). The order *Cytophagales* Leadbetter 1974, 99^{AL}. In *Bergey's Manual of Systematic Bacteriology*, vol. 3, pp. 2011–2013. Edited by J. T. Staley, M. P. Bryant, N. Pfennig & J. G. Holt. Baltimore: Williams & Wilkins.

Saitou, N. & Nei, M. (1987). The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol* **4**, 406–425.

Simon, M., Glöckner, F. O. & Amann, R. (1999). Different community structure and temperature optima of heterotrophic picoplankton in various regions of the Southern Ocean. *Aquat Microb Ecol* **18**, 275–284.

Smibert, R. M. & Krieg, N. R. (1994). Phenotypic characterization. In *Methods for General and Molecular Bacteriology*, pp. 607–654. Edited by P. Gerhardt, R. G. E. Murray, W. A. Wood & N. R. Krieg. Washington, DC: American Society for Microbiology.

Thompson, J. D., Gibson, T. J., Plewniak, F., Jeanmougin, F. & Higgins, D. G. (1997). The CLUSTAL_X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res* **25**, 4876–4882.