Microlunatus panaciterrae sp. nov., a β -glucosidase-producing bacterium isolated from soil in a ginseng field

Dong-Shan An,¹ Wan-Taek Im¹ and Min-Ho Yoon²

Correspondence

Min-Ho Yoon
mhyoon@cnu.ac.kr

¹Department of Biological Sciences, Korea Advanced Institute of Science and Technology, 373-1 Guseong-dong, Yuseong-gu, Daejeon 305-701, Republic of Korea

²Department of BioEnvironmental Chemistry, College of Agriculture and Life Sciences, Chungnam National University, Gung-dong 22, Yuseong-gu, Daejeon 305-764, Republic of Korea

A novel bacterial strain, designated Gsoil 954^T, showing β -glucosidase activity was isolated from a soil sample from a ginseng field in Pocheon Province, South Korea. The isolate was a Gram-positive, aerobic, motile, coccus-shaped, non-endospore-forming bacterium. Comparative analysis of 16S rRNA gene sequences revealed that strain Gsoil 954^T belonged to the family *Propionibacteriaceae*, the highest levels of sequence similarity being found with the type strains of *Microlunatus ginsengisoli* (96.4 %) and *Microlunatus phosphovorius* (96.2 %). The strain showed <95.9 % similarity with respect to other species within the family *Propionibacteriaceae*. In addition, the presence of LL-2,6-diaminopimelic acid in the cell-wall peptidoglycan, MK-9(H₄) as the major menaquinone and anteiso-C_{15:0} and iso-C_{15:0} as the major cellular fatty acids suggested its affiliation with the genus *Microlunatus*. The G + C content of the genomic DNA was 65.1 mol%. On the basis of its phenotypic characteristics and phylogenetic distinctiveness, strain Gsoil 954^T represents a novel species within the genus *Microlunatus*, for which the name *Microlunatus panaciterrae* sp. nov. is proposed. The type strain is Gsoil 954^T (=KCTC 13058^T=DSM 18662^T).

During a study of bacterial populations that metabolize ginsenoside Rb1 to Rg3, Rh2 or compound K, a large number of β -glucosidase-producing novel strains from various sources, such as human faeces, environmental soils, wastewater sludge and different types of soil used for ginseng cultivation, were screened. One of these β -glucosidase-producing isolates, designated strain Gsoil 954^T (isolated from soil used for ginseng cultivation) was characterized in this study. On the basis of 16S rRNA gene sequence data, this strain was shown to represent a novel species of the genus *Microlunatus* in the family *Propionibacteriaceae*. The genus *Microlunatus* was first described on the basis of the characteristics of the type species of *Microlunatus phosphovorius* by Nakamura *et al.* (1995). At the time of writing, the genus also contained another recently described species, *Microlunatus ginsengisoli* (Cui *et al.*, 2007). In this study, we characterized a novel *Microlunatus*-like strain, Gsoil 954^T, which was isolated in South Korea from soil used for ginseng cultivation.

For strain isolation, a sample of soil used for ginseng cultivation was suspended in 50 mM phosphate buffer

(pH 7.0) and the suspension was spread on modified R2A agar plates (containing, 1⁻¹, 0.25 g tryptone, 0.25 g peptone, 0.25 g yeast extract, 0.125 g malt extract, 0.125 g beef extract, 0.25 g Casamino acids, 0.25 g soytone, 0.5 g glucose, 0.3 g soluble starch, 0.2 g xylan, 0.3 g sodium pyruvate, 0.3 g K₂HPO₄, 0.05 g MgSO₄, 0.05 g CaCl₂ and 15 g agar). The plates were incubated at 30 °C for 1 month. Single colonies were purified by transferring them onto new plates (modified R2A medium) for further incubation. Strain Gsoil 954^T was one of the isolates that appeared on the modified R2A agar plates under aerobic conditions. It was routinely cultured on modified R2A agar at 30 °C and maintained as a glycerol suspension (20 %, w/v) at –70 °C.

The Gram reaction was tested by using the non-staining method as described by Buck (1982). Cell morphology and motility were investigated with a Nikon light microscope ($\times 1000$ magnification), using cells grown on R2A agar for 3 days at 30 °C. Catalase and oxidase tests were performed as outlined by Cappuccino & Sherman (2002). Acid production from 49 carbon sources was tested at 30 °C with API 50 CH in combination with API 50 CHB/E medium (bioMérieux). Growth at a variety of temperatures (4, 15, 20, 25, 30, 37, 42 and 45 °C) was assessed on R2A agar. Growth at a variety of pH values was assessed in R2A

The GenBank/EMBL/DBJ accession number for the 16S rRNA gene sequence of strain Gsoil 954^T is AB271051.

broth (Difco). Utilization of substrates as sole carbon sources and some physiological characteristics were determined with API ID32 GN, API 20NE and API 20E galleries according to the instructions of the manufacturer (bioMérieux). Phosphate uptake was measured as described by Nakamura *et al.* (1995). Tests for degradation of DNA [DNase activity; using DNase agar (Scharlau) and flooding plates with 1 M HCl], casein, chitin, starch (Atlas, 1993), lipids (Kouker & Jaeger, 1987), xylan and cellulose (Ten *et al.*, 2004) were performed and evaluated after 10 days. Salt tolerance was tested with R2A broth supplemented with 1–10 % (w/v) NaCl after 5 days incubation. Growth on nutrient agar, trypticase soy agar and MacConkey agar was also evaluated at 30 °C.

Cells of strain Gsoil 954^T were aerobic, Gram-positive, non-endospore-forming, non-motile and coccus-shaped. Colonies grown on R2A agar plates for 5 days were smooth, circular, non-glossy, yellowish in colour and 1–2 mm in diameter. Physiological and biochemical characteristics of strain Gsoil 954^T are summarized in the species description. Selected characteristics that serve to differentiate strain Gsoil 954^T from related species of the genus *Microcylunatus* are shown in Table 1.

For measurement of the G + C content of the chromosomal DNA, genomic DNA of the novel strain was extracted and purified as described by Moore & Dowhan (1995), enzymically degraded into nucleosides and then determined as described by Mesbah *et al.* (1989) using reversed-phase HPLC. Isoprenoid quinones were extracted with chloroform/methanol (2:1, v/v), evaporated under a vacuum and re-extracted in n-hexane/water (1:1, v/v). The crude n-hexane-quinone solution was purified using Sep-Pak Vac silica cartridges (Waters) and subsequently analysed using HPLC as described previously (Hiraishi *et al.*, 1996). Cellular fatty acids were determined for strains harvested from R2A (Difco) agar plates after 5 days incubation. The cellular fatty acids were saponified, methylated and extracted according to the protocol of the Sherlock Microbial Identification System (MIDI). The fatty acids were analysed using a gas chromatograph (model 6890; Hewlett Packard) and were identified using the Microbial Identification software package (Sasser, 1990). Diaminopimelic acid isomers in the cell-wall peptidoglycan were determined by using TLC after hydrolysis with 6 M HCl at 100 °C for 18 h, as described by Komagata & Suzuki (1987). Polar lipids were extracted

Table 1. Characteristics that differentiate strain Gsoil 954^T from *M. ginsengisoli* and *M. phosphovor*

Strains: 1, Gsoil 954^T; 2, *M. ginsengisoli* Gsoil 633^T (data from Cui *et al.*, 2007); 3, *M. phosphovor* KCTC 9798^T (data from Nakamura *et al.*, 1995). All of the strains were Gram-positive, non-endospore-forming, non-motile, cocci-shaped bacteria. All of the strains could reduce nitrate to nitrite under aerobic conditions. All of the strains were positive for β-glucosidase activity and utilization of D-glucose, D-mannose, L-arabinose, maltose and melibiose as sole carbon sources. All of the strains were negative for the utilization of glycerol as a carbon source. All of the strains were positive for catalase activity. +, Positive; –, negative; DPG, diphosphatidylglycerol; PE, phosphatidylethanolamine; PG, phosphatidylglycerol; PI, phosphatidylinositol; PL, unidentified phospholipid(s).

Characteristic	1	2	3
Temperature range (°C)	20–30	20–30	5–35
Oxidase	+	–	+
Nitrate reduction under anaerobic conditions	–	–	+
Phosphate uptake	–	–	+
Utilization as a sole carbon source			
Acetate	–	–	+
Cellobiose	–	+	+
Dulcitol	–	+	+
Galactose	–	+	+
Glycogen	–	+	–
Inositol	–	–	+
Lactose	–	+	–
Mannitol	–	–	+
Starch	+	–	+
D-Xylose	–	+	+
Polar lipids	PG, DPG, PE	PG, DPG, PE, PL	PI, PG, DPG, PL*
DNA G + C content (mol%)	65.1	69.9	67.9

*Data from Maszenan *et al.* (1999).

and examined using two-dimensional TLC (Minnikin *et al.*, 1977).

The G+C content of genomic DNA of strain Gsoil 954^T was 65.1 mol%. MK-9(H₄) was the predominant respiratory menaquinone. The cell-wall peptidoglycan of strain Gsoil 954^T contained LL-diaminopimelic acid. The polar lipids detected were phosphatidylethanolamine, diphosphatidylglycerol and phosphatidylglycerol; phosphatidylethanolamine is rarely found in members of the family *Propionibacteriaceae* (Maszenan *et al.*, 1999), so this difference could serve to differentiate strain Gsoil 954^T from *Microlunatus phosphovorus* (Table 2). As shown in Table 2, iso-C_{15:0} and anteiso-C_{15:0} constitute more than 60 % of the total fatty acid content of strain Gsoil 954^T and showed levels similar to those of the type strains of *Microlunatus panaciterrae* and *M. phosphovorus*. However, the presence of C_{17:0} and C_{17:1}ω8c, the absence of iso-C_{14:0} and iso-C_{15:1} G and the existence of some quantitative differences in fatty acid composition served to distinguish strain Gsoil 954^T from the recognized members of the genus *Microlunatus* (Table 2).

PCR-mediated amplification of the 16S rRNA gene and sequencing of the purified PCR product were carried out as

Table 2. Cellular fatty acid profiles of strain Gsoil 954^T and related type strains of species of the genus *Microlunatus*

Strains: 1, Gsoil 954^T; 2, *M. ginsengisoli* Gsoil 633^T; 3, *M. phosphovorus* KCTC 9798^T. Data for all strains are from this study. All strains were grown on R2A agar for 5 days at 30 °C. ND, Not detected.

Fatty acid	1	2	3
Saturated			
C _{14:0}	ND	ND	0.5
C _{15:0}	0.8	0.9	1.9
C _{16:0}	ND	ND	0.7
C _{17:0}	2.1	0.8	ND
Unsaturated			
C _{17:1} ω8c	8.8	ND	ND
Branched-chain			
anteiso-C _{13:0}	ND	1.0	ND
iso-C _{14:0}	ND	8.2	8.9
iso-C _{15:0}	17.2	13.2	17.1
iso-C _{15:1} G	ND	2.1	ND
anteiso-C _{15:0}	53.1	48.8	46.0
anteiso-C _{15:1} A	2.6	0.9	ND
iso-C _{16:0}	6.7	ND	15.3
iso-C _{16:1} H	ND	ND	1.0
anteiso-C _{16:0}	ND	ND	ND
iso-C _{17:0}	2.7	ND	2.2
anteiso-C _{17:0}	3.1	2.6	1.2
Hydroxy			
iso-C _{14:0} 3-OH	ND	ND	3.93
Summed feature 4*	3.1	ND	1.3

*Summed feature 4 comprises C_{16:1}ω7c and/or iso-C_{15:0} 2-OH.

described by Kim *et al.* (2005). Almost-complete sequences of the 16S rRNA gene were compiled using SeqMan software (DNASTAR). The 16S rRNA gene sequences of representatives of related taxa were obtained from GenBank. Multiple alignments were performed with the CLUSTAL_X program (Thompson *et al.*, 1997), gaps were edited in the BioEdit program (Hall, 1999) and evolutionary distances were calculated using the Kimura two-parameter model (Kimura, 1983). Phylogenetic trees were constructed by using a neighbour-joining method (Saitou & Nei, 1987) and maximum parsimony (Fitch, 1971) with the MEGA3 program (Kumar *et al.*, 2004); bootstrap values were based on 1000 replications (Felsenstein, 1985).

An almost-complete 16S rRNA gene sequence (1454 bp) of strain Gsoil 954^T was determined and subjected to comparative analysis. Phylogenetic analysis based on 16S rRNA gene sequences indicated that strain Gsoil 954^T belongs to the genus *Microlunatus* (Fig. 1). The closest phylogenetic neighbours of strain Gsoil 954^T are *Microlunatus ginsengisoli* Gsoil 633^T (96.4 %) and *M. phosphovorus* DSM 10555^T (96.2 %). The phylogenetic distance from any other recognized species within the family *Propionibacteriaceae* was less than 95.9 %. There is general agreement that a DNA–DNA relatedness level of approximately 70 % is the threshold value for defining species (Wayne *et al.*, 1987). On the basis of the available data, organisms that share less than 97.0 % 16S rRNA gene sequence similarity will not show more than 60 % reassociation, irrespective of the hybridization method used (Stackebrandt & Goebel, 1994; Keswani & Whitman, 2001). Thus, the phylogenetic results for strain Gsoil 954^T demonstrated that the strain was not related to any of the recognized members of the genus *Microlunatus*.

In summary, the morphological, biochemical and chemotaxonomic properties of strain Gsoil 954^T are consistent with those described for the genus *Microlunatus*. On the basis of the relatively low 16S rRNA gene sequence similarities (<97 %) and the combination of unique phenotypic characteristics (Table 1), strain Gsoil 954^T represents a novel species of the genus *Microlunatus*, for which the name *Microlunatus panaciterrae* sp. nov. is proposed.

Description of *Microlunatus panaciterrae* sp. nov.

Microlunatus panaciterrae (pa.na.ci.ter'rae. N.L. n. *Panax -acis* scientific name for ginseng; L. n. *terra* soil; N.L. gen. n. *panaciterrae* of soil of a ginseng field).

Cells are Gram-positive, strictly aerobic, non-motile, coccus-shaped and 0.3–0.7 µm in diameter after 24 h culture on R2A agar plates. Colonies grown on R2A agar for 5 days are smooth, circular, non-glossy, yellowish, convex and 1–2 mm in diameter. Grows well at 20–30 °C and at pH 5.0–9.0, but does not grow at temperatures below 15 °C or above 37 °C. Grows on nutrient agar, but not on MacConkey agar or trypticase soy agar. Growth on R2A agar occurs in the absence of NaCl and in the presence

Fig. 1. Phylogenetic tree, based on 16S rRNA gene sequences, showing the positions of strain Gsoil 954^T and related species. This tree was constructed using the neighbour-joining method (Saitou & Nei, 1987) with a Kimura (1983) two-parameter distance matrix and pairwise deletion. Filled circles indicate branches that were also recovered by using maximum-parsimony algorithms. Bootstrap percentages (based on 1000 replications) >70 % are shown at branch points. Bar, 2 substitutions per 100 nucleotide positions.

of 5.0 % (w/v) NaCl, but not with 6.0 % (w/v) NaCl. Catalase and oxidase reactions are positive. H₂S is not produced. Nitrate is reduced under aerobic conditions, but nitrite is not reduced. Anaerobic growth does not occur; nitrate is not reduced under anaerobic conditions. In API 20E tests, negative results are obtained for urease, arginine dihydrolase, lysine decarboxylase, ornithine decarboxylase and tryptophan deaminase; positive results are obtained for β -galactosidase, β -glucosidase, gelatinase and in the Voges–Proskauer reaction. Acid is produced from amygdalin, L-arabinose, D-glucose, melibiose and L-rhamnose. Acid is not produced from inositol, D-mannitol, D-sorbitol or sucrose. The following compounds are utilized as sole carbon sources: D-arabinose, citrate, D-fucose, L-fucose, D-glucose, lactate, maltose, D-mannose, melibiose, L-rhamnose, D-ribose, salicin, D-sorbitol and sucrose. The following compounds are not utilized as sole carbon sources: N-acetylglucosamine, adipate, D-adonitol, alanine, starch, amygdalin, D-arabitol, L-arabitol, arbutin, caprate, cellobiose, dulcitol, erythritol, D-fructose, galactose, glycogen, glycerol, gluconate, histidine, 3-hydroxybenzoate, 4-hydroxybenzoate, 3-hydroxybutyrate, inositol, inulin, itaconate, 2-ketogluconate, 5-ketogluconate, D-lactose, D-lyxose, malate, malonate, D-mannitol, melezitose, methyl α -D-glucopyranoside, methyl α -D-mannopyranoside, methyl β -D-xylose, phenylacetate, proline, propionate, raffinose, serine, sodium acetate, L-sorbose, suberate, D-tagatose, trehalose, turanose, valerate, xylitol and L-xylose. Xylan, chitin, cellulose, casein, olive oil and DNA are not degraded, but starch is degraded. MK-9(H₄) is the predominant menaquinone. The major cellular fatty acids are anteiso-C_{15:0}, iso-C_{15:0}, iso-C_{17:0} and iso-C_{16:0}. The cell-wall peptidoglycan contains LL-diaminopimelic acid. The polar lipids detected are phosphatidylethanolamine, diphosphatidylglycerol and phosphatidylglycerol.

The type strain, Gsoil 954^T (=KCTC 13058^T=DSM 18662^T), was isolated from soil from a ginseng field in Pocheon Province, South Korea. The DNA G+C content of the type strain is 65.1 mol% (as determined by HPLC).

Acknowledgements

We thank Jean Euzéby for his help with the etymology of the species epithet.

References

- Atlas, R. M. (1993). *Handbook of Microbiological Media*. Edited by L. C. Parks. Boca Raton, FL: CRC Press.
- Buck, J. D. (1982). Nonstaining (KOH) method for determination of Gram reactions of marine bacteria. *Appl Environ Microbiol* **44**, 992–993.
- Cappuccino, J. G. & Sherman, N. (2002). *Microbiology: a Laboratory Manual*, 6th edn. San Francisco: Pearson Education, Inc. and Benjamin Cummings.
- Cui, Y.-S., Im, W.-T., Yin, C.-R., Yang, D.-C. & Lee, S.-T. (2007). *Micrococcus ginsengisoli* sp. nov. isolated from soil of a ginseng field. *Int J Syst Evol Microbiol* **57**, 713–716.
- Felsenstein, J. (1985). Confidence limit on phylogenies: an approach using the bootstrap. *Evolution* **39**, 783–791.
- Fitch, W. M. (1971). Toward defining the course of evolution: minimum change for a specific tree topology. *Syst Zool* **20**, 406–416.
- Hall, T. A. (1999). BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symp Ser* **41**, 95–98.
- Hiraishi, A., Ueda, Y., Ishihara, J. & Mori, T. (1996). Comparative lipoquinone analysis of influent sewage and activated sludge by high-performance liquid chromatography and photodiode array detection. *J Gen Appl Microbiol* **42**, 457–469.
- Keswani, J. & Whitman, W. B. (2001). Relationship of 16S rRNA sequence similarity to DNA hybridization in prokaryotes. *Int J Syst Evol Microbiol* **51**, 667–678.

- Kim, M. K., Im, W.-T., Ohta, H., Lee, M. & Lee, S.-T. (2005).** *Sphingopyxis granuli* sp. nov., a β -glucosidase-producing bacterium in the family *Sphingomonadaceae* in α -4 subclass of the *Proteobacteria*. *J Microbiol* **43**, 152–157.
- Kimura, M. (1983).** *The Neutral Theory of Molecular Evolution*. Cambridge: Cambridge University Press.
- Komagata, K. & Suzuki, K. (1987).** Lipids and cell-wall analysis in bacterial systematics. *Methods Microbiol* **19**, 161–207.
- Kouker, G. & Jaeger, K.-E. (1987).** Specific and sensitive plate assay for bacterial lipase. *Appl Environ Microbiol* **53**, 211–213.
- Kumar, S., Tamura, K. & Nei, M. (2004).** MEGA3: integrated software for molecular evolutionary genetics analysis and sequence alignment. *Brief Bioinform* **5**, 150–163.
- Maszenan, A. M., Seviour, R. J., Patel, B. K. C., Schumann, P., Burghardt, J., Webb, R. I., Soddell, J. A. & Rees, G. N. (1999).** *Friedmanniella spumicola* sp. nov. and *Friedmanniella capsulata* sp. nov. from activated sludge foam: Gram-positive cocci that grow in aggregates of repeating groups of cocci. *Int J Syst Bacteriol* **49**, 1667–1680.
- Mesbah, M., Premachandran, U. & Whitman, W. B. (1989).** Precise measurement of the G + C content of deoxyribonucleic acid by high-performance liquid chromatography. *Int J Syst Bacteriol* **39**, 159–167.
- Minnikin, D. E., Patel, P. V., Alshamaony, L. & Goodfellow, M. (1977).** Polar lipid composition in the classification of *Nocardia* and related bacteria. *Int J Syst Bacteriol* **27**, 104–117.
- Moore, D. D. & Dowhan, D. (1995).** Preparation and analysis of DNA. In *Current Protocols in Molecular Biology*, pp. 2–11. Edited by F. W. Ausubel, R. Brent, R. E. Kingston, D. D. Moore, J. G. Seidman, J. A. Smith & K. Struhl. New York: Wiley.
- Nakamura, K., Hiraishi, A., Yoshimi, Y., Kawaharasaki, M., Masuda, K. & Kamakata, Y. (1995).** *Micrococcus phosphovorus* gen. nov., sp. nov., a new gram-positive polyphosphate-accumulating bacterium isolated from activated sludge. *Int J Syst Bacteriol* **45**, 17–22.
- Saitou, N. & Nei, M. (1987).** The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol* **4**, 406–425.
- Sasser, M. (1990).** *Identification of bacteria by gas chromatography of cellular fatty acids*, MIDI Technical Note 101. Newark, DE: MIDI Inc.
- Stackebrandt, E. & Goebel, B. M. (1994).** Taxonomic note: a place for DNA-DNA reassociation and 16S rRNA sequence analysis in the present species definition in bacteriology. *Int J Syst Bacteriol* **44**, 846–849.
- Ten, L. N., Im, W.-T., Kim, M.-K., Kang, M.-S. & Lee, S.-T. (2004).** Development of a plate technique for screening of polysaccharide-degrading microorganisms by using a mixture of insoluble chromogenic substrates. *J Microbiol Methods* **56**, 375–382.
- Thompson, J. D., Gibson, T. J., Plewniak, F., Jeanmougin, F. & Higgins, D. G. (1997).** The CLUSTAL_X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res* **25**, 4876–4882.
- Wayne, L. G., Brenner, D. J., Colwell, R. R., Grimont, P. A. D., Kandler, O., Krichevsky, M. I., Moore, L. H., Moore, W. E. C., Murray, R. G. E. & other authors (1987).** International Committee on Systematic Bacteriology. Report of the ad hoc committee on reconciliation of approaches to bacterial systematics. *Int J Syst Bacteriol* **37**, 463–464.