Sanguibacter soli sp. nov., isolated from soil of a ginseng field

Myung Kyum Kim,¹ Rama Krishna Pulla,¹ Se-Young Kim,¹ Tae-Hoo Yi,¹ Nak-Kyun Soung² and Deok-Chun Yang¹

Correspondence
Deok-Chun Yang
dcyang@khu.ac.kr

¹Korean Ginseng Center and Ginseng Genetic Resource Bank, Kyung Hee University,
1 Seocheon-dong, Kiheung-gu, Yongin, Kyunggi-do 449-701, Republic of Korea

²Laboratory of Metabolism, Center for Cancer Research, National Cancer Institute, National
Institutes of Health, Bethesda, MD 20892, USA

A Gram-positive, non-spore-forming, rod-shaped, motile bacterium, designated strain DCY22^T, was isolated from soil of a ginseng field in South Korea and characterized in order to determine its taxonomic position. 16S rRNA gene sequence analysis revealed that strain DCY22^T belonged within the family *Sanguibacteraceae*, and highest levels of sequence similarity were found with *Sanguibacter marinus* 1-19^T (96.8%), *Sanguibacter suarezii* ST-26^T (96.0%), *Sanguibacter inulinus* ST-50^T (95.9%), *Sanguibacter keddiei* ST-74^T (95.5%), *Terrabacter terrae* PPLB^T (94.0%) and *Terrabacter tumescens* DSM 20308^T (93.8%). Chemotaxonomic investigations revealed that strain DCY22^T possessed menaquinone MK-9, a common feature of members of the genus *Sanguibacter*. Predominant fatty acids were unknown ECL 13.961 (45.81%), 17:0 anteiso (23.46%), 18:0 iso (15.42%) and unknown ECL 14.966 (8.70%). The results of physiological and biochemical tests clearly demonstrated that strain DCY22^T represents a novel species of the genus *Sanguibacter*, for which the name *Sanguibacter soli* sp. nov. is proposed. The type strain is DCY22^T (=KCTC 13155^T=JCM 14841^T).

The family *Sanguibacteraceae* was proposed by Stackebrandt & Schumann (2000) with *Sanguibacter* as the type genus. *Sanguibacter* remains the only described genus within the family. The genus *Sanguibacter* was proposed by Fernández-Garayzábal *et al.* (1995) and at the time of writing comprises four recognized species, namely *Sanguibacter inulinus* (Pascual *et al.*, 1996), *Sanguibacter keddiei* (Fernández-Garayzábal *et al.*, 1995), *Sanguibacter marinus* (Huang *et al.*, 2005) and *Sanguibacter suarezii* (Fernández-Garayzábal *et al.*, 1995).

In a series of studies, we have attempted to isolate micro-organisms from soil in order to investigate the community structure based on a culture-dependent method. In the present study, one strain, designated DCY22^T, was isolated from soil in a ginseng field and was characterized based on a polyphasic approach, including 16S rRNA gene sequence analysis, together with investigations of genomic relatedness and chemotaxonomic and phenotypic properties. The

results indicate that strain DCY22^T represents a novel species of the genus *Sanguibacter*.

Strain DCY22^T was isolated from surface soil of an agricultural field where ginseng was planted. One gram of the soil was immersed in 50 ml saline solution, vortexed and serially diluted and a 100 µl aliquot was inoculated onto one-tenth-strength R2A agar (Difco). The purified colonies were tentatively identified by using analysis of the partial 16S rRNA gene sequence. Cell morphology and motility were observed with a Nikon light microscope ($\times 1000$ magnification) after incubation on diluted Luria-Bertani (LB) agar (0.5% agar) for 1 day. Gram reactions were conducted according to the non-staining method as described by Buck (1982). Oxidase activity was evaluated via the oxidation of 1% *p*-aminodimethylaniline oxalate. Catalase activity was determined by measurements of bubble production after the application of 3% (v/v) hydrogen peroxide solution. Growth at various temperatures (4, 15, 25, 30, 37 and 42 °C) was assessed on R2A agar, and growth at different pH values (pH 5.0–11.0 at intervals of 0.5 pH units) was assessed in R2A broth. Growth on nutrient agar, LB agar and trypticase soy agar (TSA) was also evaluated at 30 °C. The API 20NE, API ID32 GN, API 50CH and API ZYM microtest systems were employed in these tests following the recommendations of the manufacturer (bioMérieux).

The GenBank/EMBL/DDBJ accession number for the 16S rRNA gene sequence of strain DCY22^T is EF547937.

A maximum-parsimony tree based on 16S rRNA gene sequences showing the phylogenetic relationships between strain DCY22^T and related species and a table detailing the cellular fatty acid profiles of strain DCY22^T and related species are available as supplementary material with the online version of this paper.

Table 1. Differential phenotypic characteristics between strain DCY22^T and type strains of recognized *Sanguibacter* species

Strains: 1, *Sanguibacter soli* sp. nov. DCY22^T; 2, *S. marinus* 1-19^T (data from Huang *et al.*, 2005); 3, *S. suarezii* ST-26^T (Fernández-Garayzábal *et al.*, 1995); 4, *S. inulinus* ST-50^T (Pascual *et al.*, 1996); 5, *S. keddiei* ST-74^T (Fernández-Garayzábal *et al.*, 1995). All are Gram-positive, motile and catalase-positive. All are able to produce acid from L-arabinose, gentiobiose, fructose, galactose, D-glucose, maltose, ribose, arbutin, glycerol, sucrose, D-mannose and D-xylose, but not from melezitose, D-adonitol or D-arabinose. All have MK-9(H₄) as the major menaquinone. +, Positive; –, negative; v, variable; w, weakly positive; NA, no data available.

Characteristic	1	2	3	4	5
Cell morphology	Rod	Rod/irregular	Rod/irregular	Irregular	Irregular
Colony colour	Yellow	Pale yellow	Pale yellow	Yellow	Pale yellow
Optimum growth temperature (°C)	37	25–30	25–30	25–30	25–30
Oxidase	+	–	–	–	–
Nitrate reduction	–	+	v	–	v
Acid from:					
Gluconate	–	NA	+	NA	–
5-Ketogluconate	–	–	+	–	–
Gentiobiose	+	–	+	+	+
D-Lyxose	–	–	+	+	+
Maltose	+	NA	+	+	+
Melibiose	–	–	+	+	+
Methyl α-D-glucoside	+	v	+	+	–
Methyl α-D-mannoside	–	–	+	+	–
Methyl β-D-xyloside	–	–	v	+	+
Raffinose	–	–	+	+	NA
L-Rhamnose	–	–	+	+	+
Ribose	–	NA	+	+	+
L-Sorbose	–	–	–	–	NA
Turanose	w	NA	+	+	+
L-Xylose	–	–	+	NA	NA
D-Arabitol	–	–	–	NA	NA
L-Arabitol	–	–	NA	NA	NA
Dulcitol (galactitol)	–	–	–	NA	NA
Erythritol	–	NA	–	–	–
Glycerol	w	+	+	+	+
Mannitol	–	NA	v	–	–
Sorbitol	–	–	–	+	–
Xylitol	–	–	–	NA	–
N-Acetylglucosamine	+	NA	+	+	–
Amygdalin	–	NA	+	+	+
Inulin	–	–	+	–	–
Glycogen	+	–	+	v	+
DNA G + C content (mol%)	69.8	73.4	70.5	70.0	69.5

Isoprenoid quinones were extracted with chloroform/methanol (2:1, v/v), purified by TLC and subsequently analysed by HPLC, as described by Collins & Jones (1981) and Shin *et al.* (1996). For fatty acid methyl ester analysis, the strain was allowed to grow on TSA for 48 h at 30 °C, and then two loops of well-grown cells were harvested. Fatty acid methyl esters were prepared, separated and identified by using the Sherlock Microbial Identification System (MIDI, Inc.; Sasser, 1990).

For determination of the G + C content, genomic DNA was extracted and purified with the Qiagen Genomic-tip system 100/G and was then enzymically degraded into nucleosides. The nucleosides were analysed by using HPLC as described by Tamaoka & Komagata (1984) and Mesbah *et al.* (1989).

The 16S rRNA gene was amplified from chromosomal DNA by using the universal bacterial primer set fd1 and rP1 (Weisburg *et al.*, 1991), and the purified PCR product was sequenced by Genotec (Daejeon, Korea) (Kim *et al.*, 2005). The full 16S rRNA gene sequence was compiled with SeqMan software (DNASTAR). The 16S rRNA gene sequences of related taxa were obtained from GenBank and edited by using the BioEdit program (Hall, 1999). Multiple alignments were performed with the CLUSTAL_X program (Thompson *et al.*, 1997). Evolutionary distances were calculated according to Kimura's two-parameter model (Kimura, 1983). Phylogenetic trees were constructed via the neighbour-joining method (Saitou & Nei, 1987) and maximum-parsimony method in the MEGA 2 program (Kumar *et al.*, 2001). Bootstrap analysis with 1000

Fig. 1. Neighbour-joining tree based on 16S rRNA gene sequences, showing the phylogenetic relationships between strain DCY22^T and related species (all recognized *Sanguibacter* species and type species of other related genera). Bar, 0.01 substitutions per nucleotide position.

replicates was also conducted in order to obtain confidence levels for the branches (Felsenstein, 1985). The closest relatives of strain DCY22^T, namely the type strains of all recognized *Sanguibacter* species, and the type species of genera representing all families in the suborder *Micrococcineae* were included in the phylogenetic tree.

Strain DCY22^T was cultured on R2A agar (Difco) at 30 °C, yielding yellow, circular colonies. Strain DCY22^T was found to be an aerobic, Gram-positive, motile, rod-shaped bacterium that was able to grow at 25–42 °C but not at 4 °C. Growth was observed at pH 5–9. The physiological characteristics of strain DCY22^T are summarized in the species description below, and a comparison of differential characteristics with the type strains of recognized *Sanguibacter* species is given in Table 1.

The cellular fatty acid profile of strain DCY22^T is presented in Supplementary Table S1 (available in IJSEM Online). The major cellular fatty acids found in strain DCY22^T were unknown ECL 13.961 (45.81%), 17:0 anteiso (23.46%), 18:0 iso (15.42%), unknown ECL 14.966 (8.70%), 12:0 anteiso (3.35%) and 18:3 ω 6,9,12c (2.63%). Strain DCY22^T lacked 15:0 anteiso and 16:0, both of which are common in recognized species of the genus *Sanguibacter* and in *Terrabacter terrae* and *Terrabacter tumescens*. The fatty acids 17:0 anteiso, 18:0 iso, unknown ECL 13.961 and ECL 14.966 in DCY22^T are not major components of the profiles of recognized *Sanguibacter* and *Terrabacter* species.

Strain DCY22^T contained a menaquinone with nine isoprene units (MK-9) as the predominant isoprenoid quinone and a smaller amount of MK-8. MK-9 is commonly found in recognized *Sanguibacter* species (Pascual *et al.*, 1996; Huang *et al.*, 2005).

The 16S rRNA gene sequence of strain DCY22^T was found to be a continuous stretch of 1395 nt. Based on preliminary data, strain DCY22^T was determined to belong to the class *Actinobacteria*, suborder *Micrococcineae*, family *Sanguibacteraceae*. Highest levels of 16S rRNA gene sequence similarity were found with *S. marinus* 1-19^T (96.8%), *S. suarezii* ST-26^T (96.0%), *S. inulinus* ST-50^T (95.9%), *S. keddii* ST-74^T (95.5%), *T. terrae* PPLB^T (94.0%) and *T. tumescens* DSM 20308^T (93.8%). In the neighbour-joining phylogenetic tree (Fig. 1), strain DCY22^T clearly fell within the *Sanguibacter* lineage. A maximum-parsimony tree is available as Supplementary Fig. S1.

The G + C content of the genomic DNA of strain DCY22^T was 69.8 mol%, a value similar to that reported for recognized *Sanguibacter* species (69.5–73.4 mol%). On the basis of phenotypic, chemotaxonomic and phylogenetic data, we conclude that strain DCY22^T represents a novel species of the genus *Sanguibacter*, for which the name *Sanguibacter soli* sp. nov. is proposed.

Description of *Sanguibacter soli* sp. nov.

Sanguibacter soli (so'li. L. neut. gen. n. *soli* of soil, the source of the type strain).

Cells are Gram-positive, aerobic, motile rods when grown on R2A agar (Difco) at 30 °C for 5 days. Colonies grown on R2A agar for 5 days are yellow. Optimal growth temperature is 37 °C. Grows at pH 5–9. Oxidase- and catalase-positive. Acid is produced from L-arabinose, cellobiose, D-fructose, D-galactose, gentiobiose, glycogen, lactose, maltose, D-mannose, methyl α -D-glucoside, starch, sucrose, trehalose, turanose, D-xylose, glycerol, N-acetylglucosamine, arbutin, aesculin and salicin, but not from

gluconate, 2-ketogluconate, 5-ketogluconate, D-arabinose, D- or L-fucose, D-lyxose, melezitose, melibiose, methyl α -D-mannoside, L-rhamnose, raffinose, ribose, L-sorbose, D-tagatose, L-xylose, D-adonitol (ribitol), D-arabitol, L-arabitol, dulcitol (galactitol), erythritol, inositol, mannitol, sorbitol, xylitol, amygdalin or inulin. Positive for N-acetyl- β -glucosaminidase, acid phosphatase, alkaline phosphatase, α -chymotrypsin, cystine arylamidase, esterase (C4), esterase (C8), β -galactosidase, α -glucosidase, β -glucosidase (aesculin hydrolysis), β -glucuronidase, leucine arylamidase, lipase (C14), α -mannosidase, naphthol-AS-BI-phosphohydrolase, trypsin and valine arylamidase. Negative for arginine dihydrolase, α -fucosidase, α -galactosidase, protease (gelatin hydrolysis) and urease. Assimilates acetate, L-arabinose, D-glucose, maltose, D-mannose, sucrose, glycogen and salicin, but not 3-hydroxybenzoate, 4-hydroxybenzoate, DL-3-hydroxybutyrate, 4-hydroxybenzoate, 2-ketogluconate, 5-ketogluconate, adipate, caprate, citrate, gluconate, itaconate, L-lactate, L-malate, malonate, phenyl acetate, propionate, suberate, n-valerate, L-fucose, melibiose, L-rhamnose, D-ribose, myo-inositol, D-mannitol, D-sorbitol, N-acetyl-D-glucosamine, L-alanine, L-histidine, L-proline or L-serine. The DNA G+C content of the type strain is 69.8 mol%, as determined by HPLC. The predominant quinone is MK-9(H₄). Major cellular fatty acids include unknown ECL 13.961 (45.81 %), 17:0 anteiso (23.46 %), 18:0 iso (15.42 %), unknown ECL 14.966 (8.70 %), 12:0 anteiso (3.35 %) and 18:3 ω 6,9,12c (2.63 %).

The type strain, DCY22^T (=KCTC 13155^T =JCM 14841^T), was isolated from soil of a ginseng field in South Korea.

Acknowledgements

This research was supported by the Kyung Hee University Research Fund in 2004 (KHU-20040204) and by a BK21 research fellowship from the Korean Ministry of Education and Human Resource Development.

References

- Buck, J. D. (1982). Nonstaining (KOH) method for determination of Gram reactions of marine bacteria. *Appl Environ Microbiol* **44**, 992–993.
- Collins, M. D. & Jones, D. (1981). Distribution of isoprenoid quinone structural types in bacteria and their taxonomic implications. *Microbiol Rev* **45**, 316–354.
- Felsenstein, J. (1985). Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* **39**, 783–791.
- Fernández-Garayzabal, J. F., Dominguez, L., Pascual, C., Jones, D. & Collins, M. D. (1995). Phenotypic and phylogenetic characterization of some unknown coryneform bacteria isolated from bovine blood and milk: description of *Sanguibacter* gen. nov. *Lett Appl Microbiol* **20**, 69–75.
- Hall, T. A. (1999). BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symp Ser* **41**, 95–98.
- Huang, Y., Dai, X., He, L., Wang, Y. N., Wang, B. J., Liu, Z. & Liu, S. J. (2005). *Sanguibacter marinus* sp. nov., isolated from coastal sediment. *Int J Syst Evol Microbiol* **55**, 1755–1758.
- Kim, M. K., Im, W.-T., Ohta, H., Lee, M. & Lee, S.-T. (2005). *Sphingopyxis granuli* sp. nov., a β -glucosidase-producing bacterium in the family *Sphingomonadaceae* in α -4 subclass of the *Proteobacteria*. *J Microbiol* **43**, 152–157.
- Kimura, M. (1983). *The Neutral Theory of Molecular Evolution*. Cambridge: Cambridge University Press.
- Kumar, S., Tamura, K., Jakobsen, I.-B. & Nei, M. (2001). MEGA2: molecular evolutionary genetics analysis software. *Bioinformatics* **17**, 1244–1245.
- Mesbah, M., Premachandran, U. & Whitman, W. B. (1989). Precise measurement of the G+C content of deoxyribonucleic acid by high-performance liquid chromatography. *Int J Syst Bacteriol* **39**, 159–167.
- Montero-Barrientos, M., Rivas, R., Velázquez, E., Monte, E. & Roig, M. G. (2005). *Terrabacter terrae* sp. nov., a novel actinomycete isolated from soil in Spain. *Int J Syst Evol Microbiol* **55**, 2491–2495.
- Pascual, C., Collins, M. D., Grimont, P. A. D., Dominguez, L. & Fernández-Garayzabal, J. F. (1996). *Sanguibacter inulinus* sp. nov. *Int J Syst Bacteriol* **46**, 811–813.
- Prauser, H., Schumann, P., Rainey, F. A., Kroppenstedt, R. M. & Stackebrandt, E. (1997). *Terracoccus luteus* gen. nov., sp. nov., an LL-diaminopimelic acid-containing coccoid actinomycete from soil. *Int J Syst Bacteriol* **47**, 1218–1224.
- Saitou, N. & Nei, M. (1987). The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol* **4**, 406–425.
- Sasser, M. (1990). *Identification of bacteria by gas chromatography of cellular fatty acids*, MIDI Technical Note 101. Newark, DE: MIDI, Inc.
- Shin, Y. K., Lee, J.-S., Chun, C. O., Kim, H.-J. & Park, Y.-H. (1996). Isoprenoid quinone profiles of the *Leclercia adecarboxylata* KCTC 1036^T. *J Microbiol Biotechnol* **6**, 68–69.
- Stackebrandt, E. & Schumann, P. (2000). Description of *Bogoriellaceae* fam. nov., *Dermacoccaceae* fam. nov., *Rarobacteraceae* fam. nov. and *Sanguibacteraceae* fam. nov. and emendation of some families of the suborder *Micrococccineae*. *Int J Syst Evol Microbiol* **50**, 1279–1285.
- Tamaoka, J. & Komagata, K. (1984). Determination of DNA base composition by reversed-phase high-performance liquid chromatography. *FEMS Microbiol Lett* **25**, 125–128.
- Thompson, J. D., Gibson, T. J., Plewniak, F., Jeanmougin, F. & Higgins, D. G. (1997). The CLUSTAL_X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res* **25**, 4876–4882.
- Weisburg, W. G., Barns, S. M., Pelletier, D. A. & Lane, D. J. (1991). 16S ribosomal DNA amplification for phylogenetic study. *J Bacteriol* **173**, 697–703.