Haliea salexigens gen. nov., sp. nov., a member of the *Gammaproteobacteria* from the Mediterranean Sea

Laurent Urios,^{1,2} Laurent Intertaglia,^{1,2} Françoise Lesongeur³ and Philippe Lebaron^{1,2}

Correspondence
Philippe Lebaron
lebaron@obs-banyuls.fr

¹Université Pierre et Marie Curie – Paris 6, Laboratoire ARAGO, Avenue Fontaulé, BP 44, F-66650 Banyuls-sur-Mer, France

²CNRS, UMR 7621, Laboratoire d'Océanographie Biologique de Banyuls, Avenue Fontaulé, BP 44, F-66650 Banyuls-sur-Mer, France

³Laboratoire de Microbiologie des Environnements Extrêmes, UMR 6197, IFREMER, Centre de Brest, BP 70, F-29280 Plouzané, France

A novel aerobic, Gram-negative bacterium, designated 3X/A02/235^T, was isolated from the surface of coastal waters in the north-western Mediterranean Sea. Cells were motile, straight rods, 1.6 µm long and 0.5 µm wide, and formed cream colonies on marine agar medium. The G + C content of the genomic DNA was 61 mol%. Phylogenetic analysis of the 16S rRNA gene sequence placed the strain in the class *Gammaproteobacteria* and within the family *Alteromonadaceae*. On the basis of 16S rRNA gene sequence comparisons and physiological and biochemical characteristics, this isolate represents a novel species of a novel genus, for which the name *Haliea salexigens* gen. nov., sp. nov. is proposed. The type strain of *Haliea salexigens* is 3X/A02/235^T (=DSM 19537^T =CIP 109602^T =MOLA 286^T).

The genus *Microbulbifer* was proposed by Gonzalez *et al.* (1997) for a Gram-negative, aerobic strain that was catalase- and oxidase-positive and possessed numerous vesicles derived from the outer membrane. Four species belonging to this genus have been described: *Microbulbifer hydrolyticus* (Gonzalez *et al.*, 1997), *Microbulbifer salipaludis* (Yoon *et al.*, 2003a), *Microbulbifer maritimus* (Yoon *et al.*, 2004) and *Microbulbifer elongatus* (Yoon *et al.*, 2003b). The latter species was originally named *Pseudomonas elongata* (Humm, 1946; Palleroni, 1984), but was transferred to the genus *Microbulbifer* on the basis of 16S rRNA gene sequence similarity, cellular fatty acid composition and respiratory quinone content. Consequently, major amounts of iso-15:0 and iso-17:1ω9c as well as ubiquinone Q-8 were defined as chemical markers of this genus (Yoon *et al.*, 2004). *Microbulbifer* species have been isolated from various saline and marine environments, including salt marshes, intertidal sediments and coastal waters. In this work, a strain named 3X/A02/235^T was isolated from seawater. On the basis of its 16S rRNA gene sequence, the closest relatives were species of the genus *Microbulbifer*, but the similarity values were only

90–91 %. Thus, the taxonomic position and characteristics of this novel strain were analysed in more detail. On the basis of the findings, a novel genus belonging to the family *Alteromonadaceae*, close to the genus *Microbulbifer*, is proposed for this strain.

Samples of the surface microlayer of seawater in the bay of Banyuls-sur-Mer (42° 29' N 3° 08' E) were collected in October 2002 by submerging a metal screen (Agogué *et al.*, 2004). Subsamples were spread on marine agar 2216 plates (Difco) and incubated at 25 °C for 2 weeks. Colonies were picked and purified after three subcultures. An isolate that formed cream-coloured colonies was obtained and designated strain 3X/A02/235^T (Agogué *et al.*, 2005).

Microscopic observations (AX70; Olympus) showed that cells from isolate 3X/A02/235^T were motile rods, approximately 1.6 ± 0.3 µm long and 0.5 ± 0.2 µm wide. Cells were negatively stained for transmission electron microscopy (Raguénès *et al.*, 1997) and were shown to possess single polar flagella (see Supplementary Fig. S1, available in IJSEM Online). The Ryu KOH reaction (Powers, 1995) led to immediate cell lysis that was confirmed by microscopy (AX70; Olympus). This positive reaction indicated that the strain was Gram-negative.

Strain 3X/A02/235^T was grown in marine broth 2216 (Difco). To determine the salinity range, marine broth 2216 was prepared according to the composition provided

The GenBank/EMBL/DDBJ accession number for the 16S rRNA gene sequence of strain 3X/A02/235^T is AY576769.

An electron micrograph of a negatively stained cell of strain 3X/A02/235^T and a figure showing the effects of temperature, pH and salinity on growth of the strain are available as supplementary material with the online version of this paper.

by the manufacturer, with the appropriate NaCl concentration. For determination of the pH range, MES, PIPES, AMPSO or MOPS (Sigma) was added to marine broth 2216 to achieve the appropriate pH. Cultures were incubated at 30 °C under aerobic conditions. The methods used for the determination of growth parameters were as reported by Wery *et al.* (2001b). Growth was observed at 10–37 °C, the optimum temperature being between 25 and 30 °C (see Supplementary Fig. S2, available in IJSEM Online). The strain grew at NaCl concentrations ranging from 7 to 70 g l⁻¹, and an optimum concentration could be defined at 40 g l⁻¹ (Supplementary Fig. S2). Growth occurred at pH 5.0–9.0, the optimum being pH 8.0. Growth decreased by 61 % at pH 9.0 relative to the value obtained at pH 8.0, whereas a relative decrease of only 19 % was observed at pH 6.0 (Supplementary Fig. S2). Anaerobic growth was checked using marine agar 2216 plates in an anaerobic jar under these optimal conditions. No growth was observed after 10 days and thus the strain should be considered as strictly aerobic.

The ability of isolate 3X/A02/235^T to use different substrates was investigated using Biolog GN2 microplates (Tang *et al.*, 1998) according to the manufacturer's instructions. Positive reactions were observed for Tweens 40 and 80, pyruvic acid methyl ester, succinic acid methyl ester, β -hydroxybutyric acid, α -ketovaleric acid, succinic acid, glutamic acid, glycyl L-glutamic acid and glycerol. D-Mannose, α -cyclodextrin, acetate, aspartate, succinamic acid and L-proline produced weakly positive reactions. A comparison between strain 3X/A02/235^T and its closest relatives is presented in Table 1.

Enzyme activities were investigated using the API ZYM system (bioMérieux) according to the manufacturer's instructions. Alkaline phosphatase, leucine arylamidase, valine arylamidase, acid phosphatase and naphthol-AS-BI-phosphohydrolase exhibited positive reactions (Table 1).

An analysis of the fatty acid methyl esters was performed by the Identification Service of the Deutsche Sammlung von Mikroorganismen und Zellkulturen (DSMZ, Braunschweig, Germany). The composition for isolate 3X/A02/235^T was as follows: 16:1 ω 7c (21.2 %), 18:1 ω 7c (17.5 %), 15:1 ω 6c (5.8 %), 17:1 ω 6c (2.7 %), 17:1 ω 8c (23.9 %), 10:0 3-OH (1.8 %), 11:0 3-OH (3.3 %), 12:0 3-OH (1.1 %), iso-11:0 3-OH (3.3 %), 11:0 (1.0 %), 12:0 (1.6 %), 13:0 (1.3 %), 14:0 (1.3 %), 15:0 (4.5 %), 16:0 (2.0 %) and 17:0 (9.3 %) (Table 2). Two major fatty acids found in all *Microbulbifer* strains, iso-15:0 and iso-17:1 ω 9c, were not detected in strain 3X/A02/235^T, while 17:1 ω 8c, which is the predominant fatty acid in strain 3X/A02/235^T, was poorly represented, or not detected, within the four *Microbulbifer* species. Fatty acids 11:0, 12:0, 13:0, 15:1 ω 6c and 12:0 3-OH, which were found in strain 3X/A02/235^T, were not detected in any *Microbulbifer* species. In contrast, fatty acids 10:0, iso-11:0, iso-15:1 and iso-17:0, which are always present in *Microbulbifer* species, were not detected in strain 3X/A02/235^T.

Analyses of respiratory quinones and polar lipids were carried out by the Identification Service of the DSMZ and Brian Tindall (DSMZ). Strain 3X/A02/235^T had a ubiquinone (Q-8) system and the polar lipids were diphosphatidylglycerol, phosphatidylglycerol and an undefined aminophospholipid.

Genomic DNA was extracted as described by Wery *et al.* (2001a). The G+C content was determined by thermal denaturation using the method of Marmur & Doty (1962) and the conditions described by Raguénès *et al.* (1997). The DNA G+C content of strain 3X/A02/235^T was 61.4 $\pm$ 0.2 mol%. The 16S rRNA gene was amplified and sequenced as described by Agogué *et al.* (2005) and the sequence was analysed as described by Urios *et al.* (2006). Strain 3X/A02/235^T was found to be phylogenetically affiliated to the family *Alteromonadaceae* in the class *Gammaproteobacteria* (Fig. 1). The closest relative was *Microbulbifer salipaludis* SM-1^T, with a similarity value of 91 %.

Strain 3X/A02/235^T is distinguishable from its closest phylogenetic relatives on the basis of differences in several phenotypic properties, as shown in Tables 1 and 2. Thus we propose that strain 3X/A02/235^T represents a novel species of a novel genus belonging to the *Alteromonadaceae*. Because of the marine origin of strain 3X/A02/235^T and its salinity requirement, the name *Haliea salexigens* gen. nov., sp. nov. is proposed.

Description of *Haliea* gen. nov.

Haliea (Ha'lie.a. N.L. fem. n. *Haliea* named after Halie, a sea nymph in Greek mythology, referring to the marine source of the first strain).

Motile Gram-negative rods. The major fatty acids are 17:1 ω 8c, 16:1 ω 7c, 18:1 ω 7c and 17:0. The ubiquinone is Q-8 and the polar lipids are diphosphatidylglycerol and phosphatidylglycerol. Phylogenetically affiliated with the class *Gammaproteobacteria* within the family *Alteromonadaceae*. The type species is *Haliea salexigens*.

Description of *Haliea salexigens* sp. nov.

Haliea salexigens (sa.le'xi.gens. L. n. *sal*, *salis* salt, seawater; L. v. *exigo* to demand; N.L. part. adj. *salexigens* seawater-demanding).

Displays the following properties in addition to those given in the genus description. Produces cream colonies on marine agar 2216. Cells are 1.6 $\pm$ 0.3 μ m long and 0.5 $\pm$ 0.2 μ m wide, with single polar flagella. The G+C content of the type strain is 61 mol%. Growth occurs at 10–37 °C (optimum, 25–30 °C), at pH 5.0–9.0 (optimum, pH 8.0) and at salinities in the range 7–70 g NaCl l⁻¹ (optimum, 42 g l⁻¹). Positive reactions with Biolog GN2 plates are obtained for Tweens 40 and 80, pyruvic acid methyl ester, succinic acid methyl ester, β -hydroxybutyric acid, α -ketovaleric acid, succinic acid, glutamic acid, glycyl

Table 1. Characteristics that distinguish strain 3X/A02/235^T from type strains of *Microbulbifer* species

Strains: 1, *M. maritimus* JCM 12187^T; 2, *M. hydrolyticus* DSM 11525^T; 3, *M. salipaludis* KCCM 41586^T; 4, *M. elongatus* DSM 6810^T; 5, strain 3X/A02/235^T. The quinone in all of the strains was Q-8. Data for reference type strains were taken from Gonzalez *et al.* (1997) and Yoon *et al.* (2003a, b, 2004). +, Positive; –, negative; (+), weakly positive; ND, no data available.

Characteristic	1	2	3	4	5
Sampling environment	Marine sediment	Salt marsh	Salt marsh	Sand and seawater	Seawater
DNA G+C content (mol%)	59.9	57.7	59	ND	61.4
Motility	–	–	–	+	+
Temperature range (optimum) (°C)	15–48 (37)	10–41 (37)	10–45 (37)	(25–30)	10–37 (25–30)
Salinity range (optimum) (g NaCl l ^{–1})	<10 (2–4)	<60 (6–30)	<10 (2–3)	(2–3)	7–70 (42)
pH range (optimum)	>5 (6.5–7.5)	6.5–8.5 (7.5)	>5 (7–8)	ND	5–9 (8)
Oxidase	+	+	+	ND	+
Catalase	ND	ND	+	+	+
API ZYM tests					
Alkaline phosphatase	+	ND	+	ND	+
Leucine arylamidase	+	ND	+	ND	+
Valine arylamidase	–	ND	–	ND	+
Acid phosphatase	+	ND	+	ND	+
Naphthol-AS-BI-phosphohydrolase	+	ND	+	ND	+
N-Acetyl-β-glucosaminidase	–	ND	+	ND	–
Esterase (C4)	+	ND	+	ND	–
Lipase (C8)	+	ND	+	ND	(+)
Substrates					
Arabinose	ND	ND	–	+	–
Cellobiose	+	+	+	+	–
Fructose	–	–	–	+	–
Galactose	ND	ND	–	+	–
Glucose	+	+	+	+	–
Lactose	–	–	–	+	–
Maltose	ND	ND	+	+	–
Mannose	–	–	–	+	(+)
Rhamnose	–	–	+	–	–
Raffinose	–	ND	–	–	–
Sucrose	–	–	–	+	–
Adonitol	–	ND	–	ND	–
Glycerol	ND	–	ND	–	+
Acetate	+	+	ND	+	–
Aspartate	ND	–	ND	(+)	+
Citrate	–	–	ND	–	–
Glutamate	–	+	ND	+	(+)
β-Hydroxybutyrate	ND	+	ND	ND	+
Lactate	–	ND	ND	+	–
Propionate	ND	+	ND	+	–
Pyruvate	+	+	ND	ND	+
Succinate	+	+	ND	+	+
Alanine	ND	+	ND	(+)	–
Leucine	ND	+	ND	+	+
Proline	ND	+	ND	(+)	+
Serine	+	+	ND	–	–
Tween 80	ND	–	+	ND	+

L-glutamic acid and glycerol. Positive API ZYM reactions are obtained for the following enzyme activities: alkaline phosphatase, leucine arylamidase, valine arylamidase, acid

phosphatase and naphthol-AS-BI-phosphohydrolase. Oxidase- and catalase-positive. The major fatty acids also include 15:1ω6c and 15:0.

Table 2. Cellular fatty acid compositions (%) of strain 3X/A02/235^T and type strains of the genus *Microbulbifer*

Strains: 1, *M. maritimus* JCM 12187^T; 2, *M. hydrolyticus* DSM 11525^T; 3, *M. salipaludis* KCCM 41586^T; 4, *M. elongatus* DSM 6810^T; 5, strain 3X/A02/235^T. –, Not detected. Data for reference type strains were taken from Yoon *et al.* (2004).

Fatty acid	1	2	3	4	5
10:0	1.3	1.7	2.4	1.6	–
11:0	–	–	–	–	1.0
12:0	–	–	–	–	1.6
13:0	–	–	–	–	1.3
14:0	1.0	1.2	2.6	0.7	1.3
15:0	1.6	1.5	1.7	0.9	4.5
16:0	8.7	11.4	16.3	7.1	2.0
17:0	1.3	2.9	2.2	2.5	9.3
18:0	–	1.6	1.4	1.2	–
17:0 cyclo	2.3	5.7	–	–	–
19:0ω8c	1.4	1.0	–	–	–
iso-11:0	10.0	5.7	4.8	6.5	–
iso-15:0	25.9	24.4	19.4	20.7	–
iso-15:1	0.8	1.0	0.7	1.0	–
iso-16:0	–	–	–	0.5	–
iso-17:0	6.9	10.4	5.5	9.9	–
anteiso-17:0	–	–	–	0.8	–
iso-17:1ω9c	12.6	10.1	9.5	11.3	–
15:1ω6c	–	–	–	–	5.8
17:1ω6c	–	–	–	–	2.7
17:1ω8c	–	0.5	1.0	1.8	23.9
18:1ω5c	–	–	0.7	–	–
18:1ω7c	5.6	8.9	11.8	16.3	17.5
10:0 3-OH	1.7	1.0	1.2	1.6	1.8
11:0 3-OH	–	–	–	–	3.3
12:0 3-OH	–	–	–	–	1.1
16:0 2-OH	–	–	0.9	–	–
iso-11:0 3-OH	14.2	6.2	5.7	7.7	3.3
iso-17:0 3-OH	–	–	0.9	–	–
16:1ω7c/iso-15:0 2-OH	2.2	2.7	7.1	6.0	21.2

The type strain, 3X/A02/235^T (=DSM 19537^T =CIP 109602^T =MOLA 286^T), was isolated from the surface microlayer of seawater from the bay of Banyuls-sur-Mer.

Acknowledgements

This work was supported by the Equipe Mixte de Recherche linking the Université Pierre et Marie Curie and the Centre National de la Recherche Scientifique to the Pierre Fabre Laboratories. The project was also carried out within the framework of the MarBEF Network of Excellence ('Marine Biodiversity and Ecosystem Functioning'), which is funded in the European Community's Sixth Framework Programme (contract no. GOCE-CT-2003-505446). This publication is contribution number MPS-07061 of MarBEF. It was also partly funded by the French program 'Bio-diversité et Changement Global – project: development of a coastal microbial observatory' from the 'Institut Français de la Biodiversité' (IFB–GICC, Paris, France).

Fig. 1. Unrooted neighbour-joining phylogenetic tree (Kimura corrections), based on 16S rRNA gene sequences, showing the position of strain 3X/A02/235^T. Bootstrap percentages (based on 100 replications) are shown at branch points. Bar, 2 substitutions per 100 nucleotide positions.

References

- Agoguè, H., Casamayor, E. O., Joux, F., Obernosterer, I., Dupuy, C., Lantoine, F., Catala, P., Weinbauer, M. G., Reinthaler, T. & other authors (2004). Comparison of samplers for the biological characterization of the sea surface microlayer. *Limnol Oceanogr Methods* 2, 213–225.
- Agoguè, H., Casamayor, E. O., Bourrain, M., Obernosterer, I., Joux, F., Herndl, G. J. & Lebaron, P. (2005). A survey on bacteria inhabiting the sea surface microlayer of coastal ecosystems. *FEMS Microbiol Ecol* 54, 269–280.
- Gonzalez, J. M., Mayer, F., Moran, M. A., Hodson, R. E. & Whitman, W. B. (1997). *Microbulbifer hydrolyticus* gen. nov., sp. nov., and *Marinobacterium georgiense* gen. nov., sp. nov., two marine bacteria from a lignin-rich pulp mill waste enrichment community. *Int J Syst Bacteriol* 47, 369–376.
- Humm, H. J. (1946). Marine agar-digesting bacteria of the South Atlantic coast. *Bull Duke Univ Mar Stn* 3, 45–75.
- Marmur, J. & Doty, P. (1962). Determination of the base composition of deoxyribonucleic acid from its thermal denaturation temperature. *J Mol Biol* 5, 109–118.
- Palleroni, N. J. (1984). Genus *Pseudomonas* Migula 1894. In *Bergey's Manual of Systematic Bacteriology*, vol. 1, pp. 141–199. Edited by N. R. Krieg & J. G. Holt. Baltimore: Williams & Wilkins.
- Powers, E. M. (1995). Efficacy of the Ryu nonstaining KOH technique for rapidly determining Gram reactions of food-borne and water-borne bacteria and yeasts. *Appl Environ Microbiol* 61, 3756–3758.
- Raguénès, G., Christen, R., Guezennec, J., Pignet, P. & Barbier, G. (1997). *Vibrio diabolicus* sp. nov., a new polysaccharide-secreting organism isolated from a deep-sea hydrothermal vent polychaete annelid, *Alvinella pompejana*. *Int J Syst Bacteriol* 47, 989–995.
- Tang, Y. W., Ellis, N. M., Hopkins, M. K., Smith, D. H., Dodge, D. E. & Persing, D. H. (1998). Comparison of phenotypic and genotypic techniques for identification of unusual aerobic pathogenic gram-negative bacilli. *J Clin Microbiol* 36, 3674–3679.

- Urios, L., Agogu, H., Lesongeur, F., Stackebrandt, E. & Lebaron, P. (2006).** *Balneola vulgaris* gen. nov., sp. nov., a member of the phylum *Bacteroidetes* from the north-western Mediterranean Sea. *Int J Syst Evol Microbiol* **56**, 1883–1887.
- Wery, N., Lesongeur, F., Pignet, P., Derennes, V., Cambon-Bonavita, M.-A., Godfroy, A. & Barbier, G. (2001a).** *Marinitoga camini* gen. nov., sp. nov., a rod-shaped bacterium belonging to the order *Thermotogales*, isolated from a deep-sea hydrothermal vent. *Int J Syst Evol Microbiol* **51**, 495–504.
- Wery, N., Moricet, J.-M., Cuff, V., Jean, J., Pignet, P., Lesongeur, F., Cambon-Bonavita, M.-A. & Barbier, G. (2001b).** *Caloranaerobacter azorensis* gen. nov., sp. nov., an anaerobic thermophilic bacterium isolated from a deep-sea hydrothermal vent. *Int J Syst Evol Microbiol* **51**, 1789–1796.
- Yoon, J.-H., Kim, I.-G., Shin, D.-Y., Kang, K. H. & Park, Y.-H. (2003a).** *Microbulbifer salipaludis* sp. nov., a moderate halophile isolated from a Korean salt marsh. *Int J Syst Evol Microbiol* **53**, 53–57.
- Yoon, J. H., Kim, I. G., Kang, K. H., Oh, T. K. & Park, Y. H. (2003b).** Transfer of *Pseudomonas elongata* Humm 1946 to the genus *Microbulbifer* as *Microbulbifer elongatus* comb. nov. *Int J Syst Evol Microbiol* **53**, 1357–1361.
- Yoon, J. H., Kim, I. G., Oh, T. K. & Park, Y. H. (2004).** *Microbulbifer maritimus* sp. nov., isolated from an intertidal sediment from the Yellow Sea, Korea. *Int J Syst Evol Microbiol* **54**, 1111–1116.