Sphingobacterium siyangense sp. nov., isolated from farm soil

Rui Liu,^{1†} Huan Liu,^{2,3†} Chong-Xing Zhang,¹ Shou-Yun Yang,¹
Xiu-Hong Liu,¹ Ke-Yun Zhang¹ and Ren Lai^{1,2}

Correspondence

Ren Lai

rlai72@njau.edu.cn

¹Key Laboratory of Microbiological Engineering of Agricultural Environment, Ministry of Agriculture, Life Sciences College of Nanjing Agricultural University, Nanjing, Jiangsu 210095, PR China

²Biotoxin Department of Key Laboratory of Animal Models and Human Disease Mechanisms, Kunming Institute of Zoology, Chinese Academy of Sciences, Kunming, Yunnan 650223, PR China

³Graduate School of the Chinese Academy of Sciences, Beijing 100009, China

The taxonomic position of a novel Gram-negative strain, designated SY1^T, isolated from a farm-soil sample obtained from Jiangsu Province, PR China, was characterized by using a polyphasic approach. The cells were non-motile, non-spore-forming rods. The organism grew optimally at 30–37 °C and at pH 6.0–8.0. Based on 16S rRNA gene sequence analysis, strain SY1^T is a member of the genus *Sphingobacterium*; *Sphingobacterium multivorum* JCM 21156^T was the nearest relative (98.5% sequence similarity). The predominant fatty acids of strain SY1^T were iso-C_{15:0} (32.9%), C_{16:0} (10.9%) and summed feature 3 (iso-C_{15:0} 2-OH and/or C_{16:1ω7c}; 24.1%). The DNA G+C content was 38.5 mol%. The low level of DNA–DNA relatedness (2.2%) to *S. multivorum* JCM 21156^T in combination with differential morphological and biochemical properties demonstrated that strain SY1^T (=KCTC 22131^T=CGMCC 1.6855^T) should be classified as representing a novel species of the genus *Sphingobacterium* for which the name *Sphingobacterium siyangense* sp. nov. is proposed.

The family *Sphingobacteriaceae* of the phylum *Bacteroidetes* (previously, the *Cytophaga–Flavobacterium–Bacteroides* group) currently comprises seven genera including *Sphingobacterium* and *Pedobacter* (Steyn *et al.*, 1998) and [*Flexibacter*] *canadensis* (a misclassified *Flexibacter* species). The type genus *Sphingobacterium* was proposed originally by Yabuuchi *et al.* (1983) to include Gram-negative rods that are positive for catalase and oxidase, negative for heparinase, gelatinase and indole production and having iso-C_{15:0}, iso-C_{15:0} 2-OH, C_{16:1ω7c} and C_{17:0} 3-OH as the main fatty acids. At the time of writing, the genus comprised eight recognized species, including *Sphingobacterium multivorum*, *S. spiritivorum* and *S. mizutaii* (Yabuuchi *et al.*, 1983) [*mizutae* was corrected to *mizutaii*; Holmes *et al.* (1988)], *S. antarcticum* (Shivaji *et al.*, 1992) [*antarcticus* was corrected to *antarcticum*; Euzéby (1998)], *S. thalpophilum* and *S. faecium* (Takeuchi & Yokota, 1992), *S. daejeonense* (Kim *et al.*, 2006) and *S. composti* (Yoo *et al.*, 2007).

A Gram-negative, aerobic, non-spore-forming bacterial strain (designated SY1^T) was isolated from a farm-soil sample and subjected to a polyphasic taxonomy approach. Based on the resulting data, strain SY1^T is considered to represent a novel species of the genus *Sphingobacterium*.

Strain SY1^T was isolated using the dilution-plating technique on Luria–Bertani (LB) agar (1^{–1}: 5 g yeast extract, 10 g tryptone, 10 g NaCl; pH 7.0) incubated at 30 °C and was cultivated routinely on LB agar or in LB broth at the same temperature under aerobic conditions. Stock cultures were maintained as a glycerol suspension (20%, w/v) at –70 °C.

Amplification of the 16S rRNA gene sequence was performed as described by Cui *et al.* (2001). The sequence obtained was compared with available sequences retrieved from GenBank using the BLAST program (<http://www.ncbi.nlm.nih.gov/blast/>) to determine an approximate phylogenetic affiliation. Phylogenetic analysis was performed using the software packages PHYLIP (Felsenstein, 1993) and MEGA version 3.1 (Kumar *et al.*, 2001) after multiple alignment of data using CLUSTAL_X (Thompson *et al.*, 1997). Evolutionary distances (distance options according to the Kimura two-parameter model; Kimura, 1980, 1983) and clustering were based on the neighbour-joining (Saitou & Nei, 1987) and maximum-likelihood (Felsenstein, 1981) methods. Bootstrap analysis (1000

†These authors contributed equally to this work.

The GenBank/EMBL/DDBJ accession number for the 16S rRNA gene sequence of strain SY1^T is EU046272.

Transmission electron micrographs of cells of strain SY1^T are available as supplementary material with the online version of this paper.

replications) was used to evaluate the topology of the neighbour-joining tree (Felsenstein, 1985).

An almost complete 16S rRNA gene sequence (1489 bp) of strain SY1^T was obtained. The phylogenetic tree (Fig. 1) based on the neighbour-joining algorithm showed that strain SY1^T grouped among *Sphingobacterium* species and joined *S. multivorum* JCM 21156^T (98.5 % similarity) with a high bootstrap value of 100 %. Strain SY1^T shared <96.0 % 16S rRNA gene sequence similarity with other recognized *Sphingobacterium* species.

DNA–DNA hybridization was carried out to evaluate the genomic DNA–DNA relatedness between strain SY1^T and *S. multivorum* JCM 21156^T by using the optical renaturation rate method as described by De Ley *et al.* (1970), with the modifications described by Huß *et al.* (1983) and Escara & Hutton (1980). Renaturation rates were computed with the program TRANSFER.BAS (Jahnke, 1992). Strain SY1^T exhibited low DNA–DNA relatedness with its closest relative *S. multivorum* JCM 21156^T (2.2 %). According to Stackebrandt & Goebel (1994) and Wayne *et al.* (1987), the levels obtained for 16 rRNA gene similarity (<97 %) and DNA–DNA relatedness (<70 %) support the genomic distinction of strain SY1^T from other *Sphingobacterium* species.

Morphological features were examined by using transmission electron microscopy (Hitachi-7650) as described by Nedashkovskaya *et al.* (2005). Gram staining and endospore-forming features were investigated using light microscopy (model BH 2; Olympus) according to the method described by Gerhardt *et al.* (1994). Anaerobic growth and growth at different temperatures and pH values were investigated according to Kim *et al.* (2006) and Xu *et al.* (2005), respectively, using LB as the basal medium. NaCl tolerance was tested using LB broth containing 0–5 % (w/v) NaCl. Growth was evaluated on cetrimide agar, Simmons' citrate agar, MacConkey agar and nutrient agar. All tests were incubated at 30 °C for 2 days. Oxidase activity was tested by determining the oxidation of 1 % (w/v) tetramethyl-*p*-phenylenediamine (Merck) and catalase activity was evaluated by determining the production of oxygen bubbles in a 5 % (v/v) aqueous hydrogen peroxide solution. Hydrolysis of casein, gelatin, aesculin, starch, carboxymethylcellulose, Tween 80 and Tween 20 was tested on LB agar after 2 days incubation as described by Lányi (1987) and Gerhardt *et al.* (1994). Degradation of DNA (using DNase agar from Scharlau, supplemented with 1 M HCl) and chitin was also investigated as described by Kim *et al.* (2006). Acid production from carbohydrates was determined using the medium and method described by

Fig. 1. Comparative analysis of the 16S rRNA gene sequences of strain SY1^T and representative strains of the genus *Sphingobacterium* using the neighbour-joining method. Bootstrap values (expressed as percentages of 1000 replications) >50 % are shown at branching points. The maximum-likelihood tree showed essentially the same topology (data not shown). Bar, 0.01 substitutions per nucleotide position.

Yamaguchi & Yokoe (2000). Single carbon-source utilization was determined as described by Zhou *et al.* (2007). Some other biochemical characteristics, as given in the species description below, were tested using API 20E kits (bioMérieux) according to the instructions of the manufacturer. Antibiotic sensitivity tests were performed using the diffusion method as described by Park *et al.* (2007) on LB agar at 30 °C with filter-paper discs (8 mm diameter; Sanofi Pasteur) containing the following antibiotics (µg): streptomycin (10), ampicillin (10), chloramphenicol (30), erythromycin (15), tetracycline (30), ofloxacin (5), rifampicin (5), gentamicin (10), ceftazidime (30), kanamycin (30), carbenicillin (100) and polymyxin B (30). After 2 days incubation, the diameters of the inhibition zones were measured. Phenotypic characteristics of strain SY1^T are given in the species description and in Table 1. Transmission electron micrographs of cells of strain SY1^T are shown in Supplementary Fig. S1 (available in IJSEM Online).

Genomic DNA was extracted and purified according to Yoon *et al.* (1996) and the DNA G+C content was determined using the thermal denaturation method (Mandel & Marmur, 1968). The G+C content of the DNA of strain SY1^T was 38.5 mol%, within the range of values reported for members of the genus *Sphingobacterium* (37.3–44.2 mol%; Yoo *et al.*, 2007).

For quantitative analysis of the cellular fatty acid content, cells were harvested after 2 days growth at 30 °C on trypticase soy agar and fatty acid methyl esters were prepared and identified by following the instructions of the Microbial Identification system (MIDI), as described by Sasser (1990). The cellular fatty acids mainly comprised iso-C_{15:0} (32.9%), C_{16:0} (10.9%) and summed feature 3 (iso-C_{15:0} 2-OH and/or C_{16:1}ω7c 24.1%), typical of members of the genus *Sphingobacterium* (Takeuchi & Yokota, 1992; Steyn *et al.*, 1998). The cellular fatty acids of strain SY1^T are listed in Table 2 and compared with those

Table 1. Differential characteristics of strain SY1^T and related *Sphingobacterium* species

Strains: 1, SY1^T (*S. siyangense* sp. nov.); 2, *S. multivorum* JCM 21156^T; 3, *S. thalpophilum* ATCC 43320^T; 4, *S. faecium* DSM 11690^T; 5, *S. mizutaii* ATCC 33299^T; 6, *S. daejeonense* LMG 23402^T; 7, *S. spiritivorum* ATCC 33861^T; 8, *S. composti* DSM 18850^T. Data were from Steyn *et al.* (1998), Kim *et al.* (2006), Yoo *et al.* (2007) and this study. +, Positive; –, negative; v, results vary between references; ND, no data available.

Characteristic	1	2	3	4	5	6	7	8
Growth at:								
4 °C	+	–	–	+	–	–	–	–
42 °C	+	–	+	–	–	+	–	+
Hydrolysis of:								
DNA	+	+	–	+	+	–	+	–
Starch	+	+	+	+	+	–	+	–
Aesculin	–	+	+	+	+	–	+	+
Tween 80	–	v	–	v	+	ND	+	+
Urea	+	+	+	+	+	–	+	–
Tyrosine	–	v	–	–	+	–	–	–
Lysine	+	–	–	–	v	–	–	ND
Utilization of:								
D-Ribose	+	–	–	–	–	–	–	–
L-Sorbose	+	–	–	–	–	–	–	–
L-Rhamnose	+	–	+	+	–	–	+	–
L-Arabinose	+	+	+	+	v	–	–	+
D-Mannitol	+	–	–	–	–	–	+	–
Xylitol	+	–	–	–	–	–	–	–
Adonitol	+	–	+	–	–	+	–	–
Inulin	+	v	+	+	–	–	v	–
Gluconate	+	–	–	–	–	–	–	–
Glycerol	+	–	+	+	–	–	v	–
Acid production from:								
D-Glucose	–	+	+	+	+	+	+	–
Sucrose	–	+	+	+	+	–	+	–
Trehalose	+	–	–	–	–	–	–	+
L-Rhamnose	+	v	+	–	–	–	–	–
L-Arabinose	–	+	+	+	+	–	–	+
Nitrate reduction	+	–	+	–	–	ND	–	–
DNA G+C content (mol%)	38.5	39.9–40.5	44.0–44.2	37.3	39.3–40.0	38.7	39.8	42.3

Table 2. Fatty acid contents (%) of strain SY1^T and related *Sphingobacterium* species

Strains: 1, SY1^T (*S. siyangense* sp. nov.); 2, *S. multivorum* JCM 21156^T; 3, *S. thalpophilum* ATCC 43320^T; 4, *S. faecium* DSM 11690^T; 5, *S. mizutaii* ATCC 33299^T; 6, *S. daejeonense* LMG 23402^T; 7, *S. spiritivorum* ATCC 33861^T; 8, *S. composti* DSM 18850^T. Data from Steyn *et al.* (1998), Kim *et al.* (2006), Yoo *et al.* (2007) and this study. Fatty acids amounting to less than 1.0 % in all strains tested are not listed. tr, Trace (<1.0 %); –, not detected; ECL, equivalent chain-length.

Fatty acid	1	2	3	4	5	6	7	8
C _{14:0}	3.9	2.7	3.2	tr	tr	–	1.0	–
anteiso-C _{15:0}	2.7	–	–	tr	tr	2.6	tr	–
iso-C _{15:0}	32.9	22.2	17.7	24.6	30.0	45.6	30.1	29.5
iso-C _{15:0} 3-OH	3.0	3.2	4.3	3.7	3.0	1.5	2.2	2.3
C _{16:0}	10.9	7.8	6.0	4.5	tr	3.4	3.5	2.2
C _{16:0} 2-OH	tr	tr	3.2	–	–	–	–	–
C _{16:0} 3-OH	6.4	5.3	6.3	2.1	tr	–	2.7	1.2
C _{16:0} 10-methyl	–	–	–	1.4	–	–	–	–
C _{16:1} ω5c	tr	–	–	1.5	tr	tr	tr	–
iso-C _{17:0} 3-OH	5.9	7.1	10.0	10.0	22.1	16.6	12.5	19.7
iso-C _{17:1} ω9c	1.1	tr	–	–	3.7	2.9	1.7	2.9
Summed feature 3*	24.1	49.0	47.8	48.1	35.1	23.8	42.7	37.5
ECL 13.566	tr	–	1.4	tr	1.3	1.0	tr	–

*Summed feature 3 contains iso-C_{15:0} 2-OH and/or C_{16:1}ω7c.

of the type strains of phylogenetically related *Sphingobacterium* species.

On the basis of phylogenetic evidence and phenotypic distinctiveness (Table 1), it is proposed that strain SY1^T should be classified in the genus *Sphingobacterium* as representing a novel species, for which the name *Sphingobacterium siyangense* sp. nov. is proposed.

Description of *Sphingobacterium siyangense* sp. nov.

Sphingobacterium siyangense (si.yang.en'se. N.L. neut. adj. *siyangense* pertaining to Siyang in Jiangsu Province, China, the city where the strain was isolated).

Cells are Gram-negative, non-motile, non-spore-forming, strictly aerobic rods, 0.8–1.7 µm long and 0.7–0.9 µm wide. After 2 days incubation on LB agar, colonies are 1.0–2.0 mm in diameter, slightly yellowish, convex, circular and smooth with entire margins. Grows at 4–42 °C (optimum, 30–37 °C), pH 3.0–10.0 (optimum, pH 6.0–8.0) and 0–4 % NaCl (optimum, 0–2 %). Growth occurs on nutrient agar and cetrimide agar, but not on Simmons' citrate agar or MacConkey agar. Positive for catalase and oxidase. In API 20E tests, positive for β-galactosidase, arginine dihydrolase, arginine decarboxylase, urease and nitrate reduction. Negative for ornithine decarboxylase, lysine decarboxylase, tryptophan deaminase, indole and H₂S production and Voges–Proskauer reaction. Starch, DNA, aesculin and Tween 20 are hydrolysed, but casein, gelatin, carboxymethylcellulose, Tween 80 or chitin are not. Acid is produced from trehalose and L-rhamnose, but not from D-glucose, cellobiose, D-galactose, sucrose, D-lactose, melezitose, melibiose, raffinose, D-mannitol, D-sorbitol or

inositol. Utilizes D-xylose, melibiose, turanose, maltose, D-fructose, D-glucose, D-lactose, sucrose, L-sorbose, trehalose, D-galactose, raffinose, L-rhamnose, L-arabinose, cellobiose, D-ribose, D-mannose, melezitose, i-erythritol, adonitol, L-arabitol, xylitol, D-sorbitol, D-mannitol, salicin, inulin, dextrin, N-acetyl-D-glucosamine, gluconate, lysine, amygdalin and glycerol, but not inositol, acetate, malonate or tyrosine. Resistant to streptomycin, kanamycin and polymyxin B, but sensitive to ceftazidime, chloramphenicol, erythromycin, tetracycline, ofloxacin, rifampicin, gentamicin, ampicillin and carbenicillin. The predominant fatty acids are iso-C_{15:0} (32.9 %), C_{16:0} (10.9 %) and summed feature 3 (iso-C_{15:0} 2-OH and/or C_{16:1}ω7c; 24.1 %). The G+C content of the DNA of the type strain is 38.5 mol%.

The type strain, SY1^T (=KCTC 22131^T=CGMCC 1.6855^T), was isolated from a soil sample from Jiangsu Province, PR China.

Acknowledgements

We are grateful for the financial support of the Chinese National Natural Science Foundation (30600001) and the Jiangsu Natural Sciences Foundation (BK2005422).

References

- Cui, X. L., Mao, P. H., Tseng, M., Li, W. J., Zhang, L. P., Xu, L. H. & Jiang, C. L. (2001). *Streptomonospora salina* gen. nov., a new member of the family Nocardioaceae. *Int J Syst Evol Microbiol* **51**, 357–363.
- De Ley, J., Cattoir, H. & Reynaerts, A. (1970). The quantitative measurement of DNA hybridization from renaturation rates. *Eur J Biochem* **12**, 133–142.

- Escara, J. F. & Hutton, J. R. (1980). Thermal stability and renaturation of DNA in dimethyl sulfoxide solutions: acceleration of renaturation rate. *Biopolymers* **19**, 1315–1327.
- Euzéby, J. P. (1998). Taxonomic note: necessary correction of specific and subspecific epithets according to Rules 12c and 13b of the International Code of Nomenclature of Bacteria (1990 Revision). *Int J Syst Bacteriol* **48**, 1073–1075.
- Felsenstein, J. (1981). Evolutionary trees from DNA sequences: a maximum likelihood approach. *J Mol Evol* **17**, 368–376.
- Felsenstein, J. (1985). Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* **39**, 783–791.
- Felsenstein, J. (1993). PHYLIP (phylogeny inference package), version 3.5c. Distributed by the author. Department of Genome Sciences, University of Washington, Seattle, USA.
- Gerhardt, P., Murray, R. G. E., Wood, W. A. & Krieg, N. R. (editors) (1994). *Methods for General and Molecular Bacteriology*. Washington, DC: American Society for Microbiology.
- Holmes, B., Weaver, R. E., Steigerwalt, A. G. & Brenner, D. J. (1988). A taxonomic study of *Flavobacterium spiritivorum* and *Sphingobacterium mizutae*: proposal of *Flavobacterium yabuuchiae* sp. nov. and *Flavobacterium mizutaii* comb. nov. *Int J Syst Bacteriol* **38**, 348–353.
- Huß, V. A. R., Festl, H. & Schleifer, K. H. (1983). Studies on the spectrophotometric determination of DNA hybridization from renaturation rates. *Syst Appl Microbiol* **4**, 184–192.
- Jahnke, K. D. (1992). BASIC computer program for evaluation of spectroscopic DNA renaturation data from GILFORD SYSTEM 2600 spectrophotometer on a PC/XT/AT type personal computer. *J Microbiol Methods* **15**, 61–73.
- Kim, K.-H., Ten, L. N., Liu, Q.-M., Im, W.-K. & Lee, S.-T. (2006). *Sphingobacterium daejeonense* sp. nov., isolated from a compost sample. *Int J Syst Evol Microbiol* **56**, 2031–2036.
- Kimura, M. (1980). A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequence. *J Mol Evol* **16**, 111–120.
- Kimura, M. (1983). *The Neutral Theory of Molecular Evolution*. Cambridge: Cambridge University Press.
- Kumar, S., Tamura, K., Jakobsen, I.-B. & Nei, M. (2001). MEGA2: Molecular Evolutionary Genetics Analysis software. *Bioinformatics* **17**, 1244–1245.
- Lányi, B. (1987). Classical and rapid identification methods for medically important bacteria. *Methods Microbiol* **19**, 1–67.
- Mandel, M. & Marmur, J. (1968). Use of ultraviolet absorbance-temperature profile for determining the guanine plus cytosine content of DNA. *Methods Enzymol* **12B**, 195–206.
- Nedashkovskaya, O. I., Kim, S. B., Suzuki, M., Shevchenko, L. S., Lee, M. S., Lee, K. H., Park, M. S., Frolova, G. M., Oh, H. W. & other authors (2005). *Pontibacter actiniarum* gen. nov., sp. nov., a novel member of the phylum 'Bacteroidetes', and proposal of *Reichenbachella* gen. nov. as a replacement for the illegitimate prokaryotic generic name *Reichenbachia* Nedashkovskaya *et al.* 2003. *Int J Syst Evol Microbiol* **55**, 2583–2588.
- Park, M., Ryu, S. H., Thi Vu, T.-H., Ro, H.-S., Yun, P.-Y. & Jeon, C. O. (2007). *Flavobacterium defluvii* sp. nov., isolated from activated sludge. *Int J Syst Evol Microbiol* **57**, 233–237.
- Saitou, N. & Nei, M. (1987). The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol* **4**, 406–425.
- Sasser, M. (1990). Identification of bacteria by gas chromatography of cellular fatty acids. *USFCC News* **10**, 16.
- Shivaji, S., Ray, M. K., Rao, N. S., Saisree, L., Jagannadham, M. V., Kumar, G. S., Reddy, G. S. N. & Bhargava, P. M. (1992). *Sphingobacterium antarcticus* sp. nov., a psychrotrophic bacterium from the soils of Schirmacher Oasis, Antarctica. *Int J Syst Bacteriol* **42**, 102–106.
- Stackebrandt, E. & Goebel, B. M. (1994). Taxonomic note: a place for DNA-DNA reassociation and 16S rRNA sequence analysis in the present species definition in bacteriology. *Int J Syst Bacteriol* **44**, 846–849.
- Steyn, P. L., Segers, P., Vancanneyt, M., Sandra, P., Kersters, K. & Joubert, J. J. (1998). Classification of heparinolytic bacteria into a new genus, *Pedobacter*, comprising four species: *Pedobacter heparinus* comb. nov., *Pedobacter piscium* comb. nov., *Pedobacter africanus* sp. nov. and *Pedobacter saltans* sp. nov. Proposal of the family *Sphingobacteriaceae* fam. nov. *Int J Syst Bacteriol* **48**, 165–177.
- Takeuchi, M. & Yokota, A. (1992). Proposals of *Sphingobacterium faecium* sp. nov., *Sphingobacterium piscium* sp. nov., *Sphingobacterium heparinum* comb. nov., *Sphingobacterium thalophilum* comb. nov. and two genospecies of the genus *Sphingobacterium*, and synonymy of *Flavobacterium yabuuchiae* and *Sphingobacterium spiritivorum*. *J Gen Appl Microbiol* **38**, 465–482.
- Thompson, J. D., Gibson, T. J., Plewniak, F., Jeanmougin, F. & Higgins, D. G. (1997). The CLUSTAL_X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res* **25**, 4876–4888.
- Wayne, L. G., Brenner, D. J., Colwell, R. R., Grimont, P. A. D., Kandler, O., Krichevsky, M. I., Moore, L. H., Moore, W. E. C., Murray, R. G. E. & other authors (1987). International Committee on Systematic Bacteriology. Report of the ad hoc committee on reconciliation of approaches to bacterial systematics. *Int J Syst Bacteriol* **37**, 463–464.
- Xu, P., Li, W.-J., Tang, S.-K., Zhang, Y.-Q., Chen, G.-Z., Chen, H.-H., Xu, L.-H. & Jiang, C.-L. (2005). *Naxibacter alkalitolerans* gen. nov., sp. nov., a novel member of the family 'Oxalobacteraceae' isolated from China. *Int J Syst Evol Microbiol* **55**, 1149–1153.
- Yabuuchi, E., Kaneko, T., Yano, I., Moss, C. W. & Miyoshi, N. (1983). *Sphingobacterium* gen. nov., *Sphingobacterium spiritivorum* comb. nov., *Sphingobacterium multivorum* comb. nov., *Sphingobacterium mizutae* sp. nov., and *Flavobacterium indologenes* sp. nov.: glucose-nonfermenting gram-negative rods in CDC groups IIk-2 and IIb. *Int J Syst Bacteriol* **33**, 580–598.
- Yamaguchi, S. & Yokoe, M. (2000). A novel protein-deamidating enzyme from *Chryseobacterium proteolyticum* sp. nov., a newly isolated bacterium from soil. *Appl Environ Microbiol* **66**, 3337–3343.
- Yoo, S.-H., Weon, H.-Y., Jang, H.-B., Kim, B.-Y., Kwon, S.-W., Go, S.-J. & Stackebrandt, E. (2007). *Sphingobacterium composti* sp. nov., isolated from cotton-waste composts. *Int J Syst Evol Microbiol* **57**, 1590–1593.
- Yoon, J.-H., Kim, H., Kim, S.-B., Kim, H.-J., Kim, W. Y., Lee, S. T., Goodfellow, M. & Park, Y.-H. (1996). Identification of *Saccharomonospora* strains by the use of genomic DNA fragments and rRNA gene probes. *Int J Syst Bacteriol* **46**, 502–505.
- Zhou, Y., Dong, J., Wang, X., Huang, X., Zhang, K.-Y., Zhang, Y.-Q., Guo, Y.-F., Lai, R. & Li, W.-J. (2007). *Chryseobacterium flavum* sp. nov., isolated from a polluted soil. *Int J Syst Evol Microbiol* **57**, 1765–1769.