Tropicibacter naphthalenivorans gen. nov., sp. nov., a polycyclic aromatic hydrocarbon-degrading bacterium isolated from Semarang Port in Indonesia

Theresia Umi Harwati,^{1,2} Yuki Kasai,¹ Yumiko Kodama,¹
Dwi Susilaningsih² and Kazuya Watanabe¹

Correspondence

Yuki Kasai
ykasai@kitasato-u.ac.jp

¹Marine Biotechnology Institute, 3-75-1 Heita, Kamaishi, Iwate 026-0001, Japan

²Research Centre for Biotechnology, Indonesia Institute of Sciences (LIPI), Cibinong 11691, Indonesia

An aerobic, Gram-negative, motile bacterium, strain C02^T, was isolated from seawater obtained from Semarang Port in Indonesia. Cells of strain C02^T were peritrichously flagellated and rod-shaped. Strain C02^T was able to degrade naphthalene, alkyl naphthalenes and phenanthrene. 16S rRNA gene sequence analysis revealed that this strain was affiliated with the family *Rhodobacteraceae* in the class *Alphaproteobacteria* and was related most closely to *Marinovum algicola* FF3^T (95.7 % similarity) and *Thalassobius aestuarii* JC2049^T (95.2 %). The DNA G + C content of strain C02^T was 64.6 mol%. The major cellular fatty acids were C_{18:1}ω7c (50.9 % of the total), C_{16:0} (17.9 %), 11 methyl C_{18:1}ω7c (14.7 %), C_{18:1}ω9c (2.9 %) and C_{19:0} cyclo ω8c (2.4 %), and the predominant respiratory lipoquinone was ubiquinone-10. Based on physiological, chemotaxonomic and phylogenetic data, strain C02^T is suggested to represent a novel species of a new genus, for which the name *Tropicibacter naphthalenivorans* gen. nov., sp. nov. is proposed. The type strain of *Tropicibacter naphthalenivorans* is C02^T (=JCM 14838^T=DSM 19561^T).

Contamination of the marine environment with petroleum hydrocarbons is of great public concern owing to their toxicity to humans and marine organisms (Malins *et al.*, 1985; Meador *et al.*, 1995). A number of hydrocarbon-degrading bacteria have been isolated from the marine environment and characterized (e.g. Kasai *et al.*, 2002a, b; Ozaki *et al.*, 2006), although information regarding hydrocarbon-degrading bacteria from tropical waters is relatively scarce (Chaillan *et al.*, 2004; Zhuang *et al.*, 2003; Zinjarde & Pant, 2002). We have recently isolated a substantial number of marine bacteria from seawater obtained from Semarang Port in Indonesia, and demonstrated that some of the isolates were capable of degrading hydrocarbons (Harwati *et al.*, 2007). The present study characterizes one of these Indonesian isolates, designated strain C02^T, affiliated with the class *Alphaproteobacteria*. Based on the results of polyphasic examinations, including phenotypic, chemotaxonomic and phylogenetic analyses, we propose that this strain represents a novel species of a new genus.

The ability of strain C02^T to degrade hydrocarbons in crude oil was examined in 10 ml ONR7a medium (Dyksterhouse *et al.*, 1995) supplemented with 1 mg heat-treated Arabian Light crude oil ml⁻¹ (Dutta &

Harayama, 2000). Cells of strain C02^T grown in 10 ml marine broth 2216 (MB; Difco) up to an optical density at 600 nm of approximately 1 were harvested by centrifugation (8000 g, 10 min), washed twice with ONR7a medium and inoculated on to the medium in 50-ml tubes fitted with Teflon-lined caps. The tubes were incubated at 30 °C on a reciprocal shaker (at 90 r.p.m.) for 4 weeks. Cultures were prepared in triplicate. Non-inoculated samples were incubated similarly and served as controls. Following incubation, oil components were extracted by using chloroform, and hydrocarbon losses were analysed via GC-MS (GC-MS-QP5000; Shimadzu) as described by Kasai *et al.* (2002b). The percentage biodegradation was calculated as described by Dutta & Harayama (2000). Strain C02^T degraded 88.1 ± 1.1 % (mean ± SE) of total naphthalene, 56.0 ± 10.4 % of total C₁-alkyl naphthalenes, 22.5 ± 8.6 % of total C₂-alkyl naphthalenes and 14.5 ± 2.7 % phenanthrene, while alkanes, C_{3–4}-alkyl naphthalenes, C_{0–4}-alkyldibenzothiophenes, C_{1–6}-alkylphenanthrenes and C_{0–3}-alkylfluorenes were not significantly degraded (<10 %) (C₀–C₆ represent total carbon numbers of branched alkyl groups).

The cell morphology of strain C02^T was examined by transmission electron microscopy (Beveridge *et al.*, 1994) and motility was examined under a phase-contrast microscope. Gram staining and oxidase and catalase tests

The GenBank/EMBL/DBJ accession number for the 16S rRNA gene sequence of strain C02^T is AB302370.

were performed according to the procedures of Smibert & Krieg (1994). Growth was tested at 30 °C in MB unless otherwise stated. Salinity requirements were tested by using modified MB (Sohn *et al.*, 2004) supplemented with 0–20 % (w/v) NaCl at 30 °C. The pH range for optimal growth was determined on solid media containing MB whose pH was adjusted to 5.5–9.5. Solid media contained 1.5 % (w/v) Bactoagar (Difco). The presence of poly- β -hydroxyalkanoate was detected by using Sudan Black according to the procedures of de Lima *et al.* (1998). Susceptibility to antibiotics was determined on agar plates containing MB (MA plates) in the presence of the following antibiotics (concentrations given in parentheses; $\mu\text{g ml}^{-1}$): ampicillin (50, 100, 150 and 200), chloramphenicol (20), gentamicin (50), kanamycin sulfate (20), nalidixic acid (20, 50, 100 and 200), spectinomycin (7.5, 15 and 20), streptomycin (20) and tetracycline (10). API ZYM, API 20NE (bioMérieux) and Microlog GN2 microplates (Biolog) were used for physiological and biochemical characterization according to the manufacturers' instructions.

Cells of strain C02^T were Gram-negative rods (1.1–3.4 μm in length and 0.1–0.7 μm in width), motile by means of peritrichous flagella (Fig. 1). Strain C02^T formed white colonies on MA plates. The strain was oxidase-positive but catalase-negative. It reduced nitrate to nitrite. Cells contained poly- β -hydroxyalkanoate. Growth of strain C02^T was observed between 10 and 43 °C, with optimum

Fig. 1. Transmission electron micrograph of a cell of strain C02^T. Bar, 1 μm .

growth at 37 °C. It grew within a pH range of 6.5–8.5, with optimum growth at pH 7.6. Strain C02^T showed essential requirements for NaCl, as no growth was observed in media lacking NaCl. It grew at NaCl concentrations from 1 to 15 %, with optimum growth at 5 % NaCl. The physiological and biochemical characteristics of strain C02^T are presented in detail in the species description below. It was susceptible to ampicillin, chloramphenicol, gentamicin, nalidixic acid, kanamycin, spectinomycin, streptomycin and tetracycline.

Table 1. Cellular fatty acid contents of strain C02^T and closely related taxa

Taxa: 1, strain C02^T; 2, *Antarctobacter heliothermus* EL-219^T (data from Labrenz *et al.*, 1998); 3, *Sagittula stellata* E-37^T (Gonzalez *et al.*, 1997); 4, *Ruegeria atlantica* 1480^T (Rüger & Höfle, 1992); 5, *Marinovum algicola* FF3^T (Lafay *et al.*, 1995; Martens *et al.*, 2006); 6, *Thalassobius aestuarii* JC2049^T (Yi & Chun, 2006); 7, *Roseovarius tolerans* EL-172^T (Labrenz *et al.*, 1999); 8, *Roseobacter denitrificans* DSM 7001^T (Labrenz *et al.*, 1999); 9, *Phaeobacter daeponensis* TF-218^T (Yoon *et al.*, 2007). Values are percentages of the total fatty acids; components that represented <0.5 % in all strains were omitted. ND, Not detected.

Fatty acid	1	2	3	4	5	6	7	8	9
Straight-chain									
C _{10:0}	ND	ND	ND	2.1	ND	ND	ND	ND	ND
C _{12:0}	ND	ND	ND	2.6	ND	ND	ND	ND	1.2
C _{16:0}	17.9	2.5	8.6	6.9	1.6	6.8	6.2	2.2	8.6
C _{17:0}	0.5	ND	ND	ND	ND	1.4	ND	ND	0.6
C _{18:0}	1.7	1.0	6.8	1.0	2.7	3.0	0.8	2.3	2.4
Unsaturated									
C _{16:1}	ND	0.8	ND	ND	ND	ND	0.8	ND	ND
C _{18:1ω7c}	50.9	83.2	*	71.2	86.7	68.0	70.2	87.7	57.7
C _{18:1ω9c}	2.9	ND	ND	ND	ND	ND	ND	ND	ND
Hydroxy									
C _{10:0} 3-OH	1.4	ND	ND	0.4	ND	1.3	ND	1.8	1.7
C _{12:0} 3-OH	0.7	ND	ND	4.7	1.6	ND	ND	ND	2.6
C _{12:1} 3-OH	1.2	3.1	3.6	ND	ND	ND	3.6	ND	ND
C _{14:0} 2-OH	0.5	ND	ND	ND	ND	ND	ND	2.9	ND
C _{16:0} 2-OH	ND	ND	ND	3.4	ND	3.8	ND	ND	5.6
11 Methyl C _{18:1ω7c}	14.7	ND	ND	7.1	3.2	12.0	ND	ND	16.6
C _{19:0} cyclo ω 8c	2.4	2.4	ND	ND	ND	ND	ND	ND	ND

*Major fatty acid, but not quantified.

Cellular fatty acids and quinone were analysed at the TechnoSuruga Laboratory Co., Ltd from cells grown in MB for 24 h. The major cellular fatty acids of strain C02^T were C_{18:1}ω7c (50.9 % of the total), C_{16:0} (17.9 %), 11 methyl C_{18:1}ω7c (14.7 %), C_{18:1}ω9c (2.9 %) and C_{19:0} cyclo ω8c (2.4 %) (detailed results are given in Table 1). The major lipoquinone was ubiquinone-10 (approximately 88 % of the total).

The 16S rRNA gene sequence of strain C02^T (1363 bp) was determined previously (Harwati *et al.*, 2007). Searches for similar 16S rRNA gene sequences were conducted by using the GenBank and RDP (Maidak *et al.*, 1999) databases. Phylogenetic analysis was performed by using CLUSTAL_X (version 1.83) (Thompson *et al.*, 1997), and a phylogenetic tree was constructed by using the neighbour-joining plot program within the MEGA software package (version 3.0) (Kumar *et al.*, 2004). In the neighbour-joining phylogen-

etic tree, strain C02^T formed a separate branch within the family *Rhodobacteraceae* (Fig. 2). 16S rRNA gene sequence analysis revealed that strain C02^T was related most closely to *Marinovum algicola* FF3^T (95.7 % similarity) and *Thalassobius aestuarii* JC2049^T (95.2 %). The latter two taxa belong to the *Roseobacter* clade, which is known as one of the most abundant groups in the marine environment (Buchan *et al.*, 2005). The DNA G+C content of strain C02^T as determined by the method of Katayama-Fujimura *et al.* (1984) was 64.6 mol%.

Phenotypic characteristics of strain C02^T that can be used to differentiate it from closely related members of the *Roseobacter* clade are detailed in Table 2. Based on phylogeny, fatty acid composition and phenotypic characteristics, we conclude that strain C02^T represents a novel species of a new genus, for which the name *Tropicibacter naphthalenivorans* gen. nov., sp. nov. is proposed.

Fig. 2. Phylogenetic relationships between strain C02^T and other members of the family *Rhodobacteraceae*. The tree was constructed by using the neighbour-joining algorithm. Numbers at nodes are bootstrap percentages based on 1000 replications; only values >50 % are shown. Bar, 1 % estimated sequence divergence.

Table 2. Differential phenotypic characteristics between strain C02^T and closely related taxa of the *Roseobacter* clade

Taxa: 1, strain C02^T; 2, *Antarctobacter heliothermus* EL-219^T (data from Labrenz *et al.*, 1998); 3, *Sagittula stellata* E-37^T (Gonzalez *et al.*, 1997); 4, *Ruegeria atlantica* 1480^T (Rüger & Höfle, 1992); 5, *Marinovum algicola* FF3^T (Lafay *et al.*, 1995; Martens *et al.*, 2006); 6, *Thalassobius aestuarii* JC2049^T (Yi & Chun, 2006); 7, *Roseovarius tolerans* EL-172^T (Labrenz *et al.*, 1999); 8, *Sulfitobacter dubius* KMM 3554^T (Ivanova *et al.*, 2004); 9, *Roseobacter denitrificans* Och 114^T (Shiba, 1991); 10, *Phaeobacter daeponensis* TF-218^T (Yoon *et al.*, 2007). +, Positive; –, negative; +/–, variable; ND, no data available; NF, not found; p, polar; sp, subpolar.

Characteristic	1	2	3	4	5	6	7	8	9	10
Cell morphology	Rod	Rod	Rod	Rod	Ovoid	Ovoid	Rod	Rod	Rod or ovoid	Ovoid
Pigmentation	None	Brown–yellow	Cream	Beige	Brown–beige	Cream	Red	Yellow	Red	None
Motility	+	+/–	+*	–	+	–	+/–	+	+	+
Flagella	Peritrichous	1–3, sp	+/?	–	1–2, sp	–	NF	1, sp	1–3, sp	1, p
Salt growth range (%)	1–15	1–10	ND†	1–11	0.6–12	1–7	1–13	1–12	ND	>0–8
pH optimum	7.6	6.9–7.8	7.5	7	7.5	7	6.2–9.0	7.5–8.0	7.0–8.0	7–8
Temperature growth range (°C)	10–43	ND	10–41	5–30	10–37	15–35	3–43.5	10–30	ND	4–42
Temperature optimum (°C)	37	16–26	30	25	25–30	35	8.5–33.5	25	20–30	37
NO ₃ [–] to NO ₂ [–]	+	+	–	+	–	–	–	+	+	+
Indole production	+	–	–	ND	–	–	–	ND	+	ND
β-Galactosidase	+	ND	ND	–	+	–	ND	ND	ND	+
DNA G + C content (mol%)	64.6	62.3–62.8	65.0	55	60	61	63.3–63.4	63.7	59.6	64.9

*Motility was assumed as flagella were found in suspension, but motile cells were not seen.

†Not determined, but required NaCl for growth (Gonzalez *et al.*, 1997).

Description of *Tropicibacter* gen. nov.

Tropicibacter [Trop.ic.i.bac.ter. L. adj. *tropicus* tropical, pertaining to the tropical zone of the Earth; N.L. masc. n. *bacter* (from Gr. n. *bakterion*) rod; N.L. masc. n. *Tropicibacter* a rod belonging to the tropical zone].

Cells are Gram-negative, motile by means of peritrichous flagella and rod-shaped (1.1–3.4 µm long and 0.1–0.7 µm wide). Require sodium ions for growth. Positive for oxidase and nitrate reduction. Contain poly-β-hydroxyalkanoate. The major ubiquinone is Q-10. The predominant fatty acids are C_{18:1ω7c}, C_{16:0} and 11 methyl C_{18:1ω7c}. The type species is *Tropicibacter naphthalenivorans*.

Description of *Tropicibacter naphthalenivorans* sp. nov.

Tropicibacter naphthalenivorans (naph.tha.le.ni.vo'rans. N.L. neut. N. *naphthalenum* naphthalene; L. part. adj. *vorans* devouring; N.L. part. adj. *naphthalenivorans* degrading naphthalene).

The description is identical to that for the genus, with the following additions. Colonies on MA are circular, slightly convex, smooth, yellowish–white and 2.0–3.0 mm in diameter after 3 days incubation at 37 °C. Growth occurs at temperatures of 10–43 °C (optimum 37 °C), at pH 6.5–8.5 and at NaCl concentrations of 1–15 % (optimum 5 %). Able to degrade C_{0–2}-alkylnaphthalene and phenanthrene. Susceptible to (µg ml^{–1}) ampicillin (50), chloramphenicol

(20), gentamicin (50), kanamycin (20), nalidixic acid (20), spectinomycin (7.5), streptomycin (20) and tetracycline (10). Positive for nitrate reduction, protease, esterase (C4), esterase lipase (C8), leucine arylamidase, valine arylamidase, acid phosphatase, naphthol-AS-BI-phosphohydrolase, α-galactosidase, β-galactosidase, α-glucosidase and β-glucosidase. Negative for glucose fermentation, arginine dihydrolase, alkaline phosphatase, cystine arylamidase, trypsin, chymotrypsin, β-glucuronidase, α-mannosidase and α-fucosidase. The following Biolog GN2 test substrates score as positive: dextrin, Tween 80, L-arabinose, D-fructose, D-glucose, D-lactose, maltose, D-mannose, D-sorbitol, sucrose, lactic acid, succinic acid, L-aspartic acid, L-alanine, L-proline, serine, inosine, uridine, turanose, xylitol, trehalose, inosine, uridine, cellobiose, melibiose, L-rhamnose and acetic acid. The predominant fatty acids are C_{18:1ω7c}, C_{16:0}, 11 methyl C_{18:1ω7c}, C_{18:1ω9c} and C_{19:0} cyclo ω8c.

The type strain, C02^T (=JCM 14838^T=DSM 19561^T), was isolated from seawater in Semarang Port, Java, Indonesia. The DNA G + C content of the type strain is 64.6 mol%.

Acknowledgements

We are grateful to Professor H. G. Trüper for his help with the genus and species names. We thank Hiromi Awabuchi and Midori Satoh for their technical assistance. This work was supported by the Japan International Cooperation Agency (JICA) and Indonesian Institute of Sciences (LIPI).

References

- Beveridge, T. J., Popkin, T. J. & Cole, R. M. (1994). Electron microscopy. In *Methods for General and Molecular Bacteriology*, pp. 42–71. Edited by P. Gerhardt, R. G. E. Murray, W. A. Wood & N. R. Krieg. Washington, DC: American Society for Microbiology.
- Buchan, A., Gonzalez, J. M. & Moran, M. A. (2005). Overview of the marine *Roseobacter* lineage. *Appl Environ Microbiol* **71**, 5665–5667.
- Chaillan, F., Le Fleche, A., Bury, E., Phantavong, Y. H., Grimont, P., Salot, A. & Oudot, J. (2004). Identification and biodegradation potential of tropical aerobic hydrocarbon-degrading microorganisms. *Res Microbiol* **155**, 587–595.
- de Lima, T. C. S., Grisi, B. M. & Bonato, M. C. M. (1999). Bacteria isolated from a sugarcane agroecosystem: their potential production of polyhydroxyalcanoates and resistance to antibiotics. *Rev Microbiol* **30**, 214–224.
- Dutta, T. K. & Harayama, S. (2000). Fate of crude oil by the combination of photo-oxidation and biodegradation. *Environ Sci Technol* **34**, 1500–1505.
- Dyksterhouse, S. E., Gray, J. P., Herwig, R. P., Cano Lara, J. & Staley, J. T. (1995). *Cyclocasticus pugetii* gen. nov., sp. nov., an aromatic hydrocarbon-degrading bacterium from marine sediments. *Int J Syst Bacteriol* **45**, 116–123.
- Gonzalez, J. M., Mayer, F., Moran, M. A., Hodson, R. E. & Whitman, W. B. (1997). *Sagittula stellata* gen. nov., sp. nov., a lignin-transforming bacterium from a coastal environment. *Int J Syst Bacteriol* **47**, 773–780.
- Harwati, T. U., Kasai, Y., Kodama, Y., Susilaningsih, D. & Watanabe, K. (2007). Characterization of diverse hydrocarbon-degrading bacteria isolated from Indonesian seawater. *Microbes Environ* **22**, 412–415.
- Ivanova, E. P., Gorshkova, N. M., Sawabe, T., Zhukova, N. V., Hayashi, K., Kurilenko, V. V., Alexeeva, Y., Buljan, V., Nicolau, D. V. & other authors (2004). *Sulfitobacter delicatus* sp. nov. and *Sulfitobacter dubius* sp. nov., respectively from starfish (*Stellaster equestris*) and sea grass (*Zostera marina*). *Int J Syst Evol Microbiol* **54**, 475–480.
- Kasai, Y., Kishira, H. & Harayama, S. (2002a). Bacteria belonging to the genus *Cyclocasticus* play a primary role in the degradation of aromatic hydrocarbons released in a marine environment. *Appl Environ Microbiol* **68**, 5625–5633.
- Kasai, Y., Kishira, H., Sasaki, T., Syutsubo, K., Watanabe, K. & Harayama, S. (2002b). Predominant growth of *Alkanivorax* strains in oil-contaminated nutrient-supplemented sea water. *Environ Microbiol* **4**, 141–147.
- Katayama-Fujimura, Y., Komatsu, Y., Kuraishi, H. & Kaneko, T. (1984). Estimation of DNA base composition by high performance liquid chromatography of its nuclease P1 hydrolysate. *Agric Biol Chem* **48**, 3169–3172.
- Kumar, S., Tamura, K. & Nei, M. (2004). MEGA3: integrated software for molecular evolutionary genetics analysis and sequence alignment. *Brief Bioinform* **5**, 150–163.
- Labrenz, M., Collins, M. D., Lawson, P. A., Tindall, B. J., Braker, G. & Hirsch, P. (1998). *Antarctobacter heliothermus* gen. nov., sp. nov., a budding bacterium from hypersaline and heliothermal Ekho Lake. *Int J Syst Bacteriol* **48**, 1363–1372.
- Labrenz, M., Collins, M. D., Lawson, P. A., Tindall, B. J., Schumann, P. & Hirsch, P. (1999). *Roseovarius tolerans* gen. nov., sp. nov., a budding bacterium with variable bacteriochlorophyll *a* production from hypersaline Ekho Lake. *Int J Syst Bacteriol* **49**, 137–147.
- Lafay, B., Ruimy, R., de Trautenberg, C. R., Breittmayer, V., Gauthier, M. J. & Christen, R. (1995). *Roseobacter algicola* sp. nov., a new marine bacterium isolated from the phycosphere of the toxin-producing dinoflagellate *Prorocentrum lima*. *Int J Syst Bacteriol* **45**, 290–296.
- Maidak, B. L., Cole, J. R., Parker, C. T., Jr, Garrity, G. M., Larsen, N., Li, B., Lilburn, T. G., McCaughey, M. J., Olsen, G. J. & other authors (1999). A new version of the RDP (Ribosomal Database Project). *Nucleic Acids Res* **27**, 171–173.
- Malins, D. C., Krahn, M. M., Brown, D. W., Rhodes, L. D., Myers, M. S., McCain, B. B. & Chan, S. L. (1985). Toxic chemicals in marine sediment and biota from Mukilteo, Washington: relationships with hepatic neoplasms and other hepatic lesions in English sole (*Parophrys vetulus*). *J Natl Cancer Inst* **74**, 487–494.
- Martens, T., Heidorn, T., Pukall, R., Simon, M., Tindall, B. J. & Brinkhoff, T. (2006). Reclassification of *Roseobacter gallaeciensis* Ruiz-Ponte et al. 1998 as *Phaeobacter gallaeciensis* gen. nov., comb. nov., description of *Phaeobacter inhibens* sp. nov., reclassification of *Ruegeria algicola* (Lafay et al. 1995) Uchino et al. 1999 as *Marinovum algicola* gen. nov., comb. nov., and emended descriptions of the genera *Roseobacter*, *Ruegeria* and *Leisingera*. *Int J Syst Evol Microbiol* **56**, 1293–1304.
- Meador, J. P., Stein, J. E., Reichert, W. L. & Varanasi, U. (1995). Bioaccumulation of polycyclic aromatic hydrocarbons by marine organisms. *Rev Environ Contam Toxicol* **143**, 79–165.
- Ozaki, S., Kishimoto, N. & Fujita, T. (2006). Isolation and phylogenetic characterization of microbial consortia able to degrade aromatic hydrocarbons at high rates. *Microbes Environ* **21**, 44–52.
- Rüger, H. J. & Höfle, M. G. (1992). Marine star-shaped-aggregate-forming bacteria: *Agrobacterium atlanticum* sp. nov.; *Agrobacterium meteori* sp. nov.; *Agrobacterium ferrugineum* sp. nov., nom. rev.; *Agrobacterium gelatinovororum* sp. nov., nom. rev.; and *Agrobacterium stellulatum* sp. nov., nom. rev. *Int J Syst Bacteriol* **42**, 133–143.
- Shiba, T. (1991). *Roseobacter litoralis* gen. nov., sp. nov., and *Roseobacter denitrificans* sp. nov., aerobic pink-pigmented bacteria which contain bacteriochlorophyll *a*. *Syst Appl Microbiol* **14**, 140–145.
- Smibert, R. M. & Krieg, N. R. (1994). Phenotypic characterization. In *Methods for General and Molecular Bacteriology*, pp. 607–655. Edited by P. Gerhardt, R. G. E. Murray, W. A. Wood & N. R. Krieg. Washington, DC: American Society for Microbiology.
- Sohn, J. H., Kwon, K. K., Kang, J.-H., Jung, H.-B. & Kim, S.-J. (2004). *Novosphingobium pentaromatovorans* sp. nov., a high-molecular-mass polycyclic aromatic hydrocarbon-degrading bacterium isolated from estuarine sediment. *Int J Syst Evol Microbiol* **54**, 1483–1487.
- Thompson, J. D., Gibson, T. J., Plewniak, K., Jeanmougin, F. & Higgins, D. G. (1997). The CLUSTAL_X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res* **25**, 4876–4882.
- Yi, H. & Chun, J. (2006). *Thalassobius aestuarii* sp. nov., isolated from tidal flat sediment. *J Microbiol* **44**, 171–176.
- Yoon, J.-H., Kang, S.-J., Lee, S.-Y. & Oh, T.-K. (2007). *Phaeobacter daeponensis* sp. nov., isolated from a tidal flat of the Yellow Sea in Korea. *Int J Syst Evol Microbiol* **57**, 856–861.
- Zhuang, W. Q., Tay, J. H., Maszenan, A. M. & Tay, S. T. (2003). Isolation of naphthalene-degrading bacteria from tropical marine sediments. *Water Sci Technol* **47**, 303–308.
- Zinjarde, S. S. & Pant, A. A. (2002). Hydrocarbon degraders from tropical marine environments. *Mar Pollut Bull* **44**, 118–121.