Sphingomonas sanxanigenens sp. nov., isolated from soil

Hai-Dong Huang,^{1,2} Wei Wang,¹ Ting Ma,¹ Guo-Qiang Li,¹
Feng-Lai Liang¹ and Ru-Lin Liu¹

Correspondence

Ting Ma

tingma@nankai.edu.cn

¹Key Laboratory of Molecular Microbiology and Technology, Ministry of Education, College of Life Sciences, Nankai University, Tianjin 300071, PR China

²Tianjin Agricultural University, Tianjin 300384, PR China

Strain NX02^T, a Gram-negative, non-spore-forming, rod-shaped bacterium, was isolated from soil, and its taxonomic position was investigated using a polyphasic approach. Chemotaxonomic analysis revealed that strain NX02^T possessed Q-10 as the predominant ubiquinone, *sym*-homospermidine as the major polyamine and C_{18:1}ω7c, C_{16:0} and C_{14:0} 2-OH as the major fatty acids. The main polar lipids were sphingoglycolipid, phosphatidylcholine, phosphatidylglycerol, diphosphatidylglycerol, phosphatidylethanolamine, phosphatidylmonomethylethanolamine, phosphatidylidimethylethanolamine and an unidentified glycolipid. The DNA G + C content was 66.4 mol%. Phylogenetic analysis based on 16S rRNA gene sequences showed that strain NX02^T belongs to the α-4 subgroup of the *Proteobacteria*, exhibiting the highest sequence similarity with respect to *Sphingomonas azotifigens* NBRC 15497^T (95.9%), *Sphingomonas pituitosa* DSM 13101^T (95.8%) and *Sphingomonas dokdonensis* KCTC 12541^T (95.8%). On the basis of these results, strain NX02^T represents a novel species of the genus *Sphingomonas sensu stricto*, for which the name *Sphingomonas sanxanigenens* sp. nov. is proposed. The type strain is NX02^T (=DSM 19645^T =CGMCC 1.6417^T).

The genus *Sphingomonas* was first proposed by Yabuuchi *et al.* (1990) and belongs to the family *Sphingomonadaceae* (Kosako *et al.*, 2000). The genus description has been emended by Takeuchi *et al.* (1993, 2001), Yabuuchi *et al.* (2002) and Busse *et al.* (2003). Analysis of the phylogeny, polyamine patterns and fatty acid profiles of members of the genus led to the subdivision of *Sphingomonas* into four separate genera, i.e. *Sphingomonas sensu stricto* and three new genera, *Sphingobium*, *Novosphingobium* and *Sphingopyxis*. Strains belonging to the family *Sphingomonadaceae* have great potential for biotechnological applications, e.g. in the bioremediation and degradation of refractory contaminants and in the production of valuable biopolymers referred to as sphingans. Sphingans, such as gellan (S-60), welan (S-130), rhamsan (S-194), S-88, heteropolysaccharide S-7, S-198, S-657, NW11, PS-P4, GS-1, I-886 and HWR1 (Matsuyama *et al.*, 2003; Seo *et al.*, 2004), have a similar structure, with a linear repeating tetrasaccharide that consists of glucose, glucuronic acid, rhamnose or mannose. Aqueous solutions of sphingans are very viscous or have the ability to form gels.

In this paper, we report the results from our polyphasic taxonomic study on a novel biopolymer-producing bacterium designated strain NX02^T. On the basis of the physiological, chemotaxonomic and phylogenetic data, strain NX02^T represents a novel species of the genus *Sphingomonas sensu stricto*.

Bacteria were isolated from topsoil collected from a cornfield in Xinhe County, PR China. Isolation was performed using the standard dilution plating technique at 30 °C on NK medium (containing, per litre distilled water: 15 g sucrose, 5 g peptone, 3 g beef powder, 1 g yeast extract, 15 g agar; pH 7.0). For purification, colonies of strain NX02^T, which had a distinct mucoid appearance, were picked and then transferred onto new plates. Strain NX02^T produces a novel extracellular biopolymer called sanxan gum, which has thickening and pseudoplastic qualities. Sanxan gum consists of carbohydrates and peptides, and the monosaccharide composition of the carbohydrate portion is similar to that of sphingans (Wang *et al.*, 2008).

A 16S rRNA gene sequence fragment corresponding to positions 6–1540 in the *Escherichia coli* numbering system (Brosius *et al.*, 1978) was amplified using the universal primers 27F (5'-GAGAGTTTGATCCTGGCTCAG-3') and 1541R (5'-AAGGAGGTGATCCAGCCGCA-3'). A phylogenetic analysis was performed using MEGA, version 3.0

The GenBank/EMBL/DDBJ accession number for the 16S rRNA gene sequence of strain NX02^T is DQ789172.

Results of two-dimensional TLC of the polar lipids of strain NX02^T are available as a supplementary figure with the online version of this paper.

(Kumar *et al.*, 2004), after multiple alignment of the data by CLUSTAL_X (Thompson *et al.*, 1997). Distances (with distance options according to the Kimura-2 model) and clustering using the neighbour-joining method were determined using bootstrap percentages based on 1000 replications. The 16S rRNA gene sequence of strain NX02^T was a continuous stretch of 1480 bp. A sequence similarity search of the GenBank/EMBL databases, performed using the FASTA program, revealed that strain NX02^T belongs to the family *Sphingomonadaceae*. The highest sequence similarities were found with respect to strains of established species, including *Sphingomonas azotifigens* NBRC 15497^T (95.9%), *Sphingomonas pituitosa* DSM 13101^T (95.8%) and *Sphingomonas dokdonensis* KCTC 12541^T (95.8%). The neighbour-joining phylogenetic tree (Fig. 1) showed that strain NX02^T clearly belonged to the genus *Sphingomonas sensu stricto* lineage, where it formed a separate branch. This branching pattern demonstrated that strain NX02^T represents a novel species within the genus *Sphingomonas sensu stricto*.

Genomic DNA was isolated and purified using the method of Cashion *et al.* (1977). The G + C content of the genomic DNA was analysed using HPLC after enzymic degradation of the DNA into nucleosides (Mesbah *et al.*, 1989; Tamaoka & Komagata, 1984). DNA-DNA hybridization was carried out using the optical renaturation method (De Ley *et al.*, 1970; Huß *et al.*, 1983; Jahnke, 1992) with a UV-Vis spectrophotometer (Perkin-Elmer) under optimal hybridization conditions (25 °C below the melting temperature of DNA). The DNA G + C content was determined to be 66.4 mol%. The DNA-DNA relatedness between strain NX02^T and *S. azotifigens* NBRC 15497^T was 24 %, i.e. below the value (70 %) considered as the threshold for the delineation of a genospecies (Stackebrandt & Goebel, 1994).

Cell morphology was observed under light and transmission electron microscopy, using cells grown for 3 days at 30 °C (Fig. 2). All other physiological and biochemical tests were performed as described by Li *et al.* (2004) and Lebuhr

Fig. 1. Neighbour-joining phylogenetic tree, based on 16S rRNA gene sequences, showing the relationships between strain NX02^T and related taxa. Bootstrap percentages (based on 1000 replicates) are shown at nodes: only values greater than 50 % are shown. The sequence of *Erythrobacter longus* DSM 6997^T was used as the outgroup. Bar, 1 % sequence divergence.

Fig. 2. Transmission electron micrograph of a cell of strain NX02^T. Bar, 500 nm.

et al. (2000). Growth occurred at 4–40 °C and at pH 4.0–8.5, with optima at 28–30 °C and pH 7.0–7.5. Growth occurred in the presence of 0–1.2 % (w/v) NaCl, with optimal growth at 0–0.05 % (w/v). Phenotypic features of strain NX02^T and phylogenetically related species are shown in Table 1. Strain NX02^T could be distinguished from its close relatives on the basis of a combination of phenotypic features. Strain NX02^T was similar to *Sphingomonas desiccabilis* DSM 16792^T in being positive for nitrate reduction, but it differed with regard to the oxidase reaction, the hydrolysis of starch and aesculin and the substrate-assimilation profile. Oxidase activity is absent in strain NX02^T and *S. pituitosa* DSM 13101^T. This is an unusual biochemical trait that has been reported for only a few sphingon-producing strains of the genus *Sphingomonas* (Yabuuchi *et al.*, 1990; Pollock, 1993).

Chemotaxonomic analyses were performed as follows. Respiratory quinones and polar lipids were determined as described by Tindall (1990a, b) and Altenburger *et al.* (1996), respectively, and were carried out by Dr B. J. Tindall and the Identification Service of the Deutsche Sammlung von Mikroorganismen und Zellkulturen (Braunschweig, Germany). Polyamines were extracted as described by Busse & Auling (1988) and analysed according to Busse *et al.* (1997). Fatty acid analysis was performed using the standard method of Sasser (1990): the results were compared with the database of fatty acids in the Sherlock Microbial Identification System (MIDI). The quinone system of strain NX02^T consisted of Q-10 (78 %) and Q-11 (22 %). The polar lipids detected were sphingoglycolipid, phosphatidylcholine, phosphatidylglycerol, diphosphatidylglycerol, phosphatidylethanolamine, phosphatidylmonomethylethanolamine, phosphatidyl dimethylethanolamine and an unidentified glycolipid (the two-dimensional thin-layer chromatogram is presented in Supplementary Fig. S1, available in IJSEM Online). The

Table 1. Phenotypic characteristics of strain NX02^T and the type strains of related *Sphingomonas* species

Strains: 1, NX02^T (data from this study); 2, *S. pituitosa* DSM 13101^T (Denner *et al.*, 2001); 3, *S. trueperi* ATCC 12417^T (Kämpfer *et al.*, 1997; Denner *et al.*, 2001); 4, *S. azotifigens* NBRC 15497^T (Xie & Yokota, 2006); 5, *S. dokdonensis* KCTC 12541^T (Yoon *et al.*, 2006); 6, *S. desiccabilis* DSM 16792^T (Reddy & Garcia-Pichel, 2007); 7, *S. molluscorum* CIP 109223^T (Romanenko *et al.*, 2007). +, Positive; –, negative; ND, no data available. All of the strains were positive for catalase activity and negative for indole production and assimilation of citrate.

Characteristic	1	2	3	4	5	6	7
Oxidase	–	–	+	+	+	+	+
Nitrate reduction	+	–	–	–	–	+	–
Hydrolysis of:							
Starch	–	–	+	+	+	+	–
Aesculin	+	+	–	+	+	–	+
Gelatin	+	–	–	–	+	+	–
Assimilation of:							
Acetate	–	+	+	+	+	–	ND
L-Arabinose	+	+	+	+	–	+	+
L-Aspartate	–	–	+	+	ND	–	ND
D-Fructose	–	–	+	+	–	–	ND
D-Glucose	+	+	+	+	–	+	+
L-Glutamic acid	–	ND	+	+	–	–	ND
Lactose	–	+	+	+	ND	–	+
Maltose	–	+	+	+	–	+	+
DNA G + C content (mol%)	66.4	64.5	65.6	66.0–68.0*	66.9	ND	68.3

*Range of values for three strains, including the type strain.

polyamine patterns of strain NX02^T showed a predominance of *sym*-homospermidine and trace amounts of spermidine. The presence of 2-hydroxy fatty acids and the absence of 3-hydroxy fatty acids are common characteristics of members of the family *Sphingomonadaceae* (Busse *et al.*, 1999; Takeuchi *et al.*, 2001). The cellular fatty acids of strain NX02^T mainly comprise C_{18:1}ω7c (53.8 %), C_{16:0} (15.1 %) and C_{14:0} 2-OH (12.0 %). This profile is consistent with that of the genus *Sphingomonas sensu stricto* (Busse *et al.*, 1999). However, strain NX02^T could be easily distinguished from its closest phylogenetic neighbours, both qualitatively and quantitatively, with regard to certain fatty acids and by the higher levels of C_{18:1}ω7c (Table 2).

On the basis of morphological and chemotaxonomic data and the results of the 16S rRNA gene sequence analysis, strain NX02^T represents a novel species of the genus *Sphingomonas*, for which the name *Sphingomonas sanxanigenens* sp. nov. is proposed.

Description of *Sphingomonas sanxanigenens* sp. nov.

Sphingomonas sanxanigenens [san.xa.ni.ge'nens. N.L. n. *sanxanum* sanxan gum (an extracellular biopolymer); L.

Table 2. Cellular fatty acid profiles of strain NX02^T and the type strains of related *Sphingomonas* species

Strains: 1, NX02^T (data from this study); 2, *S. pituitosa* DSM 13101^T (Denner *et al.*, 2001); 3, *S. trueperi* ATCC 12417^T (Kämpfer *et al.*, 1997); 4, *S. azotifigens* NBRC 15497^T (Xie & Yokota, 2006); 5, *S. dokdonensis* KCTC 12541^T (Yoon *et al.*, 2006); 6, *S. desiccabilis* DSM 16792^T (Reddy & Garcia-Pichel, 2007); 7, *S. molluscorum* CIP 109223^T (Romanenko *et al.*, 2007). Values are percentages of total fatty acids; –, not detected/not reported.

Fatty acid	1	2	3	4	5	6	7
C _{14:0}	0.9	–	–	2.6	0.8	0.8	–
C _{15:0}	–	2.6	–	–	–	0.9	1.7
C _{16:0}	15.1	11.5	9.8	24.5	19.0	13.4	24.0
C _{16:1} ω5c	1.6	–	–	–	1.3	1.6	–
C _{16:1} ω7c/iso-C _{15:0}	8.3	2.1	–	1.2	14.8	18.1	–
2-OH							
C _{17:1} ω6c	1.0	13.0	13.6	1.4	–	6.8	–
C _{18:1} ω5c	–	–	4.0	1.2	–	3.7	–
C _{18:1} ω7c	53.8	54.2*	64.2*	44.7	45.7	37.8	17.7†
C _{14:0} 2-OH	12.0	14.2	6.7	13.3	12.1	4.5	12.8
11-Methyl C _{18:1} ω7c	7.3	–	–	10.9	6.3	–	–

*Reported as summed feature 7 (one or more of C_{18:1}ω7c, C_{18:1}ω9t and C_{18:1}ω12t).

†Also reported to contain 16.5% C_{18:1}ω9.

part. adj. *genens* (from L. v. *genere* to produce) producing; N.L. part. adj. *sanxanigenens* sanxan gum-producing].

Cells are rod-shaped, 0.6–0.9 µm wide and 1.5–2.0 µm long and do not exhibit flagella. After 6 days incubation at 30 °C, colonies on NK medium are white, circular, convex, mucoid and 5–6 mm in diameter. Optimum growth occurs at pH 7.0–7.5, 28–30 °C and 0–0.05% NaCl. Cells are Gram-negative and non-spore-forming. Metabolism is non-fermentative. Positive for catalase and for nitrate reduction. Negative for oxidase, urease and for indole and H₂S production and in the methyl red and Voges–Proskauer tests. Positive for acid coagulation reaction in litmus milk. Hydrolyses aesculin and gelatin but not starch. The following are used as substrates for growth: L-arabinose, cellobiose, dextrin, D-galactose, gentiobiose, α-D-glucose, D-mannose, melibiose, pyruvic acid methyl ester, D-glucuronic acid, β-hydroxybutyric acid, sebacic acid, glycyl-L-glutamic acid, raffinose, sucrose, trehalose, bromosuccinic acid, succinamic acid and urocanic acid. The major respiratory lipoquinone is Q-10. The predominant polyamine is sym-homospermidine. The polar lipids contain sphingoglycolipid, phosphatidylcholine, phosphatidylglycerol, diphosphatidylglycerol, phosphatidylethanolamine, phosphatidylmonomethylethanolamine, phosphatidyltrimethylethanolamine and an unidentified glycolipid. The cellular fatty acids mainly comprise C_{18:1}ω7c, C_{16:0} and C_{14:0} 2-OH. The DNA G + C content of the type strain is 66.4 mol%.

The type strain, NX02^T (=DSM 19645^T =CGMCC 1.6417^T), was isolated from soil.

Acknowledgements

The authors are grateful to Dr J. P. Euzéby for his help with the nomenclature. We thank Mr Y. G. Zhou for his help with deposition of the novel strain. This work was supported by the RFDP of China (no. 20070055049).

References

- Altenburger, P., Kämpfer, P., Makristathis, A., Lubitz, W. & Busse, H.-J. (1996). Classification of bacteria isolated from a medieval wall painting. *J Biotechnol* **47**, 39–52.
- Brosius, J., Palmer, J. L., Kennedy, J. P. & Noller, H. F. (1978). Complete nucleotide sequence of a 16S ribosomal RNA gene from *Escherichia coli*. *Proc Natl Acad Sci U S A* **75**, 4801–4805.
- Busse, H.-J. & Auling, G. (1988). Polyamine pattern as a chemotaxonomic marker within the *Proteobacteria*. *Syst Appl Microbiol* **11**, 1–8.
- Busse, H.-J., Bunka, S., Hensel, A. & Lubitz, W. (1997). Discrimination of members of the family *Pasteurellaceae* based on polyamine patterns. *Int J Syst Bacteriol* **47**, 698–708.
- Busse, H.-J., Kämpfer, P. & Denner, E. B. M. (1999). Chemotaxonomic characterisation of *Sphingomonas*. *J Ind Microbiol Biotechnol* **23**, 242–251.
- Busse, H.-J., Denner, E. B. M., Buczolits, S., Salkinoja-Salonen, M., Bennisar, A. & Kämpfer, P. (2003). *Sphingomonas aurantiaca* sp. nov., *Sphingomonas aerolata* sp. nov. and *Sphingomonas faeni* sp. nov., air- and dustborne and Antarctic, orange-pigmented, psychrotolerant bacteria, and emended description of the genus *Sphingomonas*. *Int J Syst Evol Microbiol* **53**, 1253–1260.
- Cashion, P., Holder-Franklin, M. A., McCully, J. & Franklin, M. (1977). A rapid method for the base ratio determination of bacterial DNA. *Anal Biochem* **81**, 461–466.
- De Ley, J., Cattoir, H. & Reynaerts, A. (1970). The quantitative measurement of DNA hybridization from renaturation rates. *Eur J Biochem* **12**, 133–142.
- Denner, E. B. M., Paukner, S., Kämpfer, P., Moore, E. R. B., Abraham, W.-R., Busse, H.-J., Wanner, G. & Lubitz, W. (2001). *Sphingomonas pituitosa* sp. nov., an exopolysaccharide-producing bacterium that secretes an unusual type of sphingolipid. *Int J Syst Evol Microbiol* **51**, 827–841.
- Huß, V. A. R., Festl, H. & Schleifer, K. H. (1983). Studies on the spectrophotometric determination of DNA hybridization from renaturation rates. *Syst Appl Microbiol* **4**, 184–192.
- Jahnke, K. D. (1992). BASIC computer program for evaluation of spectroscopic DNA renaturation data from Gilford System 2600 spectrophotometer on a PC/XT/AT type personal computer. *J Microbiol Methods* **15**, 61–73.
- Kämpfer, P., Denner, E. B. M., Meyer, S., Moore, E. R. B. & Busse, H.-J. (1997). Classification of “*Pseudomonas azotocolligans*” Anderson 1955, 132, in the genus *Sphingomonas* as *Sphingomonas trueperi* sp. nov. *Int J Syst Bacteriol* **47**, 577–583.
- Kosako, Y., Yabuuchi, E., Naka, T., Fujiwara, N. & Kobayashi, K. (2000). Proposal of *Sphingomonadaceae* fam. nov., consisting of *Sphingomonas* Yabuuchi *et al.* 1990, *Erythrobacter* Shiba and Shimidu 1982, *Erythromicrobium* Yurkov *et al.* 1994, *Porphyrobacter* Fuerst *et al.* 1993, *Zymomonas* Kluyver and van Niel 1936, and *Sandaracinobacter* Yurkov *et al.* 1997, with the type genus *Sphingomonas* Yabuuchi *et al.* 1990. *Microbiol Immunol* **44**, 563–575.

- Kumar, S., Tamura, K. & Nei, M. (2004). MEGA3: integrated software for molecular evolutionary genetics analysis and sequence alignment. *Brief Bioinform* 5, 150–163.
- Lebuhn, M., Achouak, W., Schloter, M., Berge, O., Meier, H., Barakat, M., Hartmann, A. & Heulin, T. (2000). Taxonomic characterization of *Ochrobactrum* sp. isolates from soil samples and wheat roots, and description of *Ochrobactrum tritici* sp. nov. and *Ochrobactrum grignonense* sp. nov. *Int J Syst Evol Microbiol* 50, 2207–2223.
- Li, W. J., Zhang, Y. Q., Park, D. J., Li, C. T., Xu, L. H., Kim, C. J. & Jiang, C. L. (2004). *Duganella violaceinigra* sp. nov., a novel mesophilic bacterium isolated from forest soil. *Int J Syst Evol Microbiol* 54, 1811–1814.
- Matsuyama, H., Kamesaki, T., Sasaki, R., Minami, H. & Yumoto, I. (2003). Production of two types of exopolysaccharide by *Novosphingobium rosa*. *J Biosci Bioeng* 95, 152–156.
- Mesbah, M., Premachandran, U. & Whitman, W. B. (1989). Precise measurement of the G + C content of deoxyribonucleic acid by high-performance liquid chromatography. *Int J Syst Bacteriol* 39, 159–167.
- Pollock, T. J. (1993). Gellan-related polysaccharides and the genus *Sphingomonas*. *J Gen Microbiol* 139, 1939–1945.
- Reddy, G. S. N. & Garcia-Pichel, F. (2007). *Sphingomonas mucosissima* sp. nov. and *Sphingomonas desiccabilis* sp. nov., from biological soil crusts in the Colorado Plateau, USA. *Int J Syst Evol Microbiol* 57, 1028–1034.
- Romanenko, L. A., Uchino, M., Frolova, G. M., Tanaka, N., Kalinovskaya, N. I., Latyshev, N. & Mikhailov, V. V. (2007). *Sphingomonas molluscorum* sp. nov., a novel marine isolate with antimicrobial activity. *Int J Syst Evol Microbiol* 57, 358–363.
- Sasser, M. (1990). Identification of bacteria by gas chromatography of cellular fatty acids. *USFCC Newsl* 20, 16.
- Seo, E. J., Yoo, S. H., Oh, K. W., Cha, J., Lee, H. G. & Park, C. S. (2004). Isolation of an exopolysaccharide-producing bacterium, *Sphingomonas* sp. CS101, which forms an unusual type of sphingan. *Biosci Biotechnol Biochem* 68, 1146–1148.
- Stackebrandt, E. & Goebel, B. M. (1994). Taxonomic note: a place for DNA-DNA reassociation and 16S rRNA sequence analysis in the present species definition in bacteriology. *Int J Syst Bacteriol* 44, 846–849.
- Takeuchi, M., Kawai, F., Shimada, Y. & Yokota, A. (1993). Taxonomic study of polyethylene glycerol-utilizing bacteria: emended description of the genus *Sphingomonas* and new descriptions of *Sphingomonas macrogoltabidus* sp. nov., *Sphingomonas sanguis* sp. nov. and *Sphingomonas terrae* sp. nov. *Syst Appl Microbiol* 16, 227–238.
- Takeuchi, M., Hamana, K. & Hiraiishi, A. (2001). Proposal of the genus *Sphingomonas sensu stricto* and three new genera, *Sphingobium*, *Novosphingobium* and *Sphingopyxis*, on the basis of phylogenetic and chemotaxonomic analyses. *Int J Syst Evol Microbiol* 51, 1405–1417.
- Tamaoka, J. & Komagata, K. (1984). Determination of DNA base composition by reversed-phase high-performance liquid chromatography. *FEMS Microbiol Lett* 25, 125–128.
- Thompson, J. D., Gibson, T. J., Plewniak, F., Jeanmougin, F. & Higgins, D. G. (1997). The CLUSTAL_X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res* 25, 4876–4882.
- Tindall, B. J. (1990a). Lipid composition of *Halobacterium lacusprofundi*. *FEMS Microbiol Lett* 66, 199–202.
- Tindall, B. J. (1990b). A comparative study of the lipid composition of *Halobacterium saccharovorum* from various sources. *Syst Appl Microbiol* 13, 128–130.
- Wang, W., Huang, H. D., Zhang, Y., Ma, T., Zhang, G. P., Liang, F. L. & Liu, R. L. (2008). Rheological and gelling properties of a novel biopolymer. *Microbiology (Beijing)* 35, 866–871 (in Chinese).
- Xie, C. H. & Yokota, A. (2006). *Sphingomonas azotifigens* sp. nov., a nitrogen-fixing bacterium isolated from the roots of *Oryza sativa*. *Int J Syst Evol Microbiol* 56, 889–893.
- Yabuuchi, E., Yano, I., Oyaizu, H., Hashimoto, Y., Ezaki, T. & Yamamoto, H. (1990). Proposals of *Sphingomonas paucimobilis* gen. nov. and comb. nov., *Sphingomonas parapaucimobilis* sp. nov., *Sphingomonas yanoikuyae* sp. nov., *Sphingomonas adhaesiva* sp. nov., *Sphingomonas capsulata* comb. nov., and two genospecies of the genus *Sphingomonas*. *Microbiol Immunol* 34, 99–119.
- Yabuuchi, E., Kosako, Y., Fujiwara, N., Naka, T., Matsunaga, I., Ogura, H. & Kobayashi, K. (2002). Emendation of the genus *Sphingomonas* Yabuuchi *et al.* 1990 and junior objective synonymy of the species of three genera, *Sphingobium*, *Novosphingobium* and *Sphingopyxis*, in conjunction with *Blastomonas ursincola*. *Int J Syst Evol Microbiol* 52, 1485–1496.
- Yoon, J. H., Lee, M. H., Kang, S. J., Lee, S. Y. & Oh, T. K. (2006). *Sphingomonas dokdonensis* sp. nov., isolated from soil. *Int J Syst Evol Microbiol* 56, 2165–2169.