Transfer of *Sejongia antarctica*, *Sejongia jeonii* and *Sejongia marina* to the genus *Chryseobacterium* as *Chryseobacterium antarcticum* comb. nov., *Chryseobacterium jeonii* comb. nov. and *Chryseobacterium marinum* comb. nov.

Peter Kämpfer,¹ Nicole Lodders,¹ Mario Vaneechoutte²
and Georges Wauters³

Correspondence

Peter Kämpfer
peter.kaempfer@
umwelt.uni-giessen.de

¹Institut für Angewandte Mikrobiologie, Justus-Liebig-Universität Giessen, Germany

²Laboratory Bacteriology Research, Department of Clinical Chemistry, Microbiology and Immunology, University of Ghent, Ghent, Belgium

³Microbiology Unit, Faculty of Medicine, University of Louvain, Brussels, Belgium

The genus *Sejongia* was described in 2005, with the two species *Sejongia antarctica* and *Sejongia jeonii*, mainly on the basis of 16S rRNA gene sequence analysis. At that time, these organisms formed a quite separate branch in a 16S rRNA gene sequence-based tree, but, in subsequent studies, it became obvious that the species *S. antarctica* and *S. jeonii* and a third species, *Sejongia marina*, were most closely related (>95.0 % similarity) to some *Chryseobacterium* species (e.g. *Chryseobacterium hominis*, *C. formosense* and *C. haifense*). In addition, there is no evidence for clear phenotypic (i.e. chemotaxonomic) differences between these organisms that justifies their assignment to different genera. For these reasons, a proposal is made to transfer these species to the genus *Chryseobacterium* as *Chryseobacterium antarcticum* comb. nov. (type strain AT1013^T = JCM 12381^T = IMSNU 14040^T = KCTC 12225^T), *Chryseobacterium jeonii* comb. nov. (type strain AT1047^T = JCM 12382^T = IMSNU 14049^T = KCTC 12226^T) and *Chryseobacterium marinum* comb. nov. (type strain IMCC3228^T = KCCM 42689^T = NBRC 103143^T) on the basis of 16S rRNA gene sequence data and published phenotypic data.

The genus *Chryseobacterium* was proposed by Vandamme *et al.* (1994) in the course of the subdivision of the genus *Flavobacterium*, which at that time contained a rather heterogeneous group of yellow-pigmented, Gram-negative organisms. At that time, the new genus *Chryseobacterium* contained the six species *Chryseobacterium balustinum*, *C. gleum*, *C. indologenes*, *C. indoltheticum*, *C. meningosepticum* and *C. scophthalmum*. These species were similar in many respects. They share many classical phenotypic features: they all contain menaquinone 6 (MK-6) as their major respiratory quinone and they have similar fatty acid profiles, which are characterized by large amounts of iso-15:0, iso-17:1 ω 9c, iso-17:0 3-OH and summed feature 4 (iso-15:0 2-OH and/or 16:1 ω 7c). Within this group, *C. meningosepticum* revealed the most aberrant fatty acid profile and was subsequently transferred to the new genus *Elizabethkingia* (Kim *et al.*, 2005b). Between 1994 and 2008, the number of *Chryseobacterium* species increased significantly, to 37 species with validly published names (de Beer *et al.*, 2005; Gallego *et al.*, 2006; Herzog *et al.*, 2008;

Kämpfer *et al.*, 2003, 2009; Kim *et al.*, 2005a, 2008; Li *et al.*, 2003; Park *et al.*, 2006; Quan *et al.*, 2007; Shen *et al.*, 2005; Shimomura *et al.*, 2005; Tai *et al.*, 2006; Vaneechoutte *et al.*, 2007; Weon *et al.*, 2006; Young *et al.*, 2005). This increase in the number of species was accompanied by the proposal of two novel genera, *Kaistella* (Kim *et al.*, 2004) and *Sejongia* (Yi *et al.*, 2005), that share many features of *Chryseobacterium* species. The reclassification of the sole species of the genus *Kaistella* in the genus *Chryseobacterium* is proposed elsewhere (Kämpfer *et al.*, 2009).

Recently, several *Chryseobacterium* species, i.e. *Chryseobacterium haifense* (Hantsis-Zacharov & Halpern, 2007), *C. hominis* (Vaneechoutte *et al.*, 2007) and '*C. anthropi*' (Kämpfer *et al.*, 2009), amongst others, have been described with high 16S rRNA gene sequence similarities to the species of the genus *Sejongia*. In comparisons of the 16S rRNA gene sequences of type strains, *C. haifense* showed similarities of 95.3 % to *Sejongia antarctica*, 95.8 % to *Sejongia jeonii* and 95.0 % to *Sejongia marina*, *C. hominis* showed similarities of 95.1 % to *S. antarctica*, 96.5 % to *S.*

jeonii and 95.6 % to *S. marina* and '*C. anthropi*' showed similarities of 95.3 % to *S. antarctica*, 95.9 % to *S. jeonii* and 94.8 % to *S. marina*. 16S rRNA gene sequence similarities of the type strain of *C. formosense* (Young *et al.*, 2005) were 95.7 % to *S. antarctica*, 96.3 % to *S. jeonii* and 95.9 % to *S. marina*. Similarities were calculated using the ARB software package (version December 2007; Ludwig *et al.*, 2004) and the corresponding SILVA SSURef 95 database (release July 2008; Pruesse *et al.*, 2007).

The 16S rRNA gene sequences for all strains under comparison were analysed as described above. Trees were reconstructed using the maximum-likelihood method with fastDNAmI (Olsen *et al.*, 1994) and a 50 % conservation filter (only alignment columns in which the frequency of the most abundant nucleotide is equal to or more than 50 % are included in the calculation). Tree topology was further tested with no conservation filter and with 30 and 40 % conservation filters. No significant differences could be detected between these trees. For tree reconstruction, 90 sequences were included. For better clarity, only a subset of the sequences used for treeing are shown in Fig. 1.

On the basis of these comparisons, the type strains of the three *Sejongia* species fell clearly in the *Chryseobacterium* cluster at a depth similar to other *Chryseobacterium* species groups.

Although respiratory quinones have low resolution within this group, the presence of MK-6 supports affiliation of the *Sejongia* species to the genus *Chryseobacterium*, where all species investigated to date have MK-6 as the major quinone.

The fatty acid profiles of all *Sejongia* species were very similar to those of species of *Chryseobacterium*, as reported

already by Yi *et al.* (2005). All *Sejongia* species reveal iso-15:0, anteiso-15:1 and iso-17:1 ω 9c as the major fatty acids (Yi *et al.*, 2005; Lee *et al.*, 2007), which is essentially in agreement with the fatty acid patterns of *Chryseobacterium* species (de Beer *et al.*, 2005; Gallego *et al.*, 2006; Herzog *et al.*, 2008; Kämpfer *et al.*, 2003, 2009; Kim *et al.*, 2005a, 2008; Li *et al.*, 2003; Park *et al.*, 2006; Quan *et al.*, 2007; Shen *et al.*, 2005; Shimomura *et al.*, 2005; Tai *et al.*, 2006; Vanechoutte *et al.*, 2007; Weon *et al.*, 2006; Young *et al.*, 2005).

In conclusion, the species of *Sejongia* and *Chryseobacterium* show a strong congruence in phenotypic characters, in that they all produce MK-6 as the major menaquinone and the fatty acid profiles are very similar. Only a few differences have been reported with regard to cell morphology and certain physiological tests.

For these reasons, it is proposed to reclassify *Sejongia antarctica*, *Sejongia jeonii* and *Sejongia marina* in the genus *Chryseobacterium* as the new combinations *Chryseobacterium antarcticum* comb. nov., *Chryseobacterium jeonii* comb. nov. and *Chryseobacterium marinum* comb. nov.

Description of *Chryseobacterium antarcticum* comb. nov.

Chryseobacterium antarcticum (ant.arc'ti.cum. L. neut. adj. *antarcticum* southern, named after Antarctica, the geographical origin of the type strain).

Basonym: *Sejongia antarctica* Yi *et al.* 2005.

The description is that of Yi *et al.* (2005). The type strain is AT1013^T (=IMSNU 14040^T =KCTC 12225^T =JCM 12381^T).

Fig. 1. Phylogenetic analysis based on 16S rRNA gene sequences available from the EMBL database (accession numbers in parentheses). The phylogenetic tree was constructed using the ARB software package (version December 2007; Ludwig *et al.*, 2004) and the corresponding SILVA SSURef 95 database (version July 2008; Pruesse *et al.*, 2007). Tree building was performed with 90 sequences using the maximum-likelihood method with fastDNAmI (Olsen *et al.*, 1994) and 50 % conservation filter. For better clarity, only a subset of the sequences used for treeing is shown. Bar, 0.05 substitutions per nucleotide position.

Description of *Chryseobacterium jeonii* comb. nov.

Chryseobacterium jeonii (jeo'ni.i. N.L. gen. n. *jeonii* named in honour of the late Jae Gyu Jeon, who devoted his life to polar research).

Basonym: *Sejongia jeonii* Yi *et al.* 2005.

The description is that of Yi *et al.* (2005). The type strain is AT1047^T (=IMSNU 14049^T =KCTC 12226^T =JCM 12382^T).

Description of *Chryseobacterium marinum* comb. nov.

Chryseobacterium marinum (ma.ri'num. L. neut. adj. *marinum* of the sea, marine).

Basonym: *Sejongia marina* Lee *et al.* 2007.

The description is that of Lee *et al.* (2007). The type strain is IMCC3228^T (=KCCM 42689^T =NBRC 103143^T).

References

- de Beer, H., Hugo, C. J., Jooste, P. J., Willems, A., Vancanneyt, M., Coenye, T. & Vandamme, P. A. (2005). *Chryseobacterium vrystaatense* sp. nov., isolated from raw chicken in a chicken-processing plant. *Int J Syst Evol Microbiol* 55, 2149–2153.
- Gallego, V., Garcia, M. T. & Ventosa, A. (2006). *Chryseobacterium hispanicum* sp. nov., isolated from the drinking water distribution system of Sevilla, Spain. *Int J Syst Evol Microbiol* 56, 1589–1592.
- Hantsis-Zacharov, E. & Halpern, M. (2007). *Chryseobacterium haifense* sp. nov., a psychrotolerant bacterium isolated from raw milk. *Int J Syst Evol Microbiol* 57, 2344–2348.
- Herzog, P., Winkler, I., Wolking, D., Kämpfer, P. & Lipski, A. (2008). *Chryseobacterium ureilyticum* sp. nov., *Chryseobacterium gambrini* sp. nov., *Chryseobacterium pallidum* sp. nov. and *Chryseobacterium molle* sp. nov., isolated from beer-bottling plants. *Int J Syst Evol Microbiol* 58, 26–33.
- Kämpfer, P., Dreyer, U., Neef, A., Dott, W. & Busse, H.-J. (2003). *Chryseobacterium defluvii* sp. nov., isolated from wastewater. *Int J Syst Evol Microbiol* 53, 93–97.
- Kämpfer, P., Vanechoutte, M., Lodders, N., De Baere, T., Avesani, V., Janssens, M., Busse, H.-J. & Wauters, G. (2009). Description of *Chryseobacterium anthropi* sp. nov., to accommodate clinical isolates biochemically similar to *Kaistella koreensis* and *Chryseobacterium haifense*, proposal to reclassify *Kaistella koreensis* as *Chryseobacterium koreense* comb. nov. and emended description of the genus *Chryseobacterium*. *Int J Syst Evol Microbiol* 59 (in press). doi:10.1099/ijs.0.008250-0.
- Kim, M. K., Im, W. T., Shin, Y. K., Lim, J. H., Kim, S. H., Lee, B. C., Park, M. Y., Lee, K. Y. & Lee, S. T. (2004). *Kaistella koreensis* gen. nov., sp. nov., a novel member of the *Chryseobacterium*–*Bergeyella*–*Riemerella* branch. *Int J Syst Evol Microbiol* 54, 2319–2324.
- Kim, K. K., Bae, H. S., Schumann, P. & Lee, S. T. (2005a). *Chryseobacterium daecheongense* sp. nov., isolated from freshwater lake sediment. *Int J Syst Evol Microbiol* 55, 133–138.
- Kim, K. K., Kim, M. K., Lim, J. H., Park, H. Y. & Lee, S. T. (2005b). Transfer of *Chryseobacterium meningosepticum* and *Chryseobacterium miricola* to *Elizabethkingia* gen. nov. as *Elizabethkingia meningoseptica* comb. nov. and *Elizabethkingia miricola* comb. nov. *Int J Syst Evol Microbiol* 55, 1287–1293.
- Kim, K. K., Lee, K. C., Oh, H.-M. & Lee, J.-S. (2008). *Chryseobacterium aquaticum* sp. nov., isolated from a water reservoir. *Int J Syst Evol Microbiol* 58, 533–537.
- Lee, K., Lee, H. K., Choi, T.-H. & Cho, J.-C. (2007). *Sejongia marina* sp. nov., isolated from Antarctic seawater. *Int J Syst Evol Microbiol* 57, 2917–2921.
- Li, Y., Kawamura, Y., Fujiwara, N., Naka, T., Liu, H., Huang, X., Kobayashi, K. & Ezaki, T. (2003). *Chryseobacterium miricola* sp. nov., a novel species isolated from condensation water of space station Mir. *Syst Appl Microbiol* 26, 523–528.
- Ludwig, W., Strunk, O., Westram, R., Richter, L., Meier, H., Yadukumar, Buchner, A., Lai, T., Steppi, S. & other authors (2004). ARB: a software environment for sequence data. *Nucleic Acids Res* 32, 1363–1371.
- Olsen, G. J., Matsuda, H., Hagström, R. & Overbeek, R. (1994). fastDNAMl: a tool for construction of phylogenetic trees of DNA sequences using maximum likelihood. *Comput Appl Biosci* 10, 41–48.
- Park, M. S., Jung, S. R., Lee, K. H., Lee, M. S., Do, J. O., Kim, S. B. & Bae, K. S. (2006). *Chryseobacterium soldanellicola* sp. nov. and *Chryseobacterium taeanense* sp. nov., isolated from roots of sand-dune plants. *Int J Syst Evol Microbiol* 56, 433–438.
- Pruesse, E., Quast, C., Knittel, K., Fuchs, B. M., Ludwig, W., Peplies, J. & Glöckner, F. O. (2007). SILVA: a comprehensive online resource for quality checked and aligned ribosomal RNA sequence data compatible with ARB. *Nucleic Acids Res* 35, 7188–7196.
- Quan, Z.-X., Kim, K. K., Kim, M.-K., Jin, L. & Lee, S. T. (2007). *Chryseobacterium caeni* sp. nov., isolated from bioreactor sludge. *Int J Syst Evol Microbiol* 57, 141–145.
- Shen, F. T., Kämpfer, P., Young, C. C., Lai, W. A. & Arun, A. B. (2005). *Chryseobacterium taichungense* sp. nov., isolated from contaminated soil. *Int J Syst Evol Microbiol* 55, 1301–1304.
- Shimomura, K., Kaji, S. & Hiraishi, A. (2005). *Chryseobacterium shigense* sp. nov., a yellow-pigmented, aerobic bacterium isolated from a lactic acid beverage. *Int J Syst Evol Microbiol* 55, 1903–1906.
- Tai, C. J., Kuo, H. P., Lee, F. L., Chen, H. K., Yokota, A. & Lo, C. C. (2006). *Chryseobacterium taiwanense* sp. nov., isolated from soil in Taiwan. *Int J Syst Evol Microbiol* 56, 1771–1776.
- Vandamme, P., Bernardet, J.-F., Segers, P., Kersters, K. & Holmes, B. (1994). New perspectives in the classification of the flavobacteria: description of *Chryseobacterium* gen. nov., *Bergeyella* gen. nov., and *Empedobacter* nom. rev. *Int J Syst Bacteriol* 44, 827–831.
- Vanechoutte, M., Kämpfer, P., De Baere, T., Avesani, V., Janssens, M. & Wauters, G. (2007). Description of *Chryseobacterium hominis* sp. nov. to accommodate clinical isolates biochemically similar to CDC groups II-h and II-c. *Int J Syst Evol Microbiol* 57, 2623–2628.
- Weon, H. Y., Kim, B. Y., Yoo, S. H., Kwon, S. W., Cho, Y. H., Go, S. J. & Stackebrandt, E. (2006). *Chryseobacterium wanjuense* sp. nov., isolated from greenhouse soil in Korea. *Int J Syst Evol Microbiol* 56, 1501–1504.
- Yi, H., Yoon, H. I. & Chun, J. (2005). *Sejongia antarctica* gen. nov., sp. nov. and *Sejongia jeonii* sp. nov., isolated from the Antarctic. *Int J Syst Evol Microbiol* 55, 409–416.
- Young, C. C., Kämpfer, P., Shen, F. T., Lai, W. A. & Arun, A. B. (2005). *Chryseobacterium formosense* sp. nov., isolated from the rhizosphere of *Lactuca sativa* L. (garden lettuce). *Int J Syst Evol Microbiol* 55, 423–426.