Williamsia faeni sp. nov., an actinomycete isolated from a hay meadow

Amanda L. Jones,[†] Gail D. Payne[‡] and Michael Goodfellow

School of Biology, University of Newcastle, Newcastle upon Tyne NE1 7RU, UK

Correspondence

Amanda L. Jones

Amanda.L.Jones@northumbria.ac.uk

The taxonomic status of an actinomycete isolated from soil collected from a hay meadow was determined using a polyphasic approach. The strain, designated N1350^T, had morphological and chemotaxonomic properties consistent with its classification in the genus *Williamsia* and formed a distinct phyletic line within the clade comprising the type strains of species of the genus *Williamsia* in the 16S rRNA gene tree. Strain N1350^T shared highest 16S rRNA gene sequence similarities with *Williamsia marianensis* MT8^T (98.1 %) and *Williamsia muralis* MA140-96^T (98.3 %). However, strain N1350^T was readily distinguished from the type strains of *Williamsia* species using a combination of phenotypic properties. On the basis of these data, strain N1350^T is considered to represent a novel species of the genus *Williamsia*. The name proposed for this taxon is *Williamsia faeni* sp. nov., with the type strain N1350^T (=DSM 45372^T =NCIMB 14575^T =NRRL B-24794^T).

The genus *Williamsia* (Kämpfer *et al.*, 1999) has been classified in the family *Nocardiaceae*, together with the genera *Gordonia*, *Millisia*, *Nocardia*, *Rhodococcus* and *Skermania* (Zhi *et al.*, 2009). At the time of writing, the genus encompasses five described species: *Williamsia muralis* (Kämpfer *et al.*, 1999), the type species, isolated from indoor building material of a children's day centre in Finland, *Williamsia deligens* (Yassin & Hupfer, 2006) from human blood, *Williamsia marianensis* (Pathom-Aree *et al.*, 2006) from sediment taken from the Mariana Trench in the Pacific Ocean, *Williamsia maris* (Stach *et al.*, 2004) from sediment collected from the Sea of Japan and *Williamsia serinedens* (Yassin *et al.*, 2007) from an oil-contaminated soil. The type strains of these species form a clade within the evolutionary radiation occupied by the suborder *Corynebacterineae* (Stackebrandt *et al.*, 1997; Zhi *et al.*, 2009). The genus *Williamsia* can also be distinguished from the other mycolic acid-containing genera using a combination of chemotaxonomic and morphological properties (Soddell *et al.*, 2006; Adachi *et al.*, 2007). Similarly, species of the genus *Williamsia* can be distinguished from each other by using a range of phenotypic properties (Yassin *et al.*, 2007). The aim of the present study was to determine the taxonomic position of an actinomycete, strain N1350^T, that had been recovered from a hay meadow soil and provisionally assigned to the genus *Williamsia*.

Strain N1350^T was isolated from a soil suspension inoculum on Gauze's medium 2 supplemented with ($\mu\text{g ml}^{-1}$) cycloheximide (50), nalidixic acid (10), novobiocin (10) and nystatin (50) after incubation at 30 °C for 21 days (Tan *et al.*, 2006). The soil sample had been collected from underneath the surface of Palace Leas meadow hay plot 6 (Atalan *et al.*, 2000) at Cockle Park Experimental Farm, Northumberland, UK (national grid reference NZ 200913). Strain N1350^T was maintained on glucose-yeast extract agar (GYEA; Gordon & Mihm, 1962) at room temperature and as glycerol suspensions (20 %, v/v) at –20 °C. Biomass for the chemotaxonomic and 16S rRNA gene sequence analyses was grown in shake flasks of GYE broth for 5 days at 28 °C, checked for purity and harvested by centrifugation. Cells for the chemosystematic studies were washed twice in distilled water and freeze-dried; cells for the 16S rRNA study were washed in NaCl/EDTA buffer (0.1 M NaCl, 0.1 M EDTA, pH 8.0) and stored at –20 °C until required.

The phylogenetic position of strain N1350^T was determined in a 16S rRNA gene sequence analysis. Isolation of chromosomal DNA, PCR amplification and direct sequencing of the purified products were carried out as described by Kim *et al.* (1998). The almost-complete 16S rRNA gene sequence (1441 nt) was aligned manually with corresponding sequences of representatives of genera classified in the suborder *Corynebacterineae* retrieved from the DDBJ/EMBL/GenBank databases using the pairwise alignment option and 16S rRNA secondary structural information held in the PHYDIT program (available at <http://plaza.snu.ac.kr/~jchun/phydit/>). Phylogenetic trees were inferred using the least-squares (Fitch & Margoliash, 1967), neighbour-joining (Saitou & Nei, 1987), maximum-parsimony (Kluge & Farris,

[†]Present address: School of Applied Sciences, Ellison Building, Northumbria University, Newcastle upon Tyne NE1 8ST, UK.

[‡]Present address: Agencourt Bioscience, 500 Cummings Center, Beverly, MA 01915, USA.

The GenBank/EMBL/DDBJ accession number for the 16S rRNA gene sequence of strain N1350^T is DQ157929.

1969) and maximum-likelihood (Felsenstein, 1981) tree-making algorithms from the PHYLIP suite of programs (Felsenstein, 1993) and evolution distance matrices were prepared according to Jukes & Cantor (1969). The resulting unrooted tree topologies were evaluated in a bootstrap analysis (Felsenstein, 1985) based on 1000 resamplings of the neighbour-joining dataset using the CONSENSE and SEQBOOT options from the PHYLIP package.

The 16S rRNA gene phylogenetic tree is shown in Fig. 1. Strain N1350^T fell within the 16S rRNA gene clade for the genus *Williamsia*. This association was supported by all of the tree-making algorithms and by a 100 % bootstrap value in the neighbour-joining analysis. Strain N1350^T shared highest 16S rRNA gene sequence similarities with *W. muralis* MA140-96^T (98.3 %, corresponding to 24 nucleotide differences across 1416 locations) and *W. marianensis* MT8^T (98.1 %). DNA–DNA hybridization studies were not performed between these strains as *W. marianensis* MT8^T and *W. muralis* MA140-96^T, which formed a subclade with strain N1350^T, share a higher 16S rRNA sequence similarity (99.5 %) with each other but a DNA–DNA relatedness value of only 11 % (Pathom-Aree *et al.*, 2006), which is well below the 70 % cut-off point recommended for the delineation of bacterial species (Wayne *et al.*, 1987).

Strain N1350^T was examined for key chemotaxonomic markers that are characteristic for the genus *Williamsia*. Standard procedures were used to determine the diagnostic isomers of diaminopimelic acid (Staneck & Roberts, 1974), fatty acids (Sutcliffe, 2000), isoprenoid quinones (Collins, 1994), muramic acid type (Uchida *et al.*, 1999), mycolic acids (Minnikin *et al.*, 1975), polar lipids (Minnikin *et al.*, 1984) and whole-organism sugars (Hasegawa *et al.*, 1983).

Strain N1350^T contained *meso*-diaminopimelic acid, arabinose and galactose in whole-organism hydrolysates (wall chemotype IV *sensu* Lechevalier & Lechevalier, 1970), *N*-glycolyl muramic acid and dihydrogenated menaquinones with nine isoprene units as the sole isoprenologue. The fatty acid profile included major proportions of straight-chain saturated, unsaturated and tuberculostearic acids [fatty acid type 1b *sensu* Kroppenstedt, 1985; hexadecanoic (C_{16:0}, 21 % of total), monosaturated octadecanoic (C_{18:1}, 15 %), tridecanoic (C_{13:0}, 11 %), tuberculostearic (10-methyl C_{18:0}, 8 %) and octadecanoic (C_{18:0}, 7 %) acids] and minor proportions of tetradecanoic (C_{14:0}), pentadecanoic (C_{15:0}), iso-hexadecanoic (iso-C_{16:0}) and eicosanoic (C_{20:0}) acids. Phosphatidylethanolamine, phosphatidylglycerol, diphosphatidylglycerol and phosphatidylinositol were the major polar lipids. Strain N1350^T contained mycolic acids that co-migrated with those from *W. muralis* DSM 44343^T. This chemotaxonomic profile is consistent with the classification of strain N1350^T in the genus *Williamsia* (Goodfellow & Maldonado, 2006; Yassin & Hupfer, 2006).

Strain N1350^T was examined for a range of phenotypic properties using a range of media and methods known to yield data of value for the classification and identification of mycolic-acid-containing actinomycetes (Jones *et al.*, 2008). Strain N1350^T was aerobic, Gram-staining-positive, non-acid–alcohol-fast, asporogenous and catalase-positive and used a diverse range of compounds as sole carbon sources. These properties were in line with the classification of strain N1350^T in the genus *Williamsia* (Kämpfer *et al.*, 1999; Yassin *et al.*, 2007). Strain N1350^T could be readily distinguished from the type strains of *Williamsia* species using a combination of phenotypic properties (Table 1). Strain N1350^T could also be distinguished from *W. deligens*

Fig. 1. Maximum-likelihood tree (Felsenstein, 1981) based on the nearly complete 16S rRNA gene sequence of strain N1350^T, showing the strain's relationships within the *Williamsia* clade. Asterisks indicate that the corresponding nodes were also recovered in trees generated with the least-squares (Fitch & Margoliash, 1967), maximum-parsimony (Kluge & Farris, 1969) and neighbour-joining (Saitou & Nei, 1987) algorithms. F, P and N indicate that the corresponding nodes were also recovered in trees generated with the least-squares, maximum-parsimony or neighbour-joining algorithms, respectively. Bootstrap values (>50 %) based on a neighbour-joining analysis of 1000 resampled datasets are shown at branch nodes. Bar, 50 changes.

Table 1. Phenotypic properties that distinguish strain N1350^T from the type strains of *Williamsia* species

Strains: 1, *Williamsia faeni* sp. nov. N1350^T; 2, *W. marianensis* DSM 44944^T (data from Pathom-Aree *et al.*, 2006); 3, *W. muralis* DSM 44343^T; 4, *W. deligens* DSM 44902^T; 5, *W. maris* DSM 44693^T; 6, *W. serinedens* DSM 45037^T. Data were obtained in this study unless indicated. +, Positive; −, negative; ND, not determined.

Characteristic	1	2	3	4	5	6
Growth at (°C):						
4	+	+	−	−	−	−
10	+	+	+	−	+	+
37	−	−	+	+	+	−
45	−	−	+	−	−	−
Aesculin hydrolysis	+	−	−	−	ND	−
Growth with (1% (w/v):						
Adonitol	+	−	+	−	−	+
(−)-L-Arabinose	+	+	−	−	−	+
(+)-Cellobiose	+	−	−	−	−	−
(−)-D-Galactose	+	−	−	−	−	+
myo-Inositol	+	−	−	−	−	−
(+)-Maltose	+	−	−	+	−	+
(−)-D-Mannitol	+	+	+	+	−	+
(+)-Melibiose	+	−	+	−	−	+
(+)-D-Raffinose	+	−	−	−	−	−
α-L-Rhamnose	+	+	+	−	+	−
(−)-D-Sorbitol	+	+	+	+	−	+
(+)-Sucrose	+	+	+	+	−	+
(+)-Trehalose	+	+	−	+	+	+
(+)-D-Xylose	+	−	−	+	+	+
Growth with (0.1%, w/v):						
m-Hydroxybenzoic acid	−	−	+	−	−	+
p-Hydroxybenzoic acid	+	−	−	−	−	−
Growth with (1%, v/v):						
1,2-Propanediol	+	−	−	−	−	+

DSM 44902^T and *W. serinedens* DSM 45037^T by its ability to degrade L-tyrosine (Yassin & Hupfer, 2006; Yassin *et al.*, 2007) and from *W. marianensis* DSM 44944^T by its ability to degrade tributyrin but not hypoxanthine (Pathom-Aree *et al.*, 2006).

It can be concluded from the genotypic and phenotypic data that strain N1350^T can be readily distinguished from described *Williamsia* species and, hence, should be classified as a representative of a novel species in the genus *Williamsia*. The name proposed for this taxon is *Williamsia faeni* sp. nov.

Description of *Williamsia faeni* sp. nov.

Williamsia faeni (fae'ni. L. n. *faenum* hay; L. gen. n. *faeni* of hay, referring to the isolation of the type strain from a hay meadow).

Forms coccoid elements. Produces irregular, convex, matt yellow–pink-pigmented colonies on GYEA after incubation for 5 days at 28 °C. Grows at 10–30 °C, but not at 37 °C. Exhibits chemotaxonomic markers characteristic of the

genus *Williamsia*. Hydrolyses allantoin and urea, but not arbutin. Degrades DNA, RNA, starch and uric acid, but not adenine, chitin, elastin, xanthine or xylan. Uses (−)-D-amylgdalin, (−)-D-arabinose, (+)-D-arabitol, arbutin, (−)-D-fructose, (−)-D-fucose, (−)-D-glucose, inulin, (+)-lactose, (+)-D-mannose, (+)-melibiose, methyl α-D-glucoside, (−)-D-ribose, (+)-turanose (all at 1%, w/v), butan-1,3-diol, butan-1,4-diol, butan-1-ol, butan-2,3-diol, ethanol, propan-1-ol, propan-2-ol (all at 1%, v/v), 3-methyl-1-butanol (isoamyl alcohol), benzoic acid, fumaric acid, glycerol, glycogen, (+)-L-lactic acid, L-malic acid, oleic acid, propanoic acid, pyruvic acid, sodium acetate, sodium n-butyrate, (+)-L-tartaric acid, valeric acid and xylitol (all at 0.1%, w/v) as sole carbon sources for energy and growth, but not dulcitol, salicin (all at 1%, w/v), adipic acid, citric acid, glutaric acid, malonic acid, D-mandelic acid, oxalic acid, sebamic acid, suberic acid or succinic acid (all at 0.1%, w/v). Uses acetamide, L-alanine, L-α-aminobutyric acid, L-arginine, L-gelatin, D-gluconic acid, L-glycine, histidine, L-leucine, DL-methionine, monoethanolamine, DL-norleucine, L-norvaline, DL-phenylalanine, L-proline, serine, uric acid, urea and L-valine as sole carbon and nitrogen sources, but not L-cysteine, L-glutamic acid, L-isoleucine or L-ornithine (all at 0.1%, w/v). Additional phenotypic properties are indicated in the text and in Table 1. The fatty acid profile includes major amounts of hexadecanoic (C_{16:0}), octadecenoic (C_{18:1}), tridecanoic (C_{13:0}), tuberculostearic (10-methyl C_{18:0}) and octadecanoic (C_{18:0}) acids and minor amounts of tetradecanoic (C_{14:0}), pentadecanoic (C_{15:0}), iso-hexadecanoic (iso-C_{16:0}) and eicosanoic (C_{20:0}) acids.

The type strain, N1350^T (=DSM 45372^T =NCIB 14575^T =NRRL B24794^T), was isolated from a hay meadow plot at Cockle Park Experimental Farm, Northumberland, UK.

Acknowledgements

The authors are indebted to Dr Iain Sutcliffe (University of Northumbria) for help with the fatty acid analysis of strain N1350^T.

References

- Adachi, K., Katsuta, A., Matsuda, S., Peng, X., Misawa, N., Shizuri, Y., Kroppenstedt, R. M., Yokota, A. & Kasai, H. (2007). *Smaragdicoccus niigatensis* gen. nov., sp. nov., a novel member of the suborder *Corynebacterineae*. *Int J Syst Evol Microbiol* **57**, 297–301.
- Atalan, E., Manfio, G. P., Ward, A. C., Kroppenstedt, R. M. & Goodfellow, M. (2000). Biosystematic studies on novel streptomycetes from soil. *Antonie van Leeuwenhoek* **77**, 337–353.
- Collins, M. D. (1994). Isoprenoid quinones. In *Chemical Methods in Prokaryotic Systematics*, pp. 265–309. Edited by M. Goodfellow & A. G. O'Donnell. Chichester: Wiley.
- Felsenstein, J. (1981). Evolutionary trees from DNA sequences: a maximum likelihood approach. *J Mol Evol* **17**, 368–376.
- Felsenstein, J. (1985). Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* **39**, 783–791.

- Felsenstein, J. (1993).** PHYLIP (phylogeny inference package), version 3.5c. Distributed by the author. Department of Genome Sciences, University of Washington, Seattle, USA.
- Fitch, W. M. & Margoliash, E. (1967).** Construction of phylogenetic trees. *Science* **155**, 279–284.
- Goodfellow, M. & Maldonado, L. A. (2006).** The families Dietziaceae, Gordoniaceae, Nocardiaceae and Tsukamurellaceae. In *The Prokaryotes: a Handbook on the Biology of Bacteria*, 3rd edn, vol. 3, pp. 843–888. Edited by M. Dworkin, S. Falkow, E. Rosenberg, K. H. Schleifer & E. Stackebrandt. New York: Springer.
- Gordon, R. E. & Mihm, J. M. (1962).** Identification of *Nocardia caviae* (Erikson) nov. comb. *Ann N Y Acad Sci* **98**, 628–636.
- Hasegawa, T., Takizawa, M. & Tanida, S. (1983).** A rapid analysis for chemical grouping of aerobic actinomycetes. *J Gen Appl Microbiol* **29**, 319–322.
- Jones, A. L., Koerner, R. J., Natarajan, S., Perry, J. D. & Goodfellow, M. (2008).** *Dietzia papillomatosis* sp. nov., a novel actinomycete isolated from the skin of an immunocompetent patient with confluent and reticulated papillomatosis. *Int J Syst Evol Microbiol* **58**, 68–72.
- Jukes, T. H. & Cantor, C. R. (1969).** Evolution of protein molecules. In *Mammalian Protein Metabolism*, vol. 3, pp. 21–132. Edited by H. N. Munro. New York: Academic Press.
- Kämpfer, P., Andersson, M. A., Rainey, F. A., Kroppenstedt, R. M. & Salkinoja-Salonen, M. (1999).** *Williamsia muralis* gen. nov., sp. nov., isolated from the indoor environment of a children's day care centre. *Int J Syst Bacteriol* **49**, 681–687.
- Kim, S. B., Falconer, C., Williams, E. & Goodfellow, M. (1998).** *Streptomyces thermocarboxydovorans* sp. nov. and *Streptomyces thermocarboxydus* sp. nov., two moderately thermophilic carboxydophilic species from soil. *Int J Syst Bacteriol* **48**, 59–68.
- Kluge, A. G. & Farris, J. S. (1969).** Quantitative phyletics and the evolution of anurans. *Syst Zool* **18**, 1–32.
- Kroppenstedt, R. M. (1985).** Fatty acid and menaquinone analysis of actinomycetes and related organisms. In *Chemical Methods in Bacterial Systematics* (Society for Applied Bacteriology Technical Series vol. 20), pp. 173–199. Edited by M. Goodfellow & D. E. Minnikin. New York: Academic Press.
- Lechevalier, M. P. & Lechevalier, H. A. (1970).** Chemical composition as a criterion in the classification of aerobic actinomycetes. *Int J Syst Bacteriol* **20**, 435–443.
- Minnikin, D. E., Alshamaony, L. & Goodfellow, M. (1975).** Differentiation of *Mycobacterium*, *Nocardia*, and related taxa by thin-layer chromatographic analysis of whole-organism methanolysates. *J Gen Microbiol* **88**, 200–204.
- Minnikin, D. E., O'Donnell, A. G., Goodfellow, M., Alderson, G., Athalye, M., Schaal, A. & Parlett, J. H. (1984).** An integrated procedure for the extraction of bacterial isoprenoid quinones and polar lipids. *J Microbiol Methods* **2**, 233–241.
- Pathom-Aree, W., Nogi, Y., Sutcliffe, I. C., Ward, A. C., Horikoshi, K., Bull, A. T. & Goodfellow, M. (2006).** *Williamsia marianensis* sp. nov., a novel actinomycete isolated from the Mariana Trench. *Int J Syst Evol Microbiol* **56**, 1123–1126.
- Saitou, N. & Nei, M. (1987).** The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol* **4**, 406–425.
- Soddell, J. A., Stainsby, F. M., Eales, K. L., Kroppenstedt, R. M., Seviour, R. J. & Goodfellow, M. (2006).** *Millisia brevis* gen. nov., sp. nov., an actinomycete isolated from activated sludge foam. *Int J Syst Evol Microbiol* **56**, 739–744.
- Stach, J. E., Maldonado, L. A., Ward, A. C., Bull, A. T. & Goodfellow, M. (2004).** *Williamsia maris* sp. nov., a novel actinomycete isolated from the Sea of Japan. *Int J Syst Evol Microbiol* **54**, 191–194.
- Stackebrandt, E., Rainey, F. A. & Ward-Rainey, N. L. (1997).** Proposal for a new hierarchic classification system, *Actinobacteria* classis nov. *Int J Syst Bacteriol* **47**, 479–491.
- Staneck, J. L. & Roberts, G. D. (1974).** Simplified approach to identification of aerobic actinomycetes by thin-layer chromatography. *Appl Microbiol* **28**, 226–231.
- Sutcliffe, I. C. (2000).** Characterisation of a lipomannan lipoglycan from the mycolic acid containing actinomycete *Dietzia maris*. *Antonie van Leeuwenhoek* **78**, 195–201.
- Tan, G. Y., Ward, A. C. & Goodfellow, M. (2006).** Exploration of *Amycolatopsis* diversity in soil using genus-specific primers and novel selective media. *Syst Appl Microbiol* **29**, 557–569.
- Uchida, K., Kudo, T., Suzuki, K. & Nakase, T. (1999).** A new rapid method of glycolate test by diethyl ether extraction, which is applicable to a small amount of bacterial cells of less than one milligram. *J Gen Appl Microbiol* **45**, 49–56.
- Wayne, L. G., Brenner, D. J., Colwell, R. R., Grimont, P. A. D., Kandler, O., Krichevsky, M. I., Moore, L. H., Moore, W. E. C., Murray, R. G. E. & other authors (1987).** International Committee on Systematic Bacteriology. Report of the ad hoc committee on reconciliation of approaches to bacterial systematics. *Int J Syst Bacteriol* **37**, 463–464.
- Yassin, A. F. & Hupfer, H. (2006).** *Williamsia deligens* sp. nov., isolated from human blood. *Int J Syst Evol Microbiol* **56**, 193–197.
- Yassin, A. F., Young, C. C., Lai, W. A., Hupfer, H., Arun, A. B., Shen, F. T., Rekha, P. D. & Ho, M. J. (2007).** *Williamsia serinedens* sp. nov., isolated from an oil-contaminated soil. *Int J Syst Evol Microbiol* **57**, 558–561.
- Zhi, X. Y., Li, W. J. & Stackebrandt, E. (2009).** An update of the structure and 16S rRNA gene sequence-based definition of higher ranks of the class *Actinobacteria*, with the proposal of two new suborders and four new families and emended descriptions of the existing higher taxa. *Int J Syst Evol Microbiol* **59**, 589–608.