Dietzia timorensis sp. nov., isolated from soil

Hideki Yamamura,^{1,2} Puspita Lisdiyanti,³ Roni Ridwan,³
Shanti Ratnakomala,³ Rasti Sarawati,⁴ Yulin Lestari,⁵ Evi Triana,⁶
Gina Kartina,³ Yantyati Widyastuti³ and Katsuhiko Ando²

Correspondence

Hideki Yamamura
hyamamura@yamanashi.ac.jp

¹Division of Applied Biological Sciences, Interdisciplinary Graduate School of Medicine and Engineering, University of Yamanashi, Takeda-4, Kofu 400, Japan

²NITE Biological Resource Center, Department of Biotechnology, National Institute of Technology and Evaluation, Japan

³Research Center for Biotechnology, Indonesian Institute of Sciences, Jl. Raya Bogor Km. 46, Cibinong 16911, Indonesia

⁴Soil Research Institute, Jl. Ir. H. Juanda 98, Bogor 16002, Indonesia

⁵Department of Biology, Faculty of Mathematics and Natural Sciences, Bogor Agricultural University, Jalan Raya Pajajaran, Bogor 16144, Indonesia

⁶Research Center for Biology, Indonesian Institute of Sciences, Jl. Raya Bogor Km. 46, Cibinong 16911, Indonesia

An actinomycete strain, ID05-A0528^T, was isolated using the SDS-yeast extract pre-treatment method from soil under mahogany (*Swietenia mahoganii*) trees in West Timor, Indonesia, and was examined by using a polyphasic taxonomic approach. Chemotaxonomic and phylogenetic characterizations demonstrated that the novel strain belongs to the genus *Dietzia*. 16S rRNA gene sequencing studies showed that the strain was related to *Dietzia cinnamea* (97.2%). Results of phenotypic and phylogenetic analyses determined that strain ID05-A0528^T is different from the known species of the genus *Dietzia*. It is proposed that the isolate should be classified as a representative of a novel species of the genus *Dietzia*, with the name *Dietzia timorensis* sp. nov. The type strain is ID05-A0528^T (=BTCC B-560^T =NBRC 104184^T).

The genus *Dietzia* is a member of the suborder *Corynebacterineae* (Stackebrandt *et al.*, 1997) and encompasses eight species at the time of writing, including *Dietzia papillomatosus*, *Dietzia schimae* and *Dietzia cercidiphylli* (Jones *et al.*, 2008; Li *et al.*, 2008). Known species of the genus *Dietzia* were originally isolated from several sources, including clinical materials, such as an alkaline soda lake, a perianal swab, a drain pool of a fish-egg processing plant, soil, the skin of an immunocompetent patient, and plant tissue (Duckworth *et al.*, 1998; Yumoto *et al.*, 2002; Yassin *et al.*, 2006; Mayilraj *et al.*, 2006; Jones *et al.*, 2008; Li *et al.*, 2008). Some strains identified as representing species of the genus *Dietzia* show degradation of hydrocarbons, including *n*-alkanes (Rainey *et al.*, 1995; Chaillan *et al.*, 2004; Yumoto *et al.*, 2002). Additionally, Takeishi *et al.* (2006) reported xylanolytic strains of the genus *Dietzia* isolated from the hindgut and faeces of *Trypoxylus dichotomus* larvae. Hence, the discovery of additional species of this genus will help in understanding their ecological roles and provide

bioresources for industrial applications, including bioremediation.

Strain ID05-A0528^T was isolated from a soil sample collected under mahogany trees in West Timor. The SDS-yeast extract pre-treatment method (Hayakawa & Nonomura, 1989) and humic acid-vitamin agar (Hayakawa & Nonomura, 1987) containing nalidixic acid (20 mg l⁻¹) were used in the isolation. The pre-treatment method was used to enhance the spore germination of actinomycetes and to decrease the number of non-filamentous bacteria on the isolation plates. The aim of the present study was to determine the taxonomic position of isolate ID05-A0528^T using a polyphasic approach.

The colonial properties of strain ID05-A0528^T were recorded from a modified Bennett's agar plate (Jones, 1949) that had been incubated for 14 days at 28 °C. Gram-staining was examined by using Hucker's method (Gerhardt, 1981). Motility was examined in hanging drops by light microscopy using culture grown on Bennett's agar plates. Morphology of the cells was observed using light microscopy. Tests for aesculin and arbutin hydrolysis (Williams *et al.*, 1983), nitrate reduction (Gordon & Mihm,

The GenBank/EMBL/DDBJ accession number for the 16S rRNA gene sequence of strain ID05-A0528^T is AB377289.

1962) and urea hydrolysis (Gordon *et al.*, 1974) were performed by using established procedures. Hydrolysis of complex substrates was examined by using well-established procedures (Gordon *et al.*, 1974; Isik *et al.*, 1999). The temperature for growth was determined at 5, 10, 15, 28, 37 and 45 °C on modified Bennett's agar plates (Jones, 1949). API 50 CH test kits (bioMérieux) were used to investigate several physiological and biochemical characteristics, according to the manufacturer's instructions. Diaminopimelic acid isomers and sugars in whole-cell hydrolysates were analysed based on the methods established by Hasegawa *et al.* (1983) and Schaal (1985), respectively. Standard procedures were also used for extraction and analysis of mycolic acids (Schaal, 1985), fatty acids (Tamura *et al.*, 1994), and isoprenoid quinones and polar lipids (Minnikin *et al.*, 1984), and compared with the appropriate controls. The acyl type of muramic acid was determined by the colorimetric method (Uchida & Aida, 1977). Chromosomal DNA from strain ID05-A0528^T was isolated and purified by the method of Saito & Miura (1963) with a minor modification (Hatano *et al.*, 2003). The G + C content (mol%) of DNA from the isolate was determined by HPLC, as described by Tamura *et al.* (1994).

The 16S rRNA gene from strain ID05-A0528^T was amplified by PCR following the procedures described by Tamura & Hatano (2001) and directly sequenced using an ABI Prism BigDye Terminator cycle sequencing kit (PE Applied Biosystems) and an automatic DNA sequencer (model 3130 Genetic Analyzer; PE Applied Biosystems). The 16S rRNA gene sequence obtained was aligned with reference sequences of members of the genus *Dietzia* available from EMBL/GenBank/DDBJ by using the CLUSTAL_X program (Thompson *et al.*, 1997). A phylogenetic tree was inferred using neighbour-joining tree algorithms (Saitou & Nei, 1987). Topography of the reconstructed tree was evaluated by bootstrap analysis with 1000 replicates (Felsenstein, 1985).

The chemotaxonomic characteristics of isolate ID05-A0528^T were consistent with its assignment to the genus *Dietzia*. The whole-cell hydrolysate contained *meso*-diaminopimelic acid, arabinose and galactose (wall chemotype IV *sensu* Lechevalier & Lechevalier, 1970). The major

menaquinone was MK-8 (H₂), in accordance with recognized species of the genus *Dietzia*. Strain ID05-A0528^T contained phosphatidylglycerol and trace amounts of phosphatidylinositol. Muramic acid residues of the peptidoglycan were *N*-acetylated. In addition, TLC revealed that the strain contained mycolic acids, which are equivalent in mobility ($R_f=0.45$) to those of *Dietzia natronolimnaea* NBRC105040^T; it was thus inferred that the mycolic acids in strain ID05-A0528^T contained 34–38 carbon atoms (Duckworth *et al.*, 1998). C_{16:0} (48%), C_{18:1ω9c} (39%) and 10-methyl C_{18:0} (8%) were present as major cellular fatty acids. Colonies were moderately orange-yellow.

The almost complete 16S rRNA gene sequence (1455 nt) of isolate ID05-A0528^T was compared with sequences from recognized species of the genus *Dietzia*. Results of phylogenetic tree reconstruction based on the neighbour-joining method showed that the novel strain formed a monophyletic clade with known species of this genus (Fig. 1). 16S rRNA gene sequence similarities to known species of the genus *Dietzia* were 95.5–97.2%. The closest phylogenetic relative was *Dietzia cinnamea*, sharing 97.2% sequence similarity with strain ID05-A0528^T. DNA–DNA hybridization between strain ID05-A0528^T and the nearest phylogenetic neighbours was not attempted, since Stackebrandt & Ebers (2006) recommend a 16S rRNA gene sequence similarity threshold of 98.7–99% as the point at which DNA–DNA reassociation experiments should be mandatory for testing the genomic uniqueness of a novel isolate(s). Physiological and biochemical characterization of strain ID05-A0528^T was also performed for comparison with recognized species of the genus *Dietzia* (Table 1).

On the basis of physiological and genotypic data, strain ID05-A0528^T represents a novel species within the genus *Dietzia*, for which the name *Dietzia timorensis* sp. nov. is proposed.

Description of *Dietzia timorensis* sp. nov.

Dietzia timorensis (ti.mo.ren'sis. N.L. fem. adj. *timorensis* pertaining to West Timor, Indonesia, from where the organism was first isolated).

Fig. 1. Phylogenetic tree derived from 16S rRNA gene sequences showing the relationship of strain ID05-A0528^T to species of the genus *Dietzia*. The tree was reconstructed using the neighbour-joining method and K_{nuc} values (Saitou & Nei, 1987). Asterisks indicate branches of the tree that were also recovered using the minimum evolution and maximum-parsimony methods (Takahashi & Nei, 2000). Bar, 0.01 nucleotide substitution rate (K_{nuc}).

Table 1. Differential characteristics of strain ID05-A0528^T and type strains of species of the genus *Dietzia*

Strains: 1, ID05-A0528^T; 2, *Dietzia cinnamomea* IMMIB RIV-399^T; 3, *D. maris* DSM 43672^T; 4, *D. schimae* YIM 65001^T; 5, *D. psychrocaliphila* ILA-1^T; 6, *D. cercidiphylli* YIM 65002^T; 7, *D. natronolimnaea* CBS 107.95^T; 8, *D. kunjamensis* KT30-10^T; 9, *D. papillomatosus* N 1280^T. +, Positive; –, negative. Data for reference strains from Li *et al.* (2008).

Characteristic	1	2	3	4	5	6	7	8	9
Urea hydrolysis	–	+	+	–	+	+	+	–	+
Nitrate reduction	–	+	+	+	–	–	–	+	+
Growth temperature range (°C)	10–37	22–45	10–45	10–45	10–37	10–37	10–37	10–37	10–37
Utilization as sole carbon source									
D-Adonitol	+	–	–	–	+	–	–	–	+
L-Arabinose	+	–	–	–	–	+	–	–	+
Cellobiose	+	–	–	+	+	–	+	+	+
L-Fucose	+	+	–	–	–	–	–	+	+
Inositol	+	–	–	–	+	–	–	–	+
Maltose	+	+	+	–	+	+	+	–	+
Raffinose	+	–	–	–	+	–	+	+	+
Salicin	+	+	–	–	–	–	–	–	+
Trehalose	+	–	–	–	+	–	+	+	+
D-Tagatose	+	+	–	–	–	+	–	–	+

Aerobic, Gram-positive, non-motile actinomycete that forms circular, convex, glistening, moderately orange-yellow colonies. Cells are rod- and coccoid-shaped. Diffusible pigments are not produced. Aesculin is hydrolysed. Nitrate is not reduced. Arbutin and urea are not hydrolysed. Does not degrade adenine, casein, elastin, hypoxanthine, testosterone, tyrosine, uric acid or xanthine. Grows at 10, 15, 28 and 37 °C, but does not grow at 5 or 45 °C. Growth occurs in the presence of 0–7 % NaCl (w/v). Utilizes aesculin, cellobiose, D-fructose, D-glucose, glycerol, lactose, D-mannose, sucrose, D-adonitol, amygdalin, L-arabinose, D-arabinose, L-arabitol, D-arabitol, arbutin, dulcitol, erythritol, L-fucose, D-fucose, D-galactose, gentiobiose, glycogen, inositol, inulin, D-lyxose, maltose, D-mannitol, melezitose, melibiose, methyl α -D-glucopyranoside, methyl α -D-mannopyranoside, methyl β -D-xylopyranoside, N-acetylglucosamine, potassium gluconate, potassium 2-ketogluconate, potassium 5-ketogluconate, raffinose, L-rhamnose, D-ribose, salicin, D-sorbitol, L-sorbitol, starch, D-tagatose, trehalose, turanose, xylitol, L-xylose and D-xylose. Contains *meso*-diaminopimelic acids, and arabinose and galactose are present in whole-cell hydrolysates. The acyl type of the glycan chain of the peptidoglycan is acetyl. The major fatty acids are C_{16:0}, C_{18:1} ω 9c and 10-methyl C_{18:0}. Mycolic acids are present. The polar lipid profile consists of phosphatidylglycerol and trace amounts of phosphatidylinositol. MK-8 (H₂) is the major menaquinone. The G+C content of DNA of the type strain is 65.5 mol%.

The type strain, ID05-A0528^T (=BTCC B-560^T =NBRC 104184^T), was isolated from a soil sample collected from under mahogany (*Swietenia mahogany*) trees on West Timor in Indonesia.

Acknowledgements

We thank Ms Yayoi Sakiyama, NITE Biological Resource Center, Department of Biotechnology, for analysis of chemotaxonomy. This work was conducted under a Joint Research Project between the Department of Biotechnology, National Institute of Technology and Evaluation, Japan, and the Indonesian Institute of Sciences (LIPI), representing Indonesian Government Research Institutes.

References

- Chaillan, F., Le Fleche, A., Bury, E., Phantavong, Y. H., Grimont, P., Saliot, A. & Oudot, J. (2004). Identification and biodegradation potential of tropical aerobic hydrocarbon-degrading microorganisms. *Res Microbiol* **155**, 587–595.
- Duckworth, A. W., Grant, S., Grant, W. D., Jones, B. E. & Meijer, D. (1998). *Dietzia natronolimnaea* sp. nov., a new member of the genus *Dietzia* isolated from an east African soda lake. *Extremophiles* **2**, 359–366.
- Felsenstein, J. (1985). Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* **39**, 783–791.
- Gerhardt, P., Murray, R. G. E., Costilow, R. N., Nester, E. W., Wood, W. A., Krieg, N. R. & Phillips, G. B. (editors) (1981). *Manual of Methods for General Bacteriology*. Washington, DC: American Society for Microbiology.
- Gordon, R. E. & Mihm, J. M. (1962). Identification of *Nocardia caviae* (Erikson) nov. comb. *Ann N Y Acad Sci* **98**, 628–636.
- Gordon, R. E., Barnett, D. A., Handerhan, J. E. & Pang, C. H.-N. (1974). *Nocardia coeliaca*, *Nocardia autotrophica*, and the nocardin strain. *Int J Syst Bacteriol* **24**, 54–63.
- Hasegawa, T., Takizawa, M. & Tanida, S. (1983). A rapid analysis for chemical grouping of aerobic actinomycetes. *J Gen Appl Microbiol* **29**, 319–322.
- Hatano, K., Nishii, T. & Kasai, H. (2003). Taxonomic re-evaluation of whorl-forming *Streptomyces* (formerly *Streptoverticillium*) species by using phenotypes, DNA–DNA hybridization and sequences of *gyrB*,

- and proposal of *Streptomyces luteireticuli* (ex Katoh and Arai 1957) coring., sp. nov., nom. rev. *Int J Syst Evol Microbiol* **53**, 1519–1529.
- Hayakawa, M. & Nonomura, H. (1987).** Humic acid-vitamin agar, a new medium for selective isolation of soil actinomycetes. *J Ferment Technol* **65**, 501–509.
- Hayakawa, M. & Nonomura, H. (1989).** A new method for the intensive isolation of actinomycetes from soil. *Actinomycetologica* **3**, 95–104.
- Isik, K., Chun, J., Hah, Y. C. & Goodfellow, M. (1999).** *Nocardia salmonicida* nom. rev., a fish pathogen. *Int J Syst Bacteriol* **49**, 833–837.
- Jones, J. (1949).** Fresh isolates of actinomycetes in which the presence of sporogenous aerial mycelia is a fluctuating characteristic. *J Bacteriol* **57**, 141–145.
- Jones, A. L., Koerner, R. J., Natarajan, S., Perry, J. D. & Goodfellow, M. (2008).** *Dietzia papillomatosis* sp. nov., a novel actinomycete isolated from the skin of an immunocompetent patient with confluent and reticulated papillomatosis. *Int J Syst Evol Microbiol* **58**, 68–72.
- Lechevalier, M. P. & Lechevalier, H. A. (1970).** Chemical composition as a criterion in the classification of aerobic actinomycetes. *Int J Syst Bacteriol* **20**, 435–443.
- Li, J., Zhao, G. Z., Zhang, Y. Q., Klenk, H. P., Pukall, R., Qin, S., Xu, L. H. & Li, W. J. (2008).** *Dietzia schimae* sp. nov. and *Dietzia cercidiphylli* sp. nov., from surface-sterilized plant tissues. *Int J Syst Evol Microbiol* **58**, 2549–2554.
- Mayilraj, S., Suresh, K., Kroppenstedt, R. M. & Saini, H. S. (2006).** *Dietzia kunjamensis* sp. nov., isolated from the Indian Himalayas. *Int J Syst Evol Microbiol* **56**, 1667–1671.
- Minnikin, D. E., O'Donnell, A. G., Goodfellow, M., Alderson, G., Athalye, M., Schaal, A. & Parlett, J. H. (1984).** An integrated procedure for the extraction of bacterial isoprenoid quinones and polar lipids. *J Microbiol Methods* **2**, 233–241.
- Rainey, F. A., Klatte, S., Kroppenstedt, R. M. & Stackebrandt, E. (1995).** *Dietzia*, a new genus including *Dietzia maris* comb. nov., formerly *Rhodococcus maris*. *Int J Syst Bacteriol* **45**, 32–36.
- Saito, H. & Miura, K.-I. (1963).** Preparation of transforming deoxyribonucleic acid by phenol treatment. *Biochim Biophys Acta* **72**, 619–629.
- Saitou, N. & Nei, M. (1987).** The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol* **4**, 406–425.
- Schaal, K. P. (1985).** Identification of clinically significant actinomycetes and related bacteria using chemical techniques. In *Chemical Methods in Bacterial Systematics*, pp 359–381. Edited by M. Goodfellow & D. E. Minnikin. London: Academic Press.
- Stackebrandt, E. & Ebers, J. (2006).** Taxonomic parameters revisited: tarnished gold standards. *Microbiol Today* **33**, 152–155.
- Stackebrandt, E., Rainey, F. A. & Ward-Rainey, N. L. (1997).** Proposal for a new hierarchic classification system, *Actinobacteria* classis nov. *Int J Syst Bacteriol* **47**, 479–491.
- Takahashi, K. & Nei, M. (2000).** Efficiencies of fast algorithms of phylogenetic inference under the criteria of maximum parsimony, minimum evolution, and maximum likelihood when a large number of sequences are used. *Mol Biol Evol* **17**, 1251–1258.
- Takeishi, H., Anzai, H., Urai, M., Aizawa, T., Wada, N., Iwabuchi, N., Sunairi, M. & Nakajima, M. (2006).** Xylanolytic and alkaliphilic *Dietzia* sp. isolated from larvae of the Japanese horned beetle, *Trypoxylus dichotomus*. *Actinomycetologica* **20**, 49–54.
- Tamura, T. & Hatano, K. (2001).** Phylogenetic analysis of the genus *Actinoplanes* and transfer of *Actinoplanes minutisporangius* Ruan *et al.* 1986 and '*Actinoplanes aurantiacus*' to *Cryptosporangium minutisporangium* comb. nov. and *Cryptosporangium aurantiacum* sp. nov. *Int J Syst Evol Microbiol* **51**, 2119–2125.
- Tamura, T., Nakagaito, Y., Nishii, T., Hasegawa, T., Stackebrandt, E. & Yokota, A. (1994).** A new genus of the order *Actinomycetales*, *Couchioplanes* gen. nov., with descriptions of *Couchioplanes caeruleus* (Horan and Brodsky 1986) comb. nov. and *Couchioplanes caeruleus* subsp. *azureus* subsp. nov. *Int J Syst Bacteriol* **44**, 193–203.
- Thompson, J. D., Gibson, T. J., Plewniak, F., Jeanmougin, F. & Higgins, D. G. (1997).** The CLUSTAL_X Windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res* **25**, 4876–4882.
- Uchida, K. & Aida, K. (1977).** Acyl type of bacterial cell wall: its simple identification by colorimetric method. *J Gen Appl Microbiol* **23**, 249–260.
- Williams, S. T., Goodfellow, M., Alderson, G., Wellington, E. M. H., Sneath, P. H. A. & Sackin, M. J. (1983).** Numerical classification of *Streptomyces* and related genera. *J Gen Microbiol* **129**, 1743–1813.
- Yassin, A. F., Hupfer, H. & Schaal, K. P. (2006).** *Dietzia cinnamea* sp. nov., a novel species isolated from a perianal swab of a patient with a bone marrow transplant. *Int J Syst Evol Microbiol* **56**, 641–645.
- Yumoto, I., Nakamura, A., Iwata, H., Kojima, K., Kusumoto, K., Nodasaka, Y. & Matsuyama, H. (2002).** *Dietzia psychralcaliphila* sp. nov., a novel, facultatively psychrophilic alkaliphile that grows on hydrocarbons. *Int J Syst Evol Microbiol* **52**, 85–90.