Bacillus marmarensis sp. nov., an alkaliphilic, protease-producing bacterium isolated from mushroom compost

Aziz Akin Denizci,¹ Dilek Kazan^{1,2} and Altan Erarslan³

Correspondence

Aziz Akin Denizci

akin.denizci@mam.gov.tr

¹Scientific and Technological Research Council of Turkey, Marmara Research Center, Genetic Engineering and Biotechnology Institute, PO Box 21, 41470 Gebze – Kocaeli, Turkey

²Marmara University, Faculty of Engineering, Bioengineering Department, Göztepe Campus, 34722 Kadıköy, İstanbul, Turkey

³Kocaeli University, Faculty of Arts and Sciences, Department of Chemistry, Section of Biochemistry, Umuttepe Central Campus, 41380 İzmit – Kocaeli, Turkey

A Gram-stain-positive, obligately alkaliphilic bacterium designated strain GMBE 72^T was isolated from mushroom compost from Yalova, located in the Marmara region of Turkey. Cells were aerobic, straight rods and they formed subterminal to terminal ellipsoidal endospores. The isolate was catalase-positive, oxidase-negative and motile and contained a type A1 γ peptidoglycan based on meso-diaminopimelic acid. The strain grew at pH 8.0–12.5. The major cellular fatty acid was anteiso-C_{15:0}. The genomic DNA G + C content was 40.2 mol%. Phylogenetic analyses based on 16S rRNA gene sequencing showed that strain GMBE 72^T belonged to the genus *Bacillus* and exhibited 98.2 % sequence similarity to *Bacillus pseudofirmus* DSM 8715^T. DNA–DNA reassociation was 56 % between GMBE 72^T and *B. pseudofirmus* DSM 8715^T. According to our polyphasic characterization, strain GMBE 72^T represents a novel species of the genus *Bacillus*, for which the name *Bacillus marmarensis* sp. nov. is proposed. The type strain is GMBE 72^T (=DSM 21297^T =JCM 15719^T).

Many alkaliphilic *Bacillus* species belong to the sixth rRNA group in the genus *Bacillus* (Nielsen *et al.*, 1994) and have been isolated from different habitats, as described by Agnew *et al.* (1995), Nielsen *et al.* (1995) and Lim *et al.* (2006a, b). They have been studied extensively because of their taxonomic and ecological significance and also for their use for biotechnological and industrial purposes (Fritze *et al.*, 1990; Horikoshi, 1996; Kumar & Takagi, 1999; Denizci *et al.*, 2004; Nogi *et al.*, 2005). Alkaliphilic bacteria have been isolated in order to investigate their diversity in terms of physiological adaptation to high pH and the industrial applications of their enzymes such as alkaline protease (Horikoshi, 1996, 1999; Kumar & Takagi, 1999; Yumoto *et al.*, 1998; Jasvir *et al.*, 1999; Takami *et al.*, 1999; Takami & Horikoshi, 2000; Takami & Krulwich, 2000; Gupta *et al.*, 2002; Saeki *et al.*, 2002). At the time of writing, 26 alkaliphilic and alkalitolerant *Bacillus* species have been identified (Vedder, 1934; Spanka & Fritze, 1993; Nielsen *et al.*, 1995; Agnew *et al.*, 1995; Fritze, 1996; Switzer

Blum *et al.*, 1998; Yumoto *et al.*, 1998, 2003; Olivera *et al.*, 2005; Nogi *et al.*, 2005; Ghosh *et al.*, 2007; Lee *et al.*, 2008; Borsodi *et al.*, 2008).

In this study, we describe the results of a polyphasic study aimed at the characterization of strain GMBE 72^T, which was isolated from mushroom compost from the Marmara region of Turkey during screening for alkaliphilic bacteria (Oner *et al.*, 2006). Compost samples were diluted serially with 0.9 % (w/v) saline solution and spread on agar containing 0.1 % glucose, 0.2 % peptone, 0.5 % yeast extract, 0.1 % K₂HPO₄, 0.02 % MgSO₄·7H₂O and 0.5 % skimmed milk (skimmed milk was sterilized separately). The medium was adjusted to pH 10.0 by addition of 10 % Na₂CO₃ solution after sterilization and plates were incubated for 72 h at 30 °C. A clear zone around colonies indicated protease-producing micro-organisms; isolate GMBE 72^T was selected as an alkaline protease producer on the basis of the diameter of the clear zone around its colonies. Subculturing was performed on DSMZ 31 medium (http://www.dsmz.de/microorganisms/medium/pdf/DSMZ_Medium31.pdf) for 24 h at 30 °C and the isolate was maintained as glycerol stocks at –70 °C.

Colony morphology was examined by growing strains on DSMZ 31 medium for 24 h at 30 °C. Cell morphology was

The GenBank/EMBL/DDBJ accession number for the 16S rRNA gene sequence of strain GMBE 72^T is EU621902.

Micrographs of strain GMBE 72^T and a maximum-parsimony 16S rRNA gene sequence-based tree are available as supplementary material with the online version of this paper.

investigated with light microscopy (Leica DM 4000) at $\times 1000$ magnification. Gram staining was determined with the Sigma Gram stain kit according to the manufacturer's instructions. Motility was observed after 12 and 24 h in semi-solid DSMZ 31 medium. Growth was tested in DSMZ 31 medium at 5–65 °C. The NaCl requirement for growth was studied in DSMZ 31 medium supplemented with 0–20 % (w/v) NaCl. Growth at different pH values was tested according to Nielsen *et al.* (1995). All physiological and biochemical tests were performed using the methods of Gordon *et al.* (1973). The medium was adjusted to pH 10 as described by Nielsen *et al.* (1995). Carbon source utilization tests were performed using API 50 CH galleries according to Nielsen *et al.* (1995). Tests were performed twice and unclear results are not given.

The cell-wall diamino acid type of strain GMBE 72^T was determined from whole-cell hydrolysates as described by Hasegawa *et al.* (1983). For cellular fatty acid analyses, the

strain was grown on TSA at 28 °C for 24 h and fatty acid methyl ester analysis was performed according to the instructions of the Sherlock Microbial Identification System (MIDI; Microbial ID). Determination of the DNA G+C content and DNA–DNA hybridization were performed by the Identification Service of the Deutsche Sammlung von Mikroorganismen und Zellkulturen (Braunschweig, Germany). Genomic DNA was isolated using DNAzol (Molecular Research Center, Inc.) according to the manufacturer's instructions. PCR amplification of the 16S rRNA gene was performed with universal primers 27f (5'-AGAGTTTGATCCTCAG-3') and 1385r (5'-CGG-TGTGTAGCAAGGCC-3'). An amplicon of approximately 1.4 kb was purified by using the EZ-10 Spin Column PCR purification kit (Bio Basic Inc.) according to the manufacturer's instructions. DNA sequencing using four primers, namely 27f, 357f, 907r, 1385r, was performed by the dideoxy chain-termination method with an ABI Prism 3100 DNA analyser using an ABI Prism Big Dye

Table 1. Phenotypic characteristics that serve to differentiate strain GMBE 72^T from type strains of phylogenetically related *Bacillus* species

Strains: 1, *Bacillus marmarensis* sp. nov. GMBE 72^T; 2, *B. pseudofirmus* DSM 8715^T (data from this study); 3, *B. wakoensis* N-1^T (data from Nogi *et al.*, 2005); 4, *B. krulwichiae* JCM 1169^T (Yumoto *et al.*, 2003). ND, No data available; v, variable.

Characteristic	1	2	3	4
O ₂ requirement	Aerobic	Strictly aerobic	ND	Aerobic/anaerobic
Colony morphology				
Pigmentation	Cream–yellowish	Yellow	Yellowish	Colourless
Margin	Entire	Irregular	ND	ND
Cell size (µm)	0.8–1.1 × 2–2.5	0.6–0.8 × 2.0–4.0	0.5–0.8 × 1.5–2.0	0.5–0.7 × 1.5–2.6
Growth with/at:				
12 % NaCl	+	+	–	+
14 % NaCl	–	+	–	+
pH 11	+	–	–	–
10 °C	–	+	+	–
45 °C	+	+	+	–
Hydrolysis of:				
Hippurate	–	+	ND	+
Starch	–	+	+	+
Gelatin	+	+	–	v
Casein	+	+	–	v
Tween 20	–	–	–	+
Tweens 40 and 60	–	+	–	+
Tween 80	+	–	ND	+
Reduction of nitrate	–	–	+	+
Growth on:				
D-Mannose	+	+	+	–
D-Xylose	–	+	+	+
D-Ribose	+	+	–	+
Cellobiose	+	+	+	–
D-Sorbitol	–	–	+	–
D-Galactose	–	–	–	+
DNA G+C content (mol%)	40.2	39	38.1	41

Terminator cycle sequencing ready reaction kit (Applied Biosystems) according to the manufacturer's protocol. 16S rRNA gene sequences of close relatives of strain GMBE 72^T with validly published names were retrieved from the GenBank database using BLASTN (Altschul *et al.*, 1997). A phylogenetic tree was constructed by using the software package MEGA version 4 (Kumar *et al.*, 2004) after multiple alignment of the data by CLUSTAL_X (Thompson *et al.*, 1997). Distances (distance options according to Kimura's two-parameter model; Kimura, 1980, 1983) and clustering were based on the neighbour-joining and maximum-parsimony methods. The tree topology was examined by the bootstrap method of resampling (Felsenstein, 1985) using 1000 bootstraps.

Strain GMBE 72^T formed cream–yellowish-coloured, circular, entire and convex colonies on DSMZ 31 medium. Cells were Gram-stain-positive, aerobic, motile, straight rods (0.8–1.1 × 2.0–2.5 µm). Cells were mainly single or occasionally in pairs or in short chains, with subterminal to terminal endospores (Supplementary Fig. S1, available in IJSEM Online). The strain was catalase-positive and oxidase-negative and did not reduce nitrate to nitrite. Growth was observed at 15–45 °C. The strain grew at pH 8.0–12.5; no growth was observed at pH 7.0. The strain grew at salt concentrations in the range 0–12% (w/v) NaCl. The growth and phenotypic characteristics that serve to distinguish strain GMBE 72^T from the type strains of closely related *Bacillus* species are shown in Table 1.

The cell wall contained *meso*-diaminopimelic acid in the peptidoglycan, suggesting the occurrence of peptidoglycan type A1 γ (as in the case of the great majority of members of the genus *Bacillus*; Claus & Berkeley, 1986).

The cellular fatty acid composition of GMBE 72^T was dominated by anteiso-C_{15:0} (41.1%). The type strain of the most closely related species, *Bacillus pseudofirmus* DSM 8715^T, also exhibited anteiso-C_{15:0} as the predominant fatty acid; iso-C_{15:0} was the second most abundant fatty acid in both strains (Table 2). This fatty acid profile corresponded well with the profile of the type species of the genus, *Bacillus subtilis* (Kämpfer, 1994), including the major acids anteiso-C_{15:0} and iso-C_{15:0}.

Comparative 16S rRNA gene sequencing analyses showed that strain GMBE 72^T was closely related to *B. pseudofirmus* DSM 8715^T (98.2%), *Bacillus wakoensis* N-1^T (93.4%), *Bacillus krulwichiae* AM31D^T (93.2%) and *B. subtilis* 168 (92.4%). The phylogenetic tree reconstructed using the neighbour-joining method showed that strain GMBE 72^T is a member of group 6 (Nielsen *et al.*, 1994) of the genus *Bacillus*. Strain GMBE 72^T formed a clade with *B. pseudofirmus* DSM 8715^T with a bootstrap value of 100% (Fig. 1). The maximum-parsimony phylogenetic tree showed essentially the same position for strain GMBE 72^T (Supplementary Fig. S2). The DNA G + C content of strain GMBE 72^T was 40.2 mol%.

Table 2. Fatty acid profiles of strain GMBE 72^T and *B. pseudofirmus* DSM 8715^T

Values are percentages of total fatty acid. Data were determined in this study; only fatty acids accounting for at least 1% of the total fatty acids are listed. —, Not detected (<1.0%).

Fatty acid	GMBE 72 ^T	<i>B. pseudofirmus</i> DSM 8715 ^T
iso-C _{10:0}	—	1.0
iso-C _{14:0}	5.0	3.6
C _{14:0}	1.7	1.6
iso-C _{15:0}	24.2	16.5
anteiso-C _{15:0}	41.1	50.8
C _{16:1} ω 7 <i>c</i> alcohol	6.9	5.1
iso-C _{16:0}	2.7	2.5
C _{16:1} ω 11 <i>c</i>	4.5	4.7
C _{16:0}	2.0	2.4
anteiso-C _{17:0}	4.5	4.7
iso-C _{17:0}	1.0	—
C _{18:1} ω 9 <i>c</i>	1.1	—
Summed feature*	3.7	3.1

*Summed feature containing iso-C_{17:1} I and/or anteiso-C_{17:1} B.

The mean value for DNA–DNA hybridization between strain GMBE 72^T and *B. pseudofirmus* DSM 8715^T was 56% (individual values 54.4 and 57.6%). This value for hybridization is lower than the recommended threshold value accepted for defining a novel species (Wayne *et al.*, 1987), supporting the distinct position of strain GMBE 72^T within the genus *Bacillus*. Phenotypic properties of strain GMBE 72^T, such as the ability to hydrolyse Tween 80 and

Fig. 1. Neighbour-joining phylogenetic tree based on 16S rRNA gene sequences, showing the position of strain GMBE 72^T. Bootstrap values are shown as percentages of 1000 replicates; only values above 50% are shown. Bar, 0.01 substitutions per nucleotide position.

the inability to hydrolyse starch and to grow in 14 % NaCl, also supported the view that the isolate was distinguishable from closely related *Bacillus* species (Table 1). Therefore, on the basis of physiological, biochemical and phylogenetic properties, strain GMBE 72^T represents a novel species within the genus *Bacillus*, for which the name *Bacillus marmarensis* sp. nov. is proposed.

Description of *Bacillus marmarensis* sp. nov.

Bacillus marmarensis (mar.ma.ren'sis. N.L. masc. adj. *marmarensis* pertaining to the region of Marmara, where the type strain was isolated).

Cells are aerobic, Gram-stain-positive, motile rods (0.8–1.1 × 2.0–2.5 µm) that produce ellipsoidal spores that are located subterminally to terminally. Colonies are cream–yellowish, circular, entire and convex. Catalase-positive and oxidase-negative. No growth is detected at pH 7.0–7.5. Growth occurs at pH 8–12.5 and 10–45 °C. The organism is able to grow at 12 % NaCl. It hydrolyses casein, gelatin and Tween 80, but does not hydrolyse urea, starch or Tweens 20, 40 or 60. Does not reduce nitrate to nitrite and does not utilize Simmons' citrate. Heterotrophic, utilizing D- and L-arabinose, ribose, glucose, fructose, mannose, methyl α-D-mannoside, methyl α-D-glucoside, N-acetylglucosamine, salicin, maltose, cellobiose, trehalose, sucrose, glycogen, xylitol, gentiobiose, turanose, D-arabitol and 2-ketogluconate, but not utilizing glycerol, erythritol, D- or L-xylose, adonitol, methyl β-D-xyloside, galactose, sorbose, rhamnose, dulcitol, inositol, mannitol, sorbitol, amygdalin, arbutin, aesculin, lactose, melibiose, inulin, melezitose, raffinose, D-lyxose, D-tagatose, D- or L-fucose, L-arabitol, gluconate or 5-ketogluconate. Major fatty acids are anteiso-C_{15:0} and iso-C_{15:0}. The detailed fatty acid profile is listed in Table 2. The cell-wall peptidoglycan contains meso-diaminopimelic acid. The DNA G+C content of the type strain is 40.2 mol% (determined by HPLC).

The type strain, GMBE 72^T (=DSM 21297^T =JCM 15719^T), was isolated from mushroom compost from Yalova, Turkey.

Acknowledgements

This research was supported by the Scientific and Technical Research Council of Turkey (TÜBİTAK; project number TBAG 2129-102T003). The authors thank Dr Nurettin Sahin for his valuable comments, Dr Bayram Yüksel for his valuable comments on phylogenetic analyses and Mrs Nesrin Karahan for her valuable technical assistance and the DSMZ Identification Service for performing the DNA–DNA hybridization and G+C content determination.

References

Agnew, M. D., Koval, S. F. & Jarrell, K. F. (1995). Isolation and characterization of novel alkaliphiles from bauxite-processing waste and description of *Bacillus vedderi* sp. nov., a new obligate alkaliphile. *Syst Appl Microbiol* **18**, 221–230.

Altschul, S. F., Madden, T. L., Schaffer, A. A., Zhang, J., Zhang, Z., Miller, W. & Lipman, D. J. (1997). Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res* **25**, 3389–3402.

Borsodi, A. K., Márialigeti, K., Szabó, G., Palatinszky, M., Pollák, B., Kéki, Z., Kovács, A. L., Schumann, P. & Tóth, E. M. (2008). *Bacillus aurantiacus* sp. nov., an alkaliphilic and moderately halophilic bacterium isolated from Hungarian soda lakes. *Int J Syst Evol Microbiol* **58**, 845–851.

Claus, D. & Berkeley, R. C. W. (1986). Genus *Bacillus* Cohn 1872. In *Bergey's Manual of Systematic Bacteriology*, vol. 2, pp. 1105–1140. Edited by P. H. A. Sneath, N. S. Mair, M. E. Sharpe & J. G. Holt. Baltimore: Williams & Wilkins.

Denizci, A. A., Kazan, D., Abeln, E. C. A. & Erarslan, A. (2004). Newly isolated *Bacillus clausii* GMBAE 42: an alkaline protease producer capable to grow under highly alkaline conditions. *J Appl Microbiol* **96**, 320–327.

Felsenstein, J. (1985). Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* **39**, 783–791.

Fritze, D. (1996). *Bacillus haloalkaliphilus* sp. nov. *Int J Syst Bacteriol* **46**, 98–101.

Fritze, D., Flossdorf, J. & Claus, D. (1990). Taxonomy of alkaliphilic *Bacillus* strains. *Int J Syst Bacteriol* **40**, 92–97.

Ghosh, A., Bhardwaj, M., Satyanarayana, T., Khurana, M., Mayilraj, S. & Jain, R. K. (2007). *Bacillus lehensis* sp. nov., an alkalitolerant bacterium isolated from soil. *Int J Syst Evol Microbiol* **57**, 238–242.

Gordon, R. E., Haynes, W. C. & Pang, C. H.-N. (1973). *The genus Bacillus* (Agriculture Handbook no. 427). Washington, DC: Agricultural Research Service, United States Department of Agriculture.

Gupta, R., Beg, Q. K. & Lorenz, P. (2002). Bacterial alkaline proteases: molecular approaches and industrial applications. *Appl Microbiol Biotechnol* **59**, 15–32.

Hasegawa, T., Takizawa, M. & Tanida, S. (1983). A rapid analysis for chemical grouping of aerobic actinomycetes. *J Gen Appl Microbiol* **29**, 319–322.

Horikoshi, K. (1996). Alkaliphiles – from an industrial point of view. *FEMS Microbiol Rev* **18**, 259–270.

Horikoshi, K. (1999). Alkaliphiles: some applications of their products for biotechnology. *Microbiol Mol Biol Rev* **63**, 735–750.

Jasvir, S., Navdeep, G., Devasahayam, G. & Sahoo, D. K. (1999). Studies on alkaline protease produced by *Bacillus* sp. NG312. *Appl Biochem Biotechnol* **76**, 57–63.

Kämpfer, P. (1994). Limits and possibilities of total fatty acid analysis for classification and identification of *Bacillus* species. *Syst Appl Microbiol* **17**, 86–98.

Kimura, M. (1980). A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. *J Mol Evol* **16**, 111–120.

Kimura, M. (1983). *The Neutral Theory of Molecular Evolution*. Cambridge: Cambridge University.

Kumar, C. G. & Takagi, H. (1999). Microbial alkaline proteases: from a bioindustrial viewpoint. *Biotechnol Adv* **17**, 561–594.

Kumar, S., Tamura, K. & Nei, M. (2004). MEGA3: integrated software for molecular evolutionary genetics analysis and sequence alignment. *Brief Bioinform* **5**, 150–163.

Lee, J.-C., Lee, G. S., Park, D.-J. & Kim, C.-J. (2008). *Bacillus alkalitelluris* sp. nov., an alkaliphilic bacterium isolated from sandy soil. *Int J Syst Evol Microbiol* **58**, 2629–2634.

Lim, J.-M., Jeon, C. O., Lee, S.-M., Lee, J.-C., Xu, L.-H., Jiang, C.-L. & Kim, C.-J. (2006a). *Bacillus salarius* sp. nov., a halophilic, spore-

- forming bacterium isolated from a salt lake in China. *Int J Syst Evol Microbiol* **56**, 373–377.
- Lim, J.-M., Jeon, C. O. & Kim, C.-J. (2006b). *Bacillus taeansensis* sp. nov., a halophilic Gram-positive bacterium from a solar saltern in Korea. *Int J Syst Evol Microbiol* **56**, 2903–2908.
- Nielsen, P., Rainey, F. A., Outtrup, H., Priest, F. G. & Fritze, D. (1994). Comparative 16S rDNA sequence analysis of some alkaliphilic bacilli and the establishment of a sixth rRNA group within the genus *Bacillus*. *FEMS Microbiol Lett* **117**, 61–66.
- Nielsen, P., Fritze, D. & Priest, F. G. (1995). Phenetic diversity of alkaliphilic *Bacillus* strains: proposal for nine new species. *Microbiology* **141**, 1745–1761.
- Nogi, Y., Takami, H. & Horikoshi, K. (2005). Characterization of alkaliphilic *Bacillus* strains used in industry: proposal of five novel species. *Int J Syst Evol Microbiol* **55**, 2309–2315.
- Olivera, N., Sineriz, F. & Breccia, J. D. (2005). *Bacillus patagoniensis* sp. nov., a novel alkali-tolerant bacterium from the rhizosphere of *Atriplex lampa* in Patagonia, Argentina. *Int J Syst Evol Microbiol* **55**, 443–447.
- Oner, M. N. K., Denizci, A. A., Kazan, D. & Erarslan, A. (2006). A serine alkaline protease from a newly isolated obligate alkaliphilic *Bacillus* sp. GMBAE 72. *FEBS J* **273** (Suppl. 1), 335.
- Saeki, K., Hitomi, J., Okuda, M., Hatada, Y., Kageyama, Y., Takaiwa, M., Kubota, H. & Hagihara, H. (2002). A novel species of alkaliphilic *Bacillus* that produced an oxidatively stable alkaline serine protease. *Extremophiles* **6**, 65–72.
- Spanka, R. & Fritze, D. (1993). *Bacillus cohnii* sp. nov., a new, obligately alkaliphilic, oval-spore-forming *Bacillus* species with ornithine and aspartic acid instead of diaminopimelic acid in the cell wall. *Int J Syst Bacteriol* **43**, 150–156.
- Switzer Blum, J., Burns Bindi, A., Buzzelli, J., Stolz, J. F. & Oremland, R. S. (1998). *Bacillus arsenicoselenatis*, sp. nov., and *Bacillus selenitireducens*, sp. nov.: two haloalkaliphiles from Mono Lake, California that respire oxyanions of selenium and arsenic. *Arch Microbiol* **171**, 19–30.
- Takami, H. & Horikoshi, K. (2000). Analyses of the genome of an alkaliphilic *Bacillus* strains from an industrial point of view. *Extremophiles* **4**, 99–108.
- Takami, H. & Krulwich, T. A. (2000). Reidentification of facultatively alkaliphilic *Bacillus firmus* OF4 as *Bacillus pseudofirmus* OF4. *Extremophiles* **4**, 19–22.
- Takami, H., Nogi, Y. & Horikoshi, K. (1999). Reidentification of the keratinase producing facultatively alkaliphilic *Bacillus* sp. AH-101 as *Bacillus halodurans*. *Extremophiles* **3**, 293–296.
- Thompson, J. D., Gibson, T. J., Plewniak, F., Jeanmougin, F. & Higgins, D. G. (1997). The CLUSTAL_X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res* **25**, 4876–4882.
- Vedder, A. (1934). *Bacillus alcalophilus* n. sp.; benevens enkele ervaringen met sterk alcalische voedingbodems. *Antonie van Leeuwenhoek* **1**, 141–147 (in Dutch).
- Wayne, L. G., Brenner, D. J., Colwell, R. R., Grimont, P. A. D., Kandler, O., Krichevsky, M. I., Moore, L. H., Moore, W. E. C., Murray, R. G. E. & other authors (1987). International Committee on Systematic Bacteriology. Report of the ad hoc committee on reconciliation of approaches to bacterial systematics. *Int J Syst Bacteriol* **37**, 463–464.
- Yumoto, I., Yamazaki, K., Sawabe, T., Nakano, K., Kawasaki, K., Ezura, Y. & Shinano, H. (1998). *Bacillus horti* sp. nov., a new Gram-negative alkaliphilic bacillus. *Int J Syst Bacteriol* **48**, 565–571.
- Yumoto, I., Yamaga, S., Sogabe, Y., Nodasaka, Y., Matsuyama, H., Nakajima, K. & Suemori, A. (2003). *Bacillus krulwichiae* sp. nov., a halotolerant obligate alkaliphile that utilizes benzoate and *m*-hydroxybenzoate. *Int J Syst Evol Microbiol* **53**, 1531–1536.