Thalassomonas agariperforans sp. nov., an agarolytic bacterium isolated from marine sand

Sooyeon Park,¹ Won-Chan Choi,¹ Tae-Kwang Oh¹ and Jung-Hoon Yoon^{1,2}

Correspondence

Jung-Hoon Yoon
jhyoon69@skku.edu

¹Korea Research Institute of Bioscience and Biotechnology (KRIBB), PO Box 115, Yusong, Taejeon, Republic of Korea

²Department of Food Science and Biotechnology, Sungkyunkwan University, Jangan-gu, Suwon, Republic of Korea

A Gram-staining-negative, motile, agarolytic bacterium, designated M-M1^T, was isolated from marine sand obtained from Geoje Island, South Sea, Korea, and its taxonomic position was investigated using a polyphasic taxonomic approach. Strain M-M1^T grew optimally at pH 7.0–8.0, at 30 °C and in the presence of 2 % (w/v) NaCl. It did not grow in the presence of >7 % (w/v) NaCl. Phylogenetic analysis based on 16S rRNA gene sequences showed that strain M-M1^T fell within the clade comprising members of the genus *Thalassomonas*, clustering with *Thalassomonas agarivorans* TMA1^T, *Thalassomonas loyana* CBMAI 722^T and *Thalassomonas ganghwensis* JC2041^T, with which it exhibited 16S rRNA gene sequence similarity values of 96.4, 96.0 and 94.9 % respectively. Strain M-M1^T exhibited 94.7–95.2 % 16S rRNA gene sequence similarity to the other species of the genus *Thalassomonas*. Strain M-M1^T contained Q-8 as the predominant ubiquinone and C_{16:1}ω7c and/or iso-C_{15:0} 2-OH, C_{16:0} and C_{18:1}ω7c as the major fatty acids. The DNA G + C content was 44.2 mol%. Strain M-M1^T could be differentiated from phylogenetically related species of the genus *Thalassomonas* by differences in some phenotypic properties. On the basis of the phenotypic, chemotaxonomic and phylogenetic data, strain M-M1^T is considered to represent a novel species of the genus *Thalassomonas*, for which the name *Thalassomonas agariperforans* sp. nov. is proposed. The type strain is M-M1^T (=KCTC 23343^T =CCUG 60020^T).

The genus *Thalassomonas* belongs to the order *Alteromonadales* (Bowman & McMeekin, 2005) in the class *Gammaproteobacteria* and was first proposed by Macián *et al.* (2001). At the time of writing, the genus *Thalassomonas* consists of six species with validly published names: *Thalassomonas viridans* (Macián *et al.*, 2001), *Thalassomonas ganghwensis* (Yi *et al.*, 2004), *Thalassomonas loyana* (Thompson *et al.*, 2006), *Thalassomonas agarivorans* (Jean *et al.*, 2006) and *Thalassomonas actiniarum* and *Thalassomonas haliotis* (Hosoya *et al.*, 2009). In this study, an agarolytic bacterial strain, M-M1^T, which was isolated from marine sand of Geoje island in the South Sea, Korea, is described. Comparative 16S rRNA gene sequence analysis indicated that strain M-M1^T is most phylogenetically affiliated to the genus *Thalassomonas*. The aim of the present work was to determine the exact taxonomic position of strain M-M1^T by using a polyphasic characterization that included the determination of phenotypic properties and a detailed phylogenetic investigation based on 16S rRNA gene sequences.

Strain M-M1^T was isolated by the dilution plating technique on marine agar 2216 (MA; Difco) at 25 °C. The type strains

of three species of the genus *Thalassomonas* were used as reference strains for phenotypic characterization and fatty acid analysis: *T. agarivorans* JCM 13379^T was obtained from the Japan Collection of Microorganisms, Saitama, Japan; *T. loyana* LMG 22536^T was obtained from the Laboratorium voor Microbiologie, Universiteit Gent, Belgium; and *T. ganghwensis* KCTC 12041^T was obtained from the Korean Collection for Type Cultures, Taejeon, Korea. The morphological, physiological and biochemical characteristics of strain M-M1^T were investigated by using routine cultivation on MA at 30 °C. Cell morphology was examined by using light microscopy (Nikon E600) and transmission electron microscopy (CM-20; Philips). Flagellation was determined by using transmission electron microscopy on cells from an exponentially growing culture. For this purpose, the cells were negatively stained with 1 % (w/v) phosphotungstic acid and the grids were examined after being air-dried. The Gram reaction was determined by using the bioMérieux Gram stain kit according to the manufacturer's instructions. Growth at various temperatures (4, 10, 25, 30, 37, 40 and 45 °C) was measured on MA. Growth in the absence of NaCl and in the presence of 0.5 and 1.0 % (w/v) NaCl was investigated by using trypticase soy broth prepared according to the formula of the Difco medium except that NaCl was excluded and 0.45 % (w/v) MgCl₂·6H₂O or 0.06 %

The GenBank/EMBL/DDBJ accession number for the 16S rRNA gene sequence of strain M-M1^T is HM237288.

(w/v) KCl was added. Growth at various NaCl concentrations (2.0–15.0%, w/v, in increments of 1.0%) was investigated in marine broth 2216 (MB; Difco). The pH range for growth was determined in MB adjusted to pH 4.5–9.5 (in increments of 0.5 pH units) by the addition of HCl or Na₂CO₃. Growth under anaerobic conditions was determined after incubation in a Forma anaerobic chamber on MA and on MA supplemented with 0.1% (w/v) potassium nitrate, both of which had been prepared anaerobically under nitrogen atmosphere. Catalase and oxidase activities were determined as described by Cowan & Steel (1965). Hydrolysis of agar, casein, starch, hypoxanthine, xanthine and tyrosine was performed on MA, using the substrate concentrations described by Cowan & Steel (1965). Hydrolysis of aesculin, gelatin, Tweens (20, 40, 60 and 80) and urea, and nitrate reduction were investigated as described by Lanyi (1987) with the modification that artificial seawater (containing 23.6 g NaCl, 0.64 g KCl, 4.53 g MgCl₂·6H₂O, 5.94 g MgSO₄·7H₂O and 1.3 g CaCl₂·2H₂O per litre of distilled water; Bruns *et al.*, 2001) was used for preparation of media. Cellulase and xylanase activities were investigated on MA containing 0.5% carboxymethylcellulose (Sigma) and xylan (Sigma), respectively, and detected according to the method of Teather & Wood (1982). DNase activity was examined by using DNase test agar with methyl green (Difco) with the modification that artificial seawater was used for the preparation of media. H₂S production was tested as described by Bruns *et al.* (2001). Utilization of substrates as sole carbon and energy sources was tested as described by Baumann & Baumann (1981), using supplementation with 2% (v/v) Hutner's mineral base (Cohen-Bazire *et al.*, 1957) and 1% (v/v) vitamin solution (Staley, 1968). Susceptibility to antibiotics was investigated on MA plates by using antibiotic discs containing the following: polymyxin B, 100 U; streptomycin, 50 µg; penicillin G, 20 U; chloramphenicol, 100 µg; ampicillin, 10 µg; cephalothin, 30 µg; gentamicin, 30 µg; novobiocin, 5 µg; tetracycline, 30 µg; kanamycin, 30 µg; lincomycin, 15 µg; oleandomycin, 15 µg; neomycin, 30 µg; and carbenicillin, 100 µg. Other physiological and biochemical tests were performed using the API 20E and API ZYM systems (bioMérieux) incubated at 30 °C according to the instructions of the manufacturer.

Cell biomass of strain M-M1^T for DNA extraction and for isoprenoid quinone analysis was obtained from cultures grown in MB at 30 °C. Chromosomal DNA was extracted and purified according to the method described by Yoon *et al.* (1996), with the modification that RNase T1 was used in combination with RNase A to minimize contamination by RNA. 16S rRNA gene amplification was performed using two universal primers as described previously (Yoon *et al.*, 1998) and the PCR products were purified by using a QIAquick PCR purification kit (Qiagen). Sequencing of the amplified 16S rRNA gene and phylogenetic analysis were performed as described by Yoon *et al.* (2003). Isoprenoid quinones were analysed as described by Komagata & Suzuki (1987), using reversed-phase HPLC and a YMC ODS-A (250 × 4.6 mm)

Table 1. Differential phenotypic characteristics of strain M-M1^T and the type strains of phylogenetically related species of the genus *Thalassomonas*

Strains: 1, *Thalassomonas agariperforans* sp. nov. M-M1^T; 2, *T. agarivorans* JCM 13379^T; 3, *T. ganghwensis* KCTC 12041^T; 4, *T. loyana* LMG 22536^T. All data from this study except for NaCl range for growth, growth at 10 and 40 °C, catalase, oxidase, nitrate reduction and hydrolysis of aesculin, agar, DNA, gelatin, starch and urea; these data from Yi *et al.* (2004), Jean *et al.* (2006) and Thompson *et al.* (2006). +, Positive; –, negative; w, weak. All strains are Gram-negative and positive for catalase and nitrate reduction (weak for *T. loyana*), hydrolysis of aesculin, DNA and Tweens 20, 40, 60 and 80, utilization of D-galactose, D-glucose and maltose, activity of alkaline phosphatase, leucine arylamidase and naphthol-AS-BI-phosphohydrolase (weak for *T. agarivorans* and *T. loyana*) and susceptibility to streptomycin, novobiocin, neomycin, gentamicin, carbenicillin, polymyxin B and chloramphenicol. All strains are negative for production of indole and H₂S, hydrolysis of urea, xanthine and hypoxanthine, utilization of L-arabinose, trehalose, benzoate, citrate, formate, L-malate, succinate and L-glutamate, activity of arginine dihydrolase, lysine decarboxylase and ornithine decarboxylase, lipase (C14), valine arylamidase, cystine arylamidase, trypsin, α-chymotrypsin, α-galactosidase, β-galactosidase, β-glucuronidase, α-glucosidase, β-glucosidase, α-mannosidase and α-fucosidase and susceptibility to lincomycin and penicillin G.

Characteristic	1	2	3	4
NaCl range for growth (%)	0–6	1–5	1–8	1–9
Growth at:				
10 °C	+	–	–	–
40 °C	–	–	+	–
Oxidase	+	+	+	–
Hydrolysis of:				
Agar	+	+	–	–
Casein	–	+	+	+
Gelatin	–	+	+	w
Starch	+	+	–	+
Tyrosine	–	+	+	+
Utilization of:				
Cellobiose	+	+	–	+
D-Fructose	+	–	–	–
D-Mannose	–	w	–	–
D-Xylose	+	+	–	–
Sucrose	+	–	–	–
Acetate	–	–	–	+
Pyruvate	–	+	–	+
Salicin	+	+	–	–
Enzyme activity (API ZYM)				
Esterase (C4)	–	+	+	+
Esterase lipase (C8)	–	+	–	+
Acid phosphatase	+	–	+	–
N-Acetyl-β-glucosaminidase	–	+	–	–
Susceptibility to antibiotics				
Ampicillin	+	–	–	–
Cephalothin	–	+	+	+
Kanamycin	+	–	+	w
Oleandomycin	–	+	+	+
Tetracycline	–	–	+	–
DNA G + C content (mol%)	44.2	41.0	42.0	39.3

Fig. 1. Neighbour-joining phylogenetic tree based on 16S rRNA gene sequences showing the positions of strain M-M1^T, species of the genus *Thalassomonas* and some other related taxa. Bootstrap values (expressed as percentages of 1000 replications) >50% are shown at branching points. Filled circles indicate that the corresponding nodes were also recovered in trees generated with the maximum-likelihood and maximum-parsimony algorithms. *Pseudoalteromonas aeruginosa* DSM 50071^T (GenBank accession number, X06684) was used as an outgroup (not shown). Bar, 0.01 substitutions per nucleotide position.

column. For cellular fatty acid analysis, cell mass of strain M-M1^T, *T. agarivorans* JCM 13379^T, *T. loyana* LMG 22536^T and *T. ganghwensis* KCTC 12041^T was harvested from MA plates after cultivation for 5 days at 30 °C. The fatty acids were extracted and fatty acid methyl esters were prepared according to the standard protocol of the MIDI/Hewlett Packard Microbial Identification System (Sasser, 1990). The DNA G+C content was determined using the method of Tamaoka & Komagata (1984) with the modification that DNA was hydrolysed using nuclease P1 (Sigma) and the resultant nucleotides were analysed by reversed-phase HPLC.

Morphological, cultural, physiological and biochemical characteristics of strain M-M1^T are given in the species description (see below) and in Table 1. The almost complete 16S rRNA gene sequence of strain M-M1^T determined in this study comprised 1465 nt, representing approximately 96% of the *Escherichia coli* 16S rRNA gene sequence. In the phylogenetic tree based on the neighbour-joining algorithm, strain M-M1^T fell within the clade comprising species of the genus *Thalassomonas*, forming a coherent cluster with *T. agarivorans* TMA1^T, *T. loyana* CBMAI 722^T and *T. ganghwensis* JC2041^T (Fig. 1). Strain M-M1^T exhibited 16S rRNA gene sequence similarity values of 96.4, 96.0 and 94.9% with the type strains of *T. agarivorans*, *T. loyana* and *T. ganghwensis*, respectively, and 94.7–95.2% with the other species of the genus *Thalassomonas*. The predominant isoprenoid quinone detected in strain M-M1^T was ubiquinone-8 (Q-8), at a peak area ratio of approximately 94%, in line with other species of the genus *Thalassomonas* (Yi *et al.*, 2004; Hosoya *et al.*, 2009). The cellular fatty acid profile of strain M-M1^T is shown in Table 2, together with those of *T. agarivorans* JCM 13379^T, *T. loyana* LMG 22536^T and *T. ganghwensis* KCTC 12041^T also analysed in this study. Strain M-M1^T contained large amounts of straight-chain and unsaturated fatty acids; the major components (>10% of total fatty acids) were summed feature 3 (C_{16:1}ω7c and/or iso-C_{15:0} 2-OH), C_{16:0} and C_{18:1}ω7c. The fatty acid patterns of the four strains were similar, but there were also differences in the proportions of some fatty acids (Table 2). The DNA G+C content of strain M-M1^T was 44.2 mol%. The results

obtained from chemotaxonomic analyses, showing properties that are shared by species of the genus *Thalassomonas*, were in agreement with the result of the phylogenetic analysis, i.e. that strain M-M1^T is a member of the genus *Thalassomonas* (Yi *et al.*, 2004; Jean *et al.*, 2006; Thompson *et al.*, 2006).

Strain M-M1^T could be differentiated from *T. agarivorans* TMA1^T, *T. loyana* CBMAI 722^T and *T. ganghwensis* JC2041^T by differences in several phenotypic characteristics, including hydrolysis of casein, gelatin and tyrosine, utilization of D-fructose and sucrose, activity of esterase (C4) and susceptibility to ampicillin, oleandomycin and cephalothin as shown in Table 1. Therefore, on the basis of the phylogenetic data and differential phenotypic properties, strain M-M1^T is considered to represent a novel species of the genus *Thalassomonas*, for which the name *Thalassomonas agariperforans* sp. nov. is proposed.

Description of *Thalassomonas agariperforans* sp. nov.

Thalassomonas agariperforans [a.ga.ri.per.fo'rans. Malayan n. agar agar; N.L. n. agarum agar (algal polysaccharide); L. part. adj. perforans perforating, making holes; N.L. part. adj. agariperforans making holes in agar].

Cells are Gram-staining-negative, aerobic and rod-shaped, 0.5–0.8 µm in diameter and 1.5–3.0 µm in length. Motile by means of a single polar flagellum. Colonies on MA are circular, yellowish-white, opaque and sunken into the agar. Anaerobic growth occurs on MA and on MA supplemented with nitrate. Optimal growth temperature is 30 °C; growth occurs at 4 and 37 °C, but not at 40 °C. Optimal pH for growth is between 7.0 and 8.0; growth occurs at pH 5.5, but not at pH 5.0. Growth occurs in the presence of 0–6.0% (w/v) NaCl with an optimum in the presence of approximately 2.0% (w/v) NaCl. Mg²⁺ ions are required for growth. Cellulose and xylan are hydrolysed. The predominant ubiquinone is Q-8. The major fatty acids (>10% of total fatty acids) are C_{16:1}ω7c and/or iso-C_{15:0} 2-OH, C_{16:0} and C_{18:1}ω7c. The DNA G+C content is 44.2 mol% (determined by HPLC). Other phenotypic characteristics are given in Table 1.

Table 2. Cellular fatty acid compositions (%) of strain M-M1^T and the type strains of phylogenetically related species of the genus *Thalassomonas*

Strains: 1, *Thalassomonas agariperforans* sp. nov. M-M1^T; 2, *T. agarivorans* JCM 13379^T; 3, *T. ganghwensis* KCTC 12041^T; 4, *T. loyana* LMG 22536^T. All data from this study. Fatty acids that represented <1.0% in all strains were omitted. —, Not detected.

Fatty acid	1	2	3	4
Straight-chain				
C _{10:0}	0.7	1.3	3.5	—
C _{11:0}	—	1.6	0.1	0.2
C _{12:0}	1.6	6.4	0.3	2.8
C _{13:0}	—	5.3	—	0.8
C _{14:0}	1.6	4.2	0.9	9.7
C _{15:0}	0.8	6.1	1.2	2.6
C _{16:0}	30.9	19.5	19.5	9.4
C _{17:0}	0.8	6.8	1.4	0.5
C _{18:0}	0.5	1.1	1.0	0.8
Branched				
iso-C _{13:0}	—	1.4	—	—
iso-C _{14:0}	—	1.4	0.2	0.4
iso-C _{16:0}	—	0.7	2.8	—
Unsaturated				
C _{15:1} ω8c	0.5	2.0	1.8	2.1
C _{16:1} ω9c	—	3.7	4.2	4.8
C _{17:1} ω8c	—	9.2	7.2	10.6
C _{18:1} ω7c	13.9	4.0	19.2	8.6
C _{18:1} ω9c	—	2.2	1.7	2.9
Hydroxy				
C _{10:0} 3-OH	3.7	—	1.8	—
C _{11:0} 3-OH	—	0.6	1.6	2.2
C _{12:0} 3-OH	0.7	2.2	2.5	7.1
C _{12:1} 3-OH	4.6	—	—	—
Summed features*				
1	—	1.9	—	0.6
2	—	2.5	—	1.1
3	38.9	13.1	22.6	27.2
Unknown†				
ECL 11.799	—	0.2	0.3	1.9

*Summed features represent groups of two or three fatty acids that could not be separated by GLC with the MIDI system. Summed feature 1 contained iso-C_{15:1} and/or C_{13:0} 3-OH; summed feature 2 contained iso-C_{16:1} I and/or C_{14:0} 3-OH; summed feature 3 contained C_{16:1}ω7c and/or iso-C_{15:0} 2-OH.

†Equivalent chain-length.

The type strain, M-M1^T (=KCTC 23343^T =CCUG 60020^T), was isolated from marine sand of Geoje island in the South Sea, Korea.

Acknowledgements

This work was supported by the 21C Frontier Program of Microbial Genomics and Applications (grant 11-2008-00-002-00) from the Ministry of Education, Science and Technology (MEST) of the Republic of Korea.

References

- Baumann, P. & Baumann, L. (1981). The marine Gram-negative eubacteria: genera *Photobacterium*, *Beneckeia*, *Alteromonas*, *Pseudomonas*, and *Alcaligenes*. In *The Prokaryotes*, vol. 2, pp. 1302–1331. Edited by M. P. Starr, H. Stolp, H. G. Trüper, A. Balows & H. G. Schlegel. Berlin: Springer.
- Bowman, J. P. & McMeekin, T. A. (2005). Order X. *Alteromonadales* ord. nov. In *Bergey's Manual of Systematic Bacteriology*, 2nd edn, vol. 2, part B, pp. 443. Edited by D. J. Brenner, N. R. Krieg, J. T. Staley & G. M. Garrity. New York: Springer.
- Bruns, A., Rohde, M. & Berthe-Corti, L. (2001). *Muricauda ruestringensis* gen. nov., sp. nov., a facultatively anaerobic, appendaged bacterium from German North Sea intertidal sediment. *Int J Syst Evol Microbiol* 51, 1997–2006.
- Cohen-Bazire, G., Sistrom, W. R. & Stanier, R. Y. (1957). Kinetic studies of pigment synthesis by nonsulfur purple bacteria. *J Cell Comp Physiol* 49, 25–68.
- Cowan, S. T. & Steel, K. J. (1965). *Manual for the Identification of Medical Bacteria*. London: Cambridge University Press.
- Hosoya, S., Adachi, K. & Kasai, H. (2009). *Thalassomonas actiniarum* sp. nov. and *Thalassomonas haliotis* sp. nov., isolated from marine animals. *Int J Syst Evol Microbiol* 59, 686–690.
- Jean, W. D., Shieh, W. Y. & Liu, T. Y. (2006). *Thalassomonas agarivorans* sp. nov., a marine agarolytic bacterium isolated from shallow coastal water of An-Ping Harbour, Taiwan, and emended description of the genus *Thalassomonas*. *Int J Syst Evol Microbiol* 56, 1245–1250.
- Komagata, K. & Suzuki, K. (1987). Lipid and cell wall analysis in bacterial systematics. *Methods Microbiol* 19, 161–207.
- Lanyi, B. (1987). Classical and rapid identification methods for medically important bacteria. *Methods Microbiol* 19, 1–67.
- Macián, M. C., Ludwig, W., Schleifer, K. H., Garay, E. & Pujalte, M. J. (2001). *Thalassomonas viridans* gen. nov., sp. nov., a novel marine γ-proteobacterium. *Int J Syst Evol Microbiol* 51, 1283–1289.
- Sasser, M. (1990). *Identification of bacteria by gas chromatography of cellular fatty acids*, MIDI Technical Note 101. Newark, DE: MIDI Inc.
- Staley, J. T. (1968). *Prosthecomicrobium* and *Ancalomicrobium*: new prosthecate freshwater bacteria. *J Bacteriol* 95, 1921–1942.
- Tamaoka, J. & Komagata, K. (1984). Determination of DNA base composition by reverse-phase high-performance liquid chromatography. *FEMS Microbiol Lett* 25, 125–128.
- Teather, R. M. & Wood, P. J. (1982). Use of Congo red-polysaccharide interactions in enumeration and characterization of cellulolytic bacteria from the bovine rumen. *Appl Environ Microbiol* 43, 777–780.
- Thompson, F. L., Barash, Y., Sawabe, T., Sharon, G., Swings, J. & Rosenberg, E. (2006). *Thalassomonas loyana* sp. nov., a causative agent of the white plague-like disease of corals on the Eilat coral reef. *Int J Syst Evol Microbiol* 56, 365–368.
- Yi, H., Bae, K. S. & Chun, J. (2004). *Thalassomonas ganghwensis* sp. nov., isolated from tidal flat sediment. *Int J Syst Evol Microbiol* 54, 377–380.
- Yoon, J.-H., Kim, H., Kim, S.-B., Kim, H.-J., Kim, W. Y., Lee, S. T., Goodfellow, M. & Park, Y.-H. (1996). Identification of *Saccharomonospora* strains by the use of genomic DNA fragments and rRNA gene probes. *Int J Syst Bacteriol* 46, 502–505.
- Yoon, J.-H., Lee, S. T. & Park, Y.-H. (1998). Inter- and intraspecific phylogenetic analysis of the genus *Nocardioides* and related taxa based on 16S rDNA sequences. *Int J Syst Bacteriol* 48, 187–194.
- Yoon, J.-H., Kim, I.-G., Shin, D.-Y., Kang, K. H. & Park, Y.-H. (2003). *Microbulbifer salpaludis* sp. nov., a moderate halophile isolated from a Korean salt marsh. *Int J Syst Evol Microbiol* 53, 53–57.