Pantoea rodasii sp. nov., *Pantoea rwandensis* sp. nov. and *Pantoea wallisii* sp. nov., isolated from *Eucalyptus*

Carrie L. Brady,¹ Ilse Cleenwerck,² Lorinda van der Westhuizen,³ Stephanus N. Venter,³ Teresa A. Coutinho³ and Paul De Vos²

Correspondence

Teresa A. Coutinho
teresa.coutinho@fabi.up.ac.za

¹LM-UGent, Laboratory of Microbiology, Faculty of Sciences, Ghent University, Ghent, Belgium

²BCCM/LMG Bacteria Collection, Ghent University, Ghent, Belgium

³Department of Microbiology and Plant Pathology, Forestry and Agricultural Biotechnology Institute (FABI), University of Pretoria, Pretoria, South Africa

Several Gram-negative-staining, facultatively anaerobic bacterial isolates were obtained from *Eucalyptus* seedlings showing symptoms of bacterial blight and dieback in Colombia, Rwanda and South Africa. Partial 16S rRNA gene sequencing, together with partial *gyrB* sequencing, placed the isolates in the genus *Pantoea* and indicated that they constituted three novel species. Multilocus sequence analysis (MLSA) based on partial sequences of *gyrB*, *rpoB*, *infB* and *atpD* revealed *Pantoea dispersa*, *Pantoea eucrina* and *Pantoea cyripedii* as their closest phylogenetic relatives. DNA–DNA hybridization studies confirmed the classification of the new isolates as three novel species and phenotypic tests allowed them to be differentiated from their closest phylogenetic neighbours. The names *Pantoea rodasii* sp. nov. [type strain LMG 26273^T=BD 943^T (deposited with the Plant Pathogenic and Plant Protecting Bacteria Collection, South Africa)=BCC 581^T (deposited with the Bacterial Culture Collection, Forestry and Agricultural Institute, South Africa)], *Pantoea rwandensis* sp. nov. (type strain LMG 26275^T=BD 944^T=BCC 571^T) and *Pantoea wallisii* sp. nov. (type strain LMG 26277^T=BD 946^T=BCC 682^T) are proposed.

Pantoea ananatis has been reported as the causal agent of bacterial blight and dieback of *Eucalyptus* seedlings in South Africa. Young leaves present symptoms first, with leaf spots that become water-soaked and eventually form larger necrotic lesions. Trees either fail to survive or become multi-stemmed (Coutinho *et al.*, 2002). In the last decade, similar symptoms have been observed in nurseries and plantations in Uganda, Argentina and Uruguay. The bacteria isolated from these diseased trees were identified as belonging to three novel species of the genus *Pantoea*: *P. vagans*, *P. eucalypti* and *P. deleyi* (Brady *et al.*, 2009). It has been suggested that a complex of *Pantoea* species may be responsible for bacterial blight and dieback in Africa and South America (Coutinho *et al.*, 2011). *P. ananatis*, *P. vagans*, *P. eucalypti* and *P. deleyi*, have been isolated from a wide range of *Eucalyptus* species, hybrids and clones which is of concern for the forestry industry.

Abbreviation: MLSA, multilocus sequence analysis.

The GenBank/EMBL/DBJ accession numbers for the 16S rRNA, *atpD*, *gyrB*, *infB* and *rpoB* gene sequences are JF295053–JF295058, JF295013–JF295022, JF295023–JF295032, JF295033–JF295042 and JF295043–JF295052, respectively.

Two supplementary tables are available with the online version of this paper.

As part of an on-going isolation campaign in countries of Africa, South America and Asia, *Eucalyptus* seedlings are regularly examined for symptoms of bacterial blight and dieback. Bacterial isolates obtained from the diseased material are identified using a polyphasic approach based on Gram staining, oxidation–fermentation testing, partial 16S rRNA gene- and *gyrB*-sequencing. *P. ananatis* and *P. eucalypti* are regularly isolated in South Africa, while *P. vagans* and *P. dispersa* are more commonly isolated in Colombia and Thailand, respectively (Swart, 2009). In 2006/2007, bacteria were isolated from diseased *Eucalyptus* material in Colombia, Rwanda and South Africa which could not be assigned to any of the recognized species of the genus *Pantoea*. Sequencing of the *gyrB* gene placed these isolates in the genus *Pantoea*, and indicated that they constituted three novel species (Swart, 2009). In the present study, these isolates were further examined using a polyphasic approach to confirm that they constitute three novel species of the genus *Pantoea*.

Bacteria were isolated from diseased *Eucalyptus* material as previously described (Brady *et al.*, 2009). Reference strains were obtained from the BCCM/LMG Bacteria Collection (<http://bccm.belspo.be>) and recovered on tryptic soy agar according to the provider's instructions. A list of strains

used in this study is available in Table S1 in IJSEM Online. Genomic DNA was extracted using the alkali method (Niemann *et al.*, 1997) and stored at -20°C .

Almost complete 16S rRNA gene sequences (1346 bp) were determined for two strains from each proposed novel species using the primers and conditions as previously described (Coenye *et al.*, 1999). Consensus sequences were aligned using the CLUSTAL W application in BioEdit version 7.0.9.0 (Hall, 1999) and the overhangs were trimmed. Phylogenetic trees were constructed using the maximum-parsimony and neighbour-joining methods in MEGA 5.0 (Tamura *et al.*, 2011) and PAUP 4.0b10 (Swofford, 2000), respectively. The reliability of the clusters was evaluated by bootstrap analysis with 1000 replicates. As the topology of the resulting phylogenetic trees was similar, only the maximum-parsimony tree is shown.

In the 16S rRNA gene maximum-parsimony tree (Fig. 1), the novel species formed three definite clusters corresponding to the country of isolation, each with 100% bootstrap support. The isolates from Colombia and Rwanda were more closely related to each other than to the South African isolates and formed a clade on a separate branch, while the South African isolates clustered with *P. dispersa* and *P. eucrina*. Several 'core' species of the genus *Pantoea* formed a well-supported clade with the type strain of *P. agglomerans* while the remainder clustered at a lower level. It has been demonstrated previously that the genus *Pantoea* is polyphyletic (Brady *et al.*, 2010b), making it increasingly difficult to allocate novel species to this genus based solely on 16S rRNA gene sequencing. Numerous genera within the family *Enterobacteriaceae* are polyphyletic when analysis is based on 16S rRNA gene sequences alone, and whether this gene is an appropriate choice to construct phylogenies of closely related bacterial taxa has been questioned (Naum *et al.*, 2008).

The 16S rRNA gene sequence pairwise similarity obtained was >99% amongst the Colombian isolates, >99.4% amongst the isolates from Rwanda and >99.8% between the South African isolates. The Colombian, Rwandan and South African isolates displayed more than 97.0% 16S rRNA gene sequence pairwise similarity amongst each other and with various species of the family *Enterobacteriaceae*. Based on 16S rRNA gene sequencing, the closest phylogenetic relatives of the Colombian and Rwandan isolates were *P. septica*, *P. eucrina*, *P. dispersa*, *P. cyripedii*, *Erwinia aphidicola*, *Kluyvera intermedia*, *Buttiauxella agrestis* and *Enterobacter ludwigii*. The isolates from South Africa were most closely related to *P. dispersa* and *P. eucrina*.

Multilocus Sequence Analysis (MLSA) based on partial sequences of *gyrB*, *rpoB*, *infB* and *atpD*, four protein-encoding genes, was performed on all strains belonging to the three novel species as described previously (Brady *et al.*, 2008). Consensus sequences were aligned using the CLUSTAL W application in BioEdit version 7.0.9.0 (Hall, 1999) and the overhangs were trimmed. The best-fit evolutionary model was selected by Modeltest 3.7 (Posada & Crandall,

1998) and maximum-likelihood and neighbour-joining trees were constructed in PHYML (Guindon & Gascuel, 2003) and PAUP 4.0b10 (Swofford, 2000), respectively, using the parameters determined by Modeltest. Bootstrap analysis with 1000 replicates was performed on each of the trees to gauge the reliability of the clusters. The topology of both trees was similar and therefore only the maximum-likelihood tree is shown.

The peptide sequences were also determined for each gene and a concatenated peptide sequence tree was constructed in PHYML using the parameters described previously (Brady *et al.*, 2008). In the maximum-likelihood tree based on concatenated sequences of the four genes (Fig. 2), the isolates from Colombia, Rwanda and South Africa formed three separate clusters (with 100% bootstrap values) in the strongly supported clade containing all recognized species of the genus *Pantoea*. The same topology was observed in the concatenated peptide sequence tree (data not shown) providing support at the protein level for the delineation of these isolates as three novel species of the genus *Pantoea*. The three novel species were found to share 20 of the 23 *atpD* signature nucleotides that can be used to differentiate species of the genus *Pantoea* from closely related species of the genera *Tatumella* and *Erwinia* (Brady *et al.*, 2010a). The MLSA data therefore placed the isolates in the genus *Pantoea* and suggested that they belong to three novel species. As observed in the 16S rRNA gene phylogenetic tree, the isolates from Colombia and Rwanda were more closely related to each other than to those from South Africa. Based on the MLSA data, the closest phylogenetic relatives of the three novel species were *P. eucrina*, *P. dispersa* and *P. cyripedii*.

Two isolates were selected from each novel species for DNA–DNA hybridization experiments. Isolates from Colombia and Rwanda were hybridized with each other, and a representative isolate from each proposed novel species was hybridized with the type strains of *P. septica*, *P. eucrina*, *P. dispersa*, *P. cyripedii*, *Erwinia aphidicola*, *K. intermedia*, *B. agrestis* and *Enterobacter ludwigii*. The isolates from South Africa were also hybridized amongst each other, and a representative isolate was hybridized with the type strains of *P. dispersa*, *P. eucrina* and *P. cyripedii*. Large-scale DNA extraction was performed on the strains using a modified version (Cleenwerck *et al.*, 2002) of the method described by Wilson (1987). DNA–DNA hybridizations (four replications) were performed at 45°C using the microplate method (Ezaki *et al.*, 1989) with some modifications (Cleenwerck *et al.*, 2002). Reciprocal reactions ($A \times B$ and $B \times A$) were performed for each DNA pair from all strains and their variation was within the limits for this method (Goris *et al.*, 1998). Isolates of the novel species exhibited >98% DNA–DNA relatedness when hybridized against each other, but <44% DNA–DNA relatedness was observed between the Colombian and Rwandan isolates, and <35% between these isolates and the type strains of the other species of the genus. The isolates from South Africa displayed <40% DNA–DNA relatedness to *P. dispersa*, *P.*

Fig. 1. Maximum-parsimony tree based on almost-complete 16S rRNA gene sequences of members of the genus *Pantoea* and phylogenetically related species. Bootstrap values after 1000 replicates are expressed as percentages. Species belonging to the genus *Tatumella* were included as outgroups. Bar, 5 substitutions per site.

eucrina and *P. cyripedii*, their closest phylogenetic relatives. The hybridization results confirmed that the isolates constituted three novel species and are summarized in Table S2 in IJSEM Online.

The DNA G+C contents of the novel species was measured by HPLC (Mesbah *et al.*, 1989) and were as follows: LMG 26273^T and LMG 26274, 53.2 and 53.0 mol%; LMG 26275^T and LMG 26276, 51.9 and 52.0 mol% and LMG 26277^T and LMG 26278, 55.5 and 55.6 mol%. These values fell within the G+C content range of the recently emended description of the genus *Pantoea* (Brady *et al.*, 2010b).

API 20E, API 50CHB/E (bioMérieux) and GN2 MicroPlate (Biolog) tests were performed, according to the manufacturers' instructions, on the isolates from Colombia, Rwanda and South Africa. Cell suspensions were prepared from cultures grown on tryptic soy agar at 28 °C for 12 h. API and Biolog tests were read after 24 and 48 h of incubation. Data were compared with those previously published for species of the genus *Pantoea* (Brady *et al.*, 2009, 2010a, b) and generated under the same conditions. The novel species were found to share all phenotypic traits characteristic of the genus *Pantoea* (Brady *et al.*, 2010b; Grimont & Grimont, 2005; Mergaert *et al.*, 1993). The results are listed in the species descriptions below. The

Fig. 2. Maximum-likelihood tree based on concatenated housekeeping gene sequences of strains of the genus *Pantoea*. Bootstrap values after 1000 replicates are expressed as percentages. Major clades also supported by the single gene phylogenies are indicated by *g* (*gyrB*), *r* (*rpoB*), *i* (*infB*) and *a* (*atpD*) in parentheses. *Cronobacter sakazakii* ATCC BAA-894 was included as an outgroup. Gene sequences for *C. sakazakii* were obtained from <http://www.ncbi.nlm.nih.gov>. Bar, 0.1 substitutions per site.

three novel species could be differentiated from each other by their reactions to dulcitol, sucrose, adonitol, fucose, psicose and serine. The most useful characteristics for differentiating the novel species from each other and their closest phylogenetic relatives are listed in Table 1.

The whole-cell fatty acid methyl ester composition was determined for two isolates from each novel species as well as for the type strain of the type species of the genus, *P. agglomerans*, using the Microbial Identification System, Sherlock version 3.10 (MIDI) and the TSBA50 identification library version 5.0 according to the previously published protocol (Mergaert *et al.*, 1993). A gas chromatograph (6890N; Agilent Technologies) was used for separation of the fatty acid methyl esters. Cells were harvested from cultures grown on trypticase soy agar (BBL 11768) for 24 h at 28 °C. The novel species and *P. agglomerans* displayed similar fatty acid compositions, corresponding with those available in literature (Mergaert *et al.*, 1993, 1999). The major fatty acids included C_{12:0}, C_{14:0}, C_{16:0}, C_{17:0} cyclo, C_{18:1}ω7c, and summed features 2 (iso-C_{16:1} and/or C_{14:0} 3-OH) and 3 (C_{16:1}ω7c and/or iso-C_{15:0} 2-OH). The fatty acid profiles for the novel species are presented in the species descriptions below.

Based on the genotypic and phenotypic data generated in this study, it is clear that the isolates from the diseased *Eucalyptus* seedlings in Colombia, Rwanda and South Africa constitute three novel species in the genus *Pantoea*. Therefore we propose to classify them as *Pantoea rodasii* sp. nov. (isolated from Colombia, type strain LMG 26273^T=BD 943^T), *Pantoea rwandensis* sp. nov. (isolated from Rwanda, type strain LMG 26275^T=BD 944^T) and *Pantoea wallisii* sp. nov. (isolated from South Africa, type strain LMG 26277^T=BD 946^T).

Description of *Pantoea rodasii* sp. nov.

Pantoea rodasii (ro.da'si.i. N.L. masc. gen. n. *rodasii* of Rodas, named after Carlos Rodas for his contribution to forest pathology in Colombia).

Cells are Gram-negative-staining, short rods (1 × 1.5–3 µm) occurring singly or in pairs, weakly motile and non-spore-forming. Colonies are round, smooth and convex with entire margins on tryptone soy agar and light beige in colour after incubation of 24 h at 28 °C. Facultatively anaerobic, oxidase-negative and catalase-positive. Acetoin and β-galactosidase are produced, but H₂S, urease and indole are not produced and citrate is not utilized. Tests for arginine dihydrolase, lysine decarboxylase and ornithine decarboxylase are all negative. Acid is produced from the

fermentation of glycerol, L-arabinose, D-ribose, D-xylose, D-adonitol, D-galactose, D-glucose, D-fructose, D-mannose, L-rhamnose, inositol, D-mannitol, N-acetylglucosamine, arbutin, aesculin ferric citrate, salicin, cellobiose, melibiose, sucrose, trehalose, gentiobiose, D-fucose and D-arabitol (API 50CHB/E). The following carbon sources are utilized at 28 °C by the majority of strains tested including the type strain, after 24 h incubation: Tweens 40 and 80, N-acetyl-D-glucosamine, adonitol, L-arabinose, D-arabitol, cellobiose, erythritol, D-fructose, D-galactose, gentiobiose, D-glucose, inositol, lactose, D-mannitol, D-mannose, melibiose, methyl β-D-glucoside, D-psicose, L-rhamnose, D-sorbitol, sucrose, trehalose, pyruvic acid methyl ester, succinic acid mono-methyl ester, acetic acid, formic acid, D-galactonic acid lactone, D-galacturonic acid, D-gluconic acid, D-glucosaminic acid, D-glucuronic acid, DL-lactic acid, quinic acid, D-saccharic acid, succinic acid, bromosuccinic acid, glucuronamide, D-alanine, L-alanine, L-asparagine, L-aspartic acid, L-glutamic acid, glycyl L-aspartic acid, glycyl L-glutamic acid, L-histidine, L-proline, D-serine, L-serine, urocanic acid, inosine, uridine, thymidine, glycerol, DL-α-glycerol phosphate, α-D-glucose 1-phosphate, D-glucose 6-phosphate (Biolog). Strains display the following fatty acid profile: C_{12:0} (4.5%), C_{14:0} (6.7%), C_{16:0} (27.4%), C_{17:0} cyclo (8.8%), C_{18:1}ω7c (11.0%), summed feature 2 (iso-C_{16:1} and/or C_{14:0} 3-OH) (13.9%) and summed feature 3 (C_{16:1}ω7c and/or iso-C_{15:0} 2-OH) (23.8%).

The type strain is LMG 26273^T [=BD 943^T (deposited with the Plant Pathogenic and Plant Protecting Bacteria Collection, South Africa)=BCC 581^T (deposited with the Bacterial Culture Collection, Forestry and Agricultural Biotechnology Institute, South Africa)]. The DNA G+C content of the type strain is 53.2 mol%. Strains belonging to this species were isolated from lesions on *Eucalyptus* leaves exhibiting symptoms of bacterial blight and dieback in Colombia.

Description of *Pantoea rwandensis* sp. nov.

Pantoea rwandensis (rwan.den'sis. N.L. fem. adj. *rwandensis* of or belonging to Rwanda, referring to the country of isolation).

Cells are Gram-negative-staining, short rods (1 × 2–3 µm) occurring singly or in pairs, non-motile and non-spore-forming. Colonies are round, smooth and convex with entire margins on tryptone soy agar and beige in colour after incubation of 24 h at 28 °C. Facultatively anaerobic, oxidase-negative and catalase-positive. β-Galactosidase is produced, but H₂S, urease and indole are not produced

Table 1. Phenotypic characteristics differentiating the novel species from each other, from the type species of the genus *Pantoea* and other species of the genus associated with bacterial blight and dieback of *Eucalyptus*

Species: 1, *Pantoea rodasii* sp. nov. ($n=4$); 2, *Pantoea rwandensis* sp. nov. ($n=4$); 3, *Pantoea wallisii* sp. nov. ($n=5$); 4, *P. agglomerans*; 5, *P. ananatis*; 6, *P. deleyi*; 7, *P. dispersa*; 8, *P. eucalypti*; 9, *P. eucrinea*; 10, *P. septica*; 11, *P. vagans*; 12, *P. cypridii*; 13, *P. gaviniae*; 14, *P. calida*. n , Number of strains tested; +, 90–100 % strains positive in 1–2 days; (+), 90–100 % strains positive in 1–4 days; –, negative; d, 11–89 % strains positive in 1–4 days; (d), 11–89 % strains positive in 3–4 days; ND, not determined. Data were taken from the following sources: taxa 1, 2, 3 (this study), taxa 4–12 (Brady *et al.*, 2009, 2010a, b; Grimont & Grimont, 2005), taxa 13, 14 (Popp *et al.*, 2010). Only data generated under the same conditions are listed.

Characteristic	1	2	3	4	5	6	7	8	9	10	11	12	13	14
Acid from:														
Glycerol	+	+	+	(d)	+	+	(d)	d	–	d	d	+	+	+
Dulcitol	–	+	+	–	–	–	–	–	–	–	–	–	–	–
Lactose	–	–	–	d	+	–	+	(+)	d	d	(+)	–	+	+
Sucrose	+	–	–	+	+	+	+	+	+	d	+	+	+	ND
Raffinose	–	–	–	(d)	+	–	–	–	–	–	–	–	+	ND
Utilization of:														
Adonitol	+	–	–	–	–	–	–	–	+	–	–	–	–	–
<i>i</i> -Erythritol	+	+	–	–	–	–	+	–	d	–	–	–	–	–
L-Fucose	–	+	+	–	–	–	–	–	–	d	–	–	–	–
Lactose	d	–	–	–	+	–	–	+	–	d	–	+	+	+
Lactulose	–	–	–	–	+	–	–	–	–	d	–	–	+	+
Melibiose	d	+	–	–	+	–	–	–	–	d	–	+	+	+
D-Psicose	+	+	–	+	+	+	(d)	(d)	ND	ND	–	+	ND	ND
Sucrose	+	–	–	(+)	+	+	+	+	+	d	+	+	+	+
Xylitol	–	–	–	–	–	–	–	–	+	d	–	–	–	–
Quinic acid	+	+	d	–	(+)	–	–	–	–	–	–	+	–	–
D-Serine	d	d	–	–	–	–	–	–	ND	ND	–	(+)	ND	ND

and citrate is not utilized. Negative results in tests for arginine dihydrolase, lysine decarboxylase and ornithine decarboxylase. Acid is produced from the fermentation of glycerol, D-arabinose, L-arabinose, D-ribose, D-xylose, D-galactose, D-glucose, D-fructose, D-mannose, L-rhamnose, dulcitol, inositol, D-mannitol, *N*-acetylglucosamine, arbutin, aesculin ferric citrate, salicin, cellobiose, melibiose, trehalose, gentiobiose, D-fucose, L-fucose and D-arabitol (API 50CHB/E). The following carbon sources are utilized at 28 °C by the majority of strains tested including the type strain, after 24 h incubation: Tweens 40 and 80, *N*-acetyl-D-glucosamine, L-arabinose, D-arabitol, cellobiose, erythritol, D-fructose, L-fucose, D-galactose, gentiobiose, D-glucose, inositol, D-mannitol, D-mannose, melibiose, methyl β -D-glucoside, D-psicose, L-rhamnose, trehalose, pyruvic acid methyl ester, succinic acid monomethyl ester, acetic acid, formic acid, D-galactonic acid lactone, D-galacturonic acid, D-gluconic acid, D-glucosaminic acid, D-glucuronic acid, DL-lactic acid, quinic acid, D-saccharic acid, succinic acid, bromosuccinic acid, glucuronamide, L-alanine, L-alanyl glycine, L-asparagine, L-aspartic acid, L-glutamic acid, glycyl L-aspartic acid, glycyl L-glutamic acid, L-proline, D-serine, L-serine, inosine, uridine, thymidine, glycerol, DL- α -glycerol phosphate, α -D-glucose 1-phosphate, D-glucose 6-phosphate (Biolog). Strains display the following fatty acid profile: C_{12:0} (4.2 %), C_{14:0} (6.9 %), C_{16:0} (26.1 %), C_{17:0} cyclo (7.1 %), C_{18:1} ω 7c (11.8 %), summed feature 2 (iso-C_{16:1} and/or C_{14:0} 3-OH) (14.3 %) and summed feature 3 (C_{16:1} ω 7c and/or iso-C_{15:0} 2-OH) (26.5 %).

The type strain is LMG 26275^T [=BD 944^T (deposited with the Plant Pathogenic and Plant Protecting Bacteria Collection, South Africa)=BCC 571^T (deposited with the Bacterial Culture Collection, Forestry and Agricultural Biotechnology Institute, South Africa)]. The DNA G+C content of the type strain is 51.2 mol%. Strains belonging to this species were isolated from lesions on *Eucalyptus* leaves exhibiting symptoms of bacterial blight and dieback in Rwanda.

Description of *Pantoea wallisii* sp. nov.

Pantoea wallisii (wall.li'si.i. N.L. masc. gen. n. *wallisii* of Wallis, named after F. M. Wallis for his contribution to the field of phytobacteriology in South Africa).

Cells are Gram-negative-staining, short rods (1 × 1–2.5 μ m) occurring singly or in pairs, motile and non-spore-forming. Colonies are round, smooth and convex with entire margins on tryptone soy agar and pale yellow in colour after incubation of 24 h at 28 °C. Facultatively anaerobic, oxidase-negative and catalase-positive. β -Galactosidase is produced, but H₂S, urease and indole are not produced and citrate is utilized. Negative result in tests for arginine dihydrolase, lysine decarboxylase and ornithine decarboxylase. Acid is produced from the fermentation of glycerol, D-arabinose, L-arabinose, D-ribose, D-xylose, D-galactose, D-glucose, D-fructose, D-mannose, L-rhamnose, dulcitol, inositol, D-mannitol, *N*-acetylglucosamine, arbutin, salicin, cellobiose, maltose, melibiose, trehalose,

gentiobiose, D-lyxose, D-fucose, L-fucose and D-arabitol (API 50CHB/E). The following carbon sources are utilized at 28 °C by the majority of strains tested including the type strain, after 24 h incubation: dextrin, Tweens 40 and 80, N-acetyl-D-glucosamine, L-arabinose, D-arabitol, cellobiose, D-fructose, L-fucose, D-galactose, gentiobiose, D-glucose, inositol, maltose, D-mannitol, D-mannose, methyl β -D-glucoside, L-rhamnose, trehalose, pyruvic acid methyl ester, succinic acid monomethyl ester, acetic acid, *cis*-aconitic acid, citric acid, formic acid, D-galactonic acid lactone, D-galacturonic acid, D-gluconic acid, D-glucosaminic acid, D-glucuronic acid, DL-lactic acid, quinic acid, D-saccharic acid, succinic acid, bromosuccinic acid, glucuronamide, D-alanine, L-alanine, L-alanyl glycine, L-asparagine, L-aspartic acid, L-glutamic acid, glycyl L-aspartic acid, glycyl L-glutamic acid, L-serine, inosine, uridine, thymidine, glycerol, DL- α -glycerol phosphate, α -D-glucose 1-phosphate, D-glucose 6-phosphate (Biolog). Strains display the following fatty acid profile: C_{12:0} (4.2 %), C_{14:0} (6.9 %), C_{16:0} (26.1 %), C_{17:0} cyclo (7.1 %), C_{18:1} ω 7c (11.8 %), summed feature 2 (iso-C_{16:1} and/or C_{14:0} 3-OH) (14.3 %) and summed feature 3 (C_{16:1} ω 7c and/or iso-C_{15:0} 2-OH) (26.5 %).

The type strain is LMG 26277^T [=BD 946^T (deposited with the Plant Pathogenic and Plant Protecting Bacteria Collection, South Africa)=BCC 682^T (deposited with the Bacterial Culture Collection, Forestry and Agricultural Biotechnology Institute, South Africa)]. The DNA G+C content of the type strain is 55.5 mol%. Strains belonging to this species were isolated from lesions on *Eucalyptus* leaves exhibiting symptoms of bacterial blight and dieback in South Africa.

Acknowledgements

The authors wish to acknowledge Katrien Engelbeen and Fati Thobjane for technical assistance, and Professor J. J. van der Walt and Dr J. P. Euzéby for assistance with the etymology of the species names. We also thank Mike Wingfield and Pieter de Maayer for collecting *Eucalyptus* material and Emma Steenkamp for discussions relating to the phylogenetic analyses. The BCCM/LMG Bacteria Collection is supported by the Federal Public Planning Service – Science Policy, Belgium. C. Brady is the beneficiary of a fellowship granted by the Federal Science Policy Office, Belgium.

References

- Brady, C. L., Cleenwerck, I., Venter, S. N., Vancanneyt, M., Swings, J. & Coutinho, T. A. (2008). Phylogeny and identification of *Pantoea* species associated with plants, humans and the natural environment based on multilocus sequence analysis (MLSA). *Syst Appl Microbiol* **31**, 447–460.
- Brady, C. L., Venter, S. N., Cleenwerck, I., Engelbeen, K., Vancanneyt, M., Swings, J. & Coutinho, T. A. (2009). *Pantoea vagans* sp. nov., *Pantoea eucalypti* sp. nov., *Pantoea deleyi* sp. nov. and *Pantoea anthophila* sp. nov. *Int J Syst Evol Microbiol* **59**, 2339–2345.
- Brady, C. L., Venter, S. N., Cleenwerck, I., Vandemeulebroecke, K., De Vos, P. & Coutinho, T. A. (2010a). Transfer of *Pantoea citrea*, *Pantoea punctata* and *Pantoea terreia* to the genus *Tatumella* emend. as *Tatumella citrea* comb. nov., *Tatumella punctata* comb. nov. and *Tatumella terreia* comb. nov. and description of *Tatumella morbirosee* sp. nov. *Int J Syst Evol Microbiol* **60**, 484–494.
- Brady, C. L., Cleenwerck, I., Venter, S. N., Engelbeen, K., De Vos, P. & Coutinho, T. A. (2010b). Emended description of the genus *Pantoea*, description of four species from human clinical samples, *Pantoea septica* sp. nov., *Pantoea eucrina* sp. nov., *Pantoea brenneri* sp. nov. and *Pantoea conspicua* sp. nov., and transfer of *Pectobacterium cypripedii* (Hori 1911) Brenner et al. 1973 emend. Hauben et al. 1998 to the genus *Pantoea cypripedii* comb. nov. *Int J Syst Evol Microbiol* **60**, 2430–2440.
- Cleenwerck, I., Vandemeulebroecke, K., Janssens, D. & Swings, J. (2002). Re-examination of the genus *Acetobacter*, with descriptions of *Acetobacter cerevisiae* sp. nov. and *Acetobacter malorum* sp. nov. *Int J Syst Evol Microbiol* **52**, 1551–1558.
- Coenye, T., Falsen, E., Vancanneyt, M., Hoste, B., Govan, J. R. W., Kersters, K. & Vandamme, P. (1999). Classification of *Alcaligenes faecalis*-like isolates from the environment and human clinical samples as *Ralstonia gilardii* sp. nov. *Int J Syst Bacteriol* **49**, 405–413.
- Coutinho, T. A., Preisig, O., Mergaert, J., Cnockaert, M. C., Riedel, K. H., Swings, J. & Wingfield, M. J. (2002). Bacterial blight and dieback of *Eucalyptus* species, hybrids, and clones in South Africa. *Plant Dis* **86**, 20–25.
- Coutinho, T. A., Venter, S. N., Roux, J., Zhou, X. D. & Wingfield, M. J. (2011). Bacterial diseases of *Eucalyptus*. Proceedings of the ACP APPS 2011 Congress, Darwin, Australia 27–29 April 2011.
- Ezaki, T., Hashimoto, Y. & Yabuuchi, E. (1989). Fluorometric deoxyribonucleic acid-deoxyribonucleic acid hybridization in micro-dilution wells as an alternative to membrane filter hybridization in which radioisotopes are used to determine genetic relatedness among bacterial strains. *Int J Syst Bacteriol* **39**, 224–229.
- Goris, J., Suzuki, K.-I., De Vos, P., Nakase, T. & Kersters, K. (1998). Evaluation of a microplate DNA-DNA hybridization method compared with the initial renaturation method. *Can J Microbiol* **44**, 1148–1153.
- Grimont, P. A. D. & Grimont, F. (2005). Genus: *Pantoea*. In *Bergey's Manual of Systematic Bacteriology, Volume Two, The Proteobacteria, Part B, The Gammaproteobacteria*, 2nd edn, pp. 713–720. Edited by D. J. Brenner, N. R. Krieg & J. T. Staley. New York: Springer.
- Guindon, S. & Gascuel, O. (2003). A simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood. *Syst Biol* **52**, 696–704.
- Hall, T. A. (1999). BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symp Ser* **41**, 95–98.
- Mergaert, J., Verdonck, L. & Kersters, K. (1993). Transfer of *Erwinia ananas* (synonym, *Erwinia uredovora*) and *Erwinia stewartii* to the genus *Pantoea* emend. as *Pantoea ananas* (Serrano 1928) comb. nov. and *Pantoea stewartii* (Smith 1898) comb. nov., respectively, and description of *Pantoea stewartii* subsp. *indologenes* subsp. nov. *Int J Syst Bacteriol* **43**, 162–173.
- Mergaert, J., Hauben, L., Cnockaert, M. C. & Swings, J. (1999). Reclassification of non-pigmented *Erwinia herbicola* strains from trees as *Erwinia billingiae* sp. nov. *Int J Syst Bacteriol* **49**, 377–383.
- Mesbah, M., Premachandran, U. & Whitman, W. B. (1989). Precise measurement of the G+C content of deoxyribonucleic acid by high-performance liquid chromatography. *Int J Syst Bacteriol* **39**, 159–167.
- Naum, M., Brown, E. W. & Mason-Gamer, R. J. (2008). Is 16S rDNA a reliable phylogenetic marker to characterize relationships below the family level in the *Enterobacteriaceae*? *J Mol Evol* **66**, 630–642.
- Niemann, S., Pühler, A., Tichy, H.-V., Simon, R. & Selbitschka, W. (1997). Evaluation of the resolving power of three different DNA

fingerprinting methods to discriminate among isolates of a natural *Rhizobium meliloti* population. *J Appl Microbiol* **82**, 477–484.

Popp, A., Cleenwerck, I., Iversen, C., De Vos, P. & Stephan, R. (2010). *Pantoea gaviniae* sp. nov. and *Pantoea calida* sp. nov., isolated from infant formula and an infant formula production environment. *Int J Syst Evol Microbiol* **60**, 2786–2792.

Posada, D. & Crandall, K. A. (1998). MODELTEST: testing the model of DNA substitution. *Bioinformatics* **14**, 817–818.

Swart, L. (2009). *Pantoea* and *Xanthomonas* species associated with bacterial blight and die-back of *Eucalyptus*. MSc thesis. University of Pretoria, Pretoria, South Africa.

Swofford, D. L. (2000). PAUP*: Phylogenetic analysis using parsimony (and other methods) Version 4.0. Sunderland, MA: Sinauer Associates.

Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M. & Kumar, S. (2011). MEGA5: Molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol Biol Evol* **28**, 2731–2739.

Wilson, K. (1987). Preparation of genomic DNA from bacteria. In *Current Protocols in Molecular Biology*, pp. 2.4.1–2.4.5. Edited by F. M. Ausubel, R. Brent, R. E. Kingston, D. D. Moore, J. G. Seidman, J. A. Smith & K. Struhl. New York: Green Publishing & Wiley-Interscience.